

Fig S1

Smith et al, Kidney Medicine, "Estimated GFR Trajectories in Pediatric and Adult Nephrotic Syndrome: Results From the Nephrotic Syndrome Study Network (NEPTUNE)"

Patients enrolled in NEPTUNE

N=1033

Patients deemed
ineligible after
enrollment
N=312

Eligible patients
N=721

Patients excluded:

- Less than one year of follow-up (145)
- <3 eGFR measurements (n= 6)
- Missing disease diagnosis (n=106)
- Missing baseline eGFR (n=11)

Analyzable patients
N=453




Fig S3




Table S1: Rates of progression to ESKD or 40% decline in eGFR across categories of PNL

PNL Category	Years from Baseline					Overall
	1	2	3	4	5	
Rate of ESRD or 40% decline in eGFR per 1000 person months						
0-10%	5.7	6.7	6.3	5.5	5.5	6.6
10-50%	6.8	7.7	6.9	6.3	9.2	9.2
50-100%	10.3	11.4	11.3	14.7	9.6	13.7
eGFR slope per month						
0-10%	-0.20	-0.26	-0.23	-0.17	-0.16	-0.24
10-50%	-0.28	-0.35	-0.33	-0.25	-0.25	-0.43
50-100%	-0.44	-0.45	-0.37	-0.13	-0.10	-0.53