



1 Article

2 Plasmonic Metasensors based on 2D hybrid 3 atomically thin perovskite nanomaterials

4 Shuwen Zeng^{1,2*†}, Guozhen Liang^{2†}, Alexandre Gheno¹, Sylvain Vedraïne¹, Bernard Ratier¹,
5 Ho-Pui Ho³, Nanfang Yu^{2*}

6 ¹ XLIM Research Institute, UMR 7252 CNRS/University of Limoges, 123 Avenue Albert Thomas, Limoges,
7 France;

8 ² Department of Applied Physics and Applied Mathematics, Columbia University, New York, USA

9 ³ Department of Biomedical Engineering, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong;

10 † These authors contributed equally to this work.


11 * Correspondence: shuwen.zeng@unilim.fr (S.Z.), ny2214@columbia.edu (N.Y.)

12 Received: date; Accepted: date; Published: date

13

14

15




16

17

18 **Figure S1.** Experimental measurement results showing the broad absorption spectra of $\text{CH}_3\text{NH}_3\text{PbI}_3$ (MAPbI₃)
19 perovskites.

20

21




22

23

24

Figure S2. SPR curve widths tuned by the groove depth of 2D Perovskite-based metasurface structure.



25

26

Figure S3. SPR curves with Reflectivity tuned by the groove width of the metasurface structure with 2D perovskite layers, corresponding to the zoom-in Figure 10b.

27

28

29

30


31

32

33

34

35



36

37

38


39

40

41


42

Figure S4. Experimental and simulation results based on three-layer graphene/gold metasurfaces. Theoretical curves of a) reflectance with respect to various incident angles in the air, c) differential phase and e) differential GH shift between p-polarized light and s-polarized light of glycerol solutions with different concentrations. Experimental curves of b) reflectance, d) differential phase and f) differential GH shift. The resonance angle is 44.10° in good agreement with the theoretical one. The phase and GH shift shows high sensitivity as $22,842 \text{ deg/RIU}$ and $7,500 \mu\text{m/RIU}$ respectively.



43
44
45
46
47

Figure S5. (a) and (b) Experimental and simulation data, respectively, of GH shift in a device based on a bilayer WS_2 .



48
49
50
51
52

Figure S6. The averaged GH shift of BSA molecules with molar concentration from 1aM to 1mM based on bilayer WS_2 /gold metasurfaces. The insert curve gives the binding trace of BSA molecules during a short time.