

Table S1. Univariate linear regression predicting criticism and emotional overinvolvement (FQ parents) - slopes presented only.

Predictor	FQ-CC b(SE)	t(p)	FQ-EOI b(SE)	t(p)
Female sex (parent) ¹	-1.921(.191)	-1.613 (.109)	-1.826 (.092)	-1.672 (.097)
Age parent	0.043 (.099)	0.433 (.666)	0.058 (.090)	0.640 (.523)
GHQ total score	0.496 (.117)	4.239 (<.001)	0.852 (.621)	9.541 (<.001)
EDSIS total score	0.220 (.026)	8.458 (<.001)	0.250 (.021)	12.171 (<.001)
CASK total score	-0.204(.029)	-7.145 (<.001)	-0.183 (.497)	-6.927 (<.001)
BDI total score	0.202 (.059)	3.414 (.001)	0.393 (.032)	8.568 (<.001)
STAI state score	0.133 (.039)	3.400 (.001)	0.237 (.032)	7.513 (<.001)
STAI trait score	0.154 (.044)	3.472 (.001)	0.261 (.037)	7.115 (<.001)
URICA precontemp.	-0.125(.099)	-1.253 (.212)	-0.270 (.089)	-3.036 (.003)
URICA contemp.	0.213 (.078)	2.721 (.007)	0.340 (.067)	5.057 (<.001)
URICA action	0.242 (.087)	2.775 (.006)	0.274 (.078)	3.498 (.001)
Time spent caregiving ²	0.538 (.995)	0.540 (.590)	0.739 (.940)	0.786 (.433)
Age patient	0.233 (.275)	0.849 (.397)	0.419 (.250)	1.675 (.096)
Treatment type ³	1.993 (.914)	2.180 (.031)	0.852 (.850)	1.003 (.317)
ED duration	0.062 (.034)	1.807 (.073)	0.038 (.032)	1.207 (.230)
BMI Perc.cat ⁴	1.220 (.910)	1.341 (.182)	0.852 (.833)	1.022 (.308)
EDE total score	-0.386(.318)	-1.212 (.227)	0.068 (.847)	0.080 (.936)
EDI total score	0.007 (.011)	0.674 (.501)	0.014 (.010)	1.477 (.142)

¹ 1 = female, 2 = male; ² 1 = < 3h/day, 2 = ≥ 3h/day; ³ 1 = inpatient, 2 = outpatient, ⁴ 1 = ≤ 1st percentile, 2 = > 1st percentile; Abbreviations: BDI Beck Depression Inventory, CASK Caregiver Skills Scale, contemp. contemplation, EDE Eating Disorder Examination Interview, EDI-2 Eating Disorder Inventory-2, EDSIS Eating Disorders Symptoms Impact Scale, FQ-CC Family Questionnaire Criticism Score, FQ-EOI Family Questionnaire Emotional Overinvolvement Score, GHQ General Health Questionnaire, Perc.cat Percentile category, STAI State/Trait Anxiety Inventory, URICA University of Rhode Island Change Assessment Scale.

Table S2. Univariate linear regression predicting criticism and emotional overinvolvement (FEICS, patients) - slopes presented only.

Predictor	FEICS-CC b(SE)	t(p)	FEICS-EOI b(SE)	t(p)
Age parent	0.102 (.074)	1.381 (.170)	0.120 (.073)	1.646 (.102)
GHQ total score	0.043 (.096)	0.450 (.654)	0.149 (.093)	1.596 (.113)
EDSIS total score	0.039 (.025)	1.563 (.120)	0.039 (.024)	1.584 (.115)
CASK total score	-0.047 (.026)	-1.809 (.073)	-0.005 (.026)	-0.196 (.845)
BDI total score	0.011 (.049)	0.223 (.824)	0.073 (.049)	1.510 (.134)
STAI state score	-0.048 (.031)	-1.562 (.121)	0.036 (.031)	1.136 (.258)
STAI trait score	0.030 (.035)	0.859 (.392)	0.030 (.036)	0.852 (.396)
URICA precontemp.	0.035 (.079)	0.451 (.653)	-0.149 (.075)	-1.992 (.048)
URICA contemp.	-0.082 (.062)	-1.323 (.188)	0.090 (.060)	1.494 (.138)
URICA action	-0.052 (.070)	-0.750 (.454)	0.106 (.067)	1.588 (.115)
Time spent caregiving ¹	-0.640 (.799)	-0.802 (.424)	1.359 (.774)	1.755 (.081)
Age patient	0.275 (.210)	1.304 (.194)	0.235 (.208)	1.130 (.260)
Treatment type ²	0.731 (.721)	1.014 (.312)	-0.739 (.710)	-1.041 (.299)
ED duration	0.062 (.029)	2.134 (.035)	-0.035 (.029)	-1.227 (.222)
BMI Perc.cat ³	0.432 (.709)	0.609 (.544)	-1.574 (.686)	-2.293 (.023)
EDE total score	0.777 (.243)	3.202 (.002)	-0.240 (.246)	-0.975 (.331)
EDI total score	0.039 (.008)	5.098 (<.001)	-0.009 (.008)	-1.083 (.281)

¹ 1 = < 3h/day, 2 = ≥ 3h/day; ² 1 = inpatient, 2 = outpatient, ³ 1 = ≤ 1st percentile, 2 = > 1st percentile; Abbreviations: BDI Beck Depression Inventory, CASK Caregiver Skills Scale, contemp. contemplation, EDE Eating Disorder Examination Interview, EDI-2 Eating Disorder Inventory-2, EDSIS Eating Disorders Symptoms Impact Scale, FQ-CC Family Questionnaire Criticism Score, FQ-EOI Family Questionnaire Emotional Overinvolvement Score, GHQ General Health Questionnaire, Perc.cat Percentile category, precontemp. precontemplation, STAI State/Trait Anxiety Inventory, URICA University of Rhode Island Change Assessment Scale.