

Applied Microbiology and Biotechnology

Characterization of the Genes Responsible for Rubber Degradation in *Actinoplanes* sp. Strain OR16.

Namiko Gibu ¹, Tomoka Arata ¹, Saya Kuboki ¹, Dao Viet Linh ^{1#}, Masao Fukuda ^{1\$},
Alexander Steinbüchel ^{2,3}, Daisuke Kasai ^{1*}

¹*Department of Bioengineering, Nagaoka University of Technology, Nagaoka, Japan*, ²*Institut für
Molekulare Mikrobiologie und Biotechnologie, Westfälische Wilhelms-Universität Münster, Münster,
Germany*, ³*Environmental Science Department, King Abdulaziz University, Jeddah, Saudi Arabia*.

*Corresponding author: Daisuke Kasai

Department of Bioengineering
Nagaoka University of Technology
Nagaoka, Niigata 940-2188, Japan
E-mail: dkasai1@vos.nagaokaut.ac.jp




Fig. S1. Effect of temperature (A, C, and E) and pH (B, D, and F) on Lcp activity. Relative oxygen consumption activities of Lcp1-his (A and B), Lcp2-his (C and D), and Lcp3-his (E and F) were indicated. Each value is the average \pm standard deviation of three independent experiments performed in parallel.