

Supplementary Materials

Impaired mesocorticolimbic connectivity underlies increased mechanical pain sensitivity
in chronic low back pain

Siyi Yu¹, Wen Li², Wei Shen¹, Robert R. Edwards³, Randy L. Gollub¹, Georgia Wilson¹, Joel Park¹,
Ana Ortiz¹, Jin Cao¹, Jessica Gerber⁴, Ishtiaq Mawla¹, Suk-Tak Chan⁴, Jeungchan Lee⁴, Ajay D.
Wasan⁵, Vitaly Napadow⁴, Ted J. Kaptchuk⁶, Bruce Rosen⁴, Jian Kong^{1,4}

¹Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA,
USA

²Department of Psychology, Florida State University, Tallahassee, FL, USA

³Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Harvard
Medical School, Boston, MA, USA

⁴Department of Radiology, Martinos Center for Biomedical Imaging, Massachusetts General Hospital,
Harvard Medical School, Charlestown, MA, USA

⁵Department of Anesthesiology, Center for Pain Research, University of Pittsburgh, Pittsburgh, PA, USA

⁶Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

Correspondence to:

Jian Kong,

Department of Psychiatry,

Massachusetts General Hospital,

Charlestown, MA 02129

E-mail: jkong2@mgh.harvard.edu

Table S1. Medication usage in cLBP group


Medication	Number of patients
Opioid analgesics ^a	6
Benzodiazepines ^b	3
Muscle relaxants ^c	5

Notes:

^a Oxycodone, 5mg, TID; Hydrocodone, 5mg, PRN or TID; Tramadol, 50mg, TID or PRN; Percocet, 10mg, TID or PRN.

^b Lorazepam, 1-3mg, PRN; clonazepam, 0.5mg, PRN, Valium, 10mg, PRN.

^c Flexeril/Cyclobenzaprine, 5-10mg, TID.



Supplementary Figure 1. Pain ratings before and after manipulation for each patient