

Table S1. Relationship between tumor location and clinicopathological characteristics of the 117 colorectal cancer patients

Characteristic (n=117)	Right-sided (n=30)	%	Left-sided (n=87)	%	Total	P-value
Sex						0.5119
Male	18	60%	58	67%	76	
Female	12	40%	29	33%	41	
Age (years), median (range)	76 (46-86)		68 (29-89)			0.0360*
≤70	12	40%	54	62%	66	
>70	18	60%	33	38%	51	
Tumor depth						0.8970
T1-2	3	10%	8	9%	11	
T3-4	27	90%	79	91%	106	
Lymph node metastasis						0.0078*
Negative	23	77%	43	49%	66	
Positive	7	23%	44	51%	51	
Tumor differentiation						0.4757
Well/moderate	25	83%	77	89%	102	
Others	5	17%	10	11%	15	
Lymphatic invasion						0.2191
Negative	17	57%	38	44%	55	
Positive	13	43%	49	56%	62	
Venous invasion						0.7832
Negative	8	27%	21	24%	29	
Positive	22	73%	66	76%	88	
Perineural invasion						0.2421
Negative	26	87%	67	77%	93	
Positive	4	13%	20	23%	24	
Mismatch repair						0.0157*
Dificient	6	20%	4	5%	10	
Proficient	24	80%	83	95%	107	
Adjuvant chemotherapy						0.0885
No	23	77%	52	60%	75	
Yes	7	23%	35	40%	42	
Recurrence						0.8956
No	8	27%	24	28%	33	
Yes	22	73%	63	72%	84	