

### Supplementary material

**Table S1** Baseline Characteristics of patients with and without follow-up.

	Patients with follow-up (n=104)	Patients without follow-up (n=125)	<i>P</i> value
Age, y (mean $\pm$ SD)	$64.0 \pm 9.8$	$60.6 \pm 9.6$	0.009*
Sex, female (%)	34 (32.7)	45 (36.0)	0.600
Education, y (median $\pm$ IQR)	$8.5 \pm 7.0$	$7.0 \pm 8.0$	0.465
BMI, kg/m <sup>2</sup> (mean $\pm$ SD)	$24.4 \pm 2.8$	$24.4 \pm 3.1$	0.924
Baseline NIHSS score (median $\pm$ IQR)	$1.0 \pm 2.0$	$1.0 \pm 2.0$	0.038*
Baseline MMSE score (median $\pm$ IQR)	$27.5 \pm 3.0$	$27.0 \pm 3.0$	0.097
Baseline MoCA score (mean $\pm$ SD)	$22.0 \pm 8.0$	$22.0 \pm 6.0$	0.385

SD, standard deviation; IQR, inter-quartile range; BMI, body mass index; NIHSS, National Institutes of Health Stroke Scale; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment; \**P* < 0.05.