

SUPPLEMENTAL MATERIAL

Table S1. External analysis of the clinical decisions after knowledge of sFlt-1/PIGF > 655 according to the presence of preeclampsia at inclusion.

Clinical decisions after knowledge of sFlt-1/PIGF >655	Total (n=237)	Preeclampsia at inclusion		
		No (n=52)	Yes (n=185)	p
Hospitalization				
- No	30 (12.7)	15 (28.8)	15 (8.1)	<0.01
- Previously hospitalized	149 (62.9)	19 (36.5)	130 (70.3)	
- Yes, following current protocols	44 (18.6)	11 (21.2)	33 (17.8)	
- Yes, conditioned by the knowledge of the sFlt-1/PIGF	14 (5.9)	7 (13.5)	7 (3.8)	
Fetal maturation				
- No	36 (15.2)	14 (26.9)	22 (11.9)	0.02
- Yes, previously indicated	146 (61.6)	25 (48.1)	121 (65.4)	
- Yes, following current protocols	41 (17.3)	8 (15.4)	32 (17.3)	
- Yes, conditioned by the knowledge of the sFlt-1/PIGF	14 (5.9)	5 (9.6)	6 (3.2)	
Indication of delivery				
- Spontaneous onset of labor	2 (0.8)	0 (0)	2 (0.1)	0.99
- Iatrogenic, following current protocols	212 (89.5)	47 (90.4)	165 (89.2)	
- Iatrogenic, conditioned by the knowledge of the sFlt-1/PIGF	23 (9.7)	5 (9.6)	18 (9.7)	
Overall adequacy				
- Correct, according to current protocols	205 (86.5)	41 (78.8)	164 (88.6)	0.22
- Correct but not following current protocols	20 (8.4)	8 (15.4)	12 (6.5)	
- Incorrect	7 (3.0)	2 (3.8)	5 (2.7)	
- Unable to establish	5 (2.1)	1 (1.9)	4 (2.2)	

Figure S1. Time until delivery according to the presence of PE at inclusion or its subsequent development.

