

SUPPLEMENTAL MATERIAL

Table S1. Angiogenic Biomarkers Stratified by Diagnosis.

	ASPRIN	NO ASPIRIN	P-Value
Patients with Preeclampsia	N = 11	N = 39	
Activin, ng/ml	8.40 (5.84, 24.46)	24.95 (11.89, 36.49)	0.01
Follistatin, ng/ml	16.15 (10.93, 21.10)	13.29 (6.71, 18.27)	0.19
FSTL3, ng/ml	36.71 (26.60, 61.69)	45.93 (37.51, 65.34)	0.18
Activin/Follistatin	0.76 (0.31, 1.70)	1.89 (0.85, 3.63)	0.01
Activin/FSTL3	0.23 (0.22, 0.29)	0.49 (0.28, 0.66)	0.01
Patients with Superimposed Preeclampsia	N = 14	N = 28	
Activin, ng/ml	6.10 (3.47, 9.93)	9.31 (6.30, 12.13)	0.06
Follistatin, ng/ml	9.06 (5.89, 14.25)	11.43 (8.79, 16.42)	0.34
FSTL3, ng/ml	34.48 (26.07, 50.02)	38.56 (24.15, 47.98)	0.96
Activin/Follistatin	0.53 (0.27, 0.73)	0.75 (0.39, 1.01)	0.19
Activin/FSTL3	0.15 (0.07, 0.27)	0.21 (0.16, 0.28)	0.07

Data is presented as median (quartile 1, quartile 3) or n (%) depending on variable type. These results are assessed between groups with a Wilcoxon Rank Sum or chi-square test, respectively. BP = blood pressure, ECHO = echocardiogram