

Supplementary Online Content

Bao W, Liu B, Rong S, Dai SY, Trasande L, Lehmler H-J. Association between bisphenol A exposure and risk of all-cause and cause-specific mortality in US adults. *JAMA Netw Open.* 2020;3(8):e2011620. doi:10.1001/jamanetworkopen.2020.11620

eTable. Stratified Analyses for the Association of Urinary BPA Levels With Cause-Specific Mortality

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Stratified Analyses for the Association of Urinary BPA Levels With Cause-Specific Mortality

Variable		Tertiles of urinary BPA levels			P for interaction
		Tertile 1	Tertile 2	Tertile 3	
CVD mortality					
Age	<65 years	1 (reference)	0.70 (0.22-2.18) [†]	1.67 (0.52-5.31)	0.71
	≥65 years	1 (reference)	1.17 (0.49-2.79)	1.71 (0.51-5.73)	
Sex	Male	1 (reference)	1.08 (0.35-3.29)	1.75 (0.50-6.19)	0.49
	Female	1 (reference)	0.98 (0.37-2.63)	1.11 (0.30-4.16)	
Race/ethnicity	Whites	1 (reference)	1.72 (0.71-4.16)	1.83 (0.51-6.49)	0.72
	Non-whites	1 (reference)	0.26 (0.07-1.00)	0.80 (0.35-1.82)	
Diet quality [#]	Lower	1 (reference)	1.45 (0.58-3.61)	0.91 (0.28-2.94)	0.46
	Higher	1 (reference)	0.64 (0.19-2.19)	2.27 (0.72-7.22)	
Physical activity [*]	Lower	1 (reference)	1.25 (0.55-2.84)	0.93 (0.30-2.90)	0.63
	Higher	1 (reference)	0.60 (0.13-2.71)	2.13 (0.40-11.36)	
Obesity	BMI < 30	1 (reference)	0.95 (0.32-2.80)	1.28 (0.50-3.29)	0.14
	BMI ≥ 30	1 (reference)	1.55 (0.43-5.66)	1.73 (0.29-10.2)	
Cancer mortality					
Age	<65 years	1 (reference)	1.03 (0.33-3.13)	1.08 (0.31-3.71)	0.29
	≥65 years	1 (reference)	1.00 (0.40-2.47)	1.03 (0.36-2.92)	
Sex	Male	1 (reference)	0.68 (0.29-1.59)	0.82 (0.36-1.87)	0.17
	Female	1 (reference)	1.75 (0.45-6.81)	1.28 (0.27-6.11)	
Race/ethnicity	Whites	1 (reference)	1.08 (0.30-3.81)	0.92 (0.27-3.10)	0.46

	Non-whites	1 (reference)	1.12 (0.45-2.81)	1.11 (0.48-2.60)	
Diet quality [#]	Lower	1 (reference)	1.64 (0.56-4.78)	1.24 (0.36-4.23)	0.21
	Higher	1 (reference)	0.40 (0.10-1.59)	0.60 (0.17-2.12)	
Physical activity [*]	Lower	1 (reference)	1.10 (0.28-4.26)	0.49 (0.10-2.37)	0.94
	Higher	1 (reference)	1.06 (0.40-2.81)	1.89 (0.74-4.80)	
Obesity	BMI < 30	1 (reference)	1.45 (0.57-3.68)	1.47 (0.59-3.64)	0.65
	BMI ≥ 30	1 (reference)	0.77 (0.17-3.59)	0.59 (0.14-2.44)	

[#] Lower or higher diet quality was defined as the healthy eating index (HEI) < the median score or ≥ the median score, respectively.

^{*} Lower or higher physical activity level was defined as below or meeting the physical activity guidelines, respectively.

Adjustment for age, sex, race/ethnicity, urinary creatinine levels, education, family income status, smoking, alcohol drinking, physical activity, total energy intake, HEI2010 score, and BMI.

[†] HR (95% CI).