Appendix to Evaluation of a city-wide school-located influenza vaccination program in Oakland, California with respect to vaccination coverage, school absences, and laboratory-confirmed influenza: a matched cohort study

## S3 Table. Percentage of elementary students whose caregiver reported they were vaccinated for influenza by vaccine type in each district

Site	Vaccine type	2014-15	2015-16	2016-17	2017-18
Comparison	Shot	48.4 (46.1, 50.8)	52.2 (50.7, 53.7)	51.5 (48.6, 54.4)	53.4 (50.7, 56.1)
Comparison	Spray	12.4 (10.3, 14.6)	11.6 (9.9, 13.3)	a	a
	Error/Missing/				
Comparison	Don't know	3.2 (2.6, 3.8)	2.1 (1.6, 2.6)	4.9 (4.1 <i>,</i> 5.7)	3.0 (2.3, 3.8)
Intervention	Shot	35.8 (32.9, 38.6)	39.1 (35.5, 42.8)	52.4 (49.2, 55.6)	64.4 (59.7, 69.0)
Intervention	Spray	18.6 (14.5, 22.6)	22.5 (18.5, 26.5)	a	a
	Error/Missing/				
Intervention	Don't know	4.9 (3.8, 6.0)	5.9 (4.5, 7.4)	11.6 (9.2, 13.9)	3.1 (2.2, 3.9)

<sup>&</sup>lt;sup>a</sup> The spray (live attenuated) vaccine was not available 2016-17 or 20.