

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-036268
Article Type:	Protocol
Date Submitted by the Author:	08-Dec-2019
Complete List of Authors:	Zhou, Lin-yue ; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Zhang, Yuan ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Tian, Yuan; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Fu, Xiaoxu; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Wang, Li-zhen ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Xie, Chun-guang; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine
Keywords:	DIABETES & ENDOCRINOLOGY, MENTAL HEALTH, COMPLEMENTARY MEDICINE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Lin-yue Zhou,^{1,2} Yuan Zhang,² Yuan Tian,² Xiaoxu Fu,^{1,2} Li-zhen Wang,² Chun-guang Xie*^{1,2}

Author affiliations

¹Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Chengdu, Sichuan Province, China

²School of Clinical Medicine, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan Province, China

*Correspondence to:

Corresponding Author: Professor Chun-guang Xie

Address: Department of Endocrinology, Hospital of Chengdu University of Traditional Chinese Medicine, No 39 Shi-er-qiao Road, Chengdu, Sichuan Province, China, 610072

Tel: (86)13438826824

E-mail: killuya233@stu.cdutcm.edu.cn

Word count: 1978 words (excluding title page, abstract, references, figures and tables)

ABSTRACT

Introduction About 463 million adults aged 20 to 79 have diabetes globally and mental disorders often exist in diabetic patients as comorbidities. The comorbidities can lead to deterioration of diseases, increased difficulty in treatment, even as well as the terrible mortality. Music intervention has been applied in the treatment of the comorbidities for 12 years, but there are no any recommendations due to lacking of evidence. Thus, a meta-analysis is necessary to be conducted to evaluate the effect of music intervention.

Methods and analysis We will retrieve the following nine online electronic databases: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical Database (CBM), from their inception until August 2019. Only randomized controlled trials (RCTs) of music intervention for depression or anxiety with diabetes will be involved. The primary outcomes include depression score and anxiety score measured by scales, and the secondary outcome is safety. Data extraction will be implemented by 2 researchers independently. Risk of bias will be evaluated through Cochrane Collaboration's Risk of Bias tool. Eventually, all the data will be analyzed via the software Review Manager V.5.3.

Ethics and dissemination This meta-analysis will provide information about music intervention applied to treat depression or anxiety in diabetic patients. There is no required ethical approval because this is a protocol for meta-analysis. The results of this study will be published to a peer reviewed journal.

Trial registration number CRD42019146439

Strengths and limitations of this study

· This meta-analysis will provide evidence of the effectiveness and safety of music intervention applied to treat depression or anxiety in diabetic patients for the first time.

· This systematic review will furnish evidence reference for guideline of relevant areas.

· This meta-analysis will be implemented on the basis of rigorous international gold-standard methodology and integrated search strategy.

· A language bias may exist in this meta-analysis because it will include studies only in English and Chinese on account of the language barrier.

INTRODUCTION

Description of the conditions

According to the newest “IDF Diabetes Atlas (9th edition)” published by International Diabetes Federation (IDF), about 463 million adults aged 20 to 79 have diabetes globally (1 in 11).[1] It is well known that diabetes has a significant influence on life quality because it can cause a lot of complications such as retinopathy, renal disease, cardiovascular disease, neurological disease and foot ulcers which may result in lower extremity amputation.[2]

In addition to physical complications, some mental disorders often exist in diabetic patients as comorbidities. A few studies suggested that depression was more prevalent in diabetic patients compared to those without.[3,4] At the same time, depression could also increase the risk of incident of diabetes,[5-7] which suggested that depression had a bidirectional association with diabetes. And the relation between anxiety and diabetes was confirmed to be the same. People with diabetes were more susceptible to concurrent anxiety[8,9] while reversely anxiety was a risk factor of diabetic incidents.[10] Individuals with chronic diseases and also comorbid depression or anxiety have obviously higher health-care utilization, functional disability, work absence and mortality risk than those without.[11,12]

Description of the intervention

When it comes to the treatment of diabetes with depression or anxiety, conventional glucose-lowering treatment combined with drug therapy or psychological intervention are the routine treatments.[13,14] Music intervention is also a successful therapy using music for treatment by professionals, which can be divided into two major categories, namely music therapy and music medicine, although the distinction between them are often neglected.[15] A lot of studies suggested that music, no matter what type of it is, demonstrated a remarkable effect in alleviating depression and anxiety among patients with various diseases.[16-18] As for diabetes, recent studies showed that music intervention could improve fasting blood-glucose and systolic blood pressure in diabetic patients.[19,20] Good control of blood glucose, blood lipids and blood pressure can effectively

1
2
3
4 reduce the incidence of serious complications of diabetes.[21]
5
6

7 8 **How the intervention might work**

9
10 Music psychology researches have shown the special potential of music that music listening can
11 give rise to complex emotional experiences[22, 23] and strengthen ability to deal with complex
12 negative emotions in everyday life.[24] The proposed mechanisms of music intervention on
13 depression and anxiety span several domains. Firstly, music intervention can exert
14 neurophysiological and neurochemical effects, such as triggering endorphin release[25] and
15 stimulating cognitive functioning.[26] Secondly, music intervention can promote positive self-
16 emotions including enhancing the sense of happiness[27] and activating self-efficacy and self-
17 confidence through experiences of success.[28] Lastly, music intervention can build up sociability,
18 such as interaction skills[29] and interpersonal relationship.[30] Furthermore, there are published
19 studies indicated that music intervention combined with diabetic routine treatments could improve
20 depression score, anxiety score, fasting glucose and glycosylated hemoglobin (HbA1c).[31,32]
21
22
23
24
25
26
27
28
29
30
31
32

33 **Why it is important to this review**

34
35 From the above, many studies indicated the bidirectional relationship between diabetes and mental
36 disorders. Music intervention, as a therapeutic method, was applied to alleviate depression and
37 anxiety and improve related indicators of diabetes. By searching literatures, it can be found that
38 some reviews evaluated the effect of music intervention on patients with mental disorders,[33-35]
39 while no related systematic review and meta-analysis can be searched about the effect of music
40 intervention on mental health of patients with diabetes. Therefore, if music intervention can be
41 proved to be both valid and with few side effects through this study, it is worth being promoted and
42 applied in clinic all over the world to benefit more people since this method is easy to be operated.
43
44
45
46
47
48
49
50
51

52 **Objectives**

53
54 The purpose of this review is to systematically assess the effect of music intervention on depression
55
56
57
58
59
60

1
2
3
4 and anxiety of diabetic patients. We also look forward to furnish evidence reference for related
5
6 guideline.

7 8 9 **METHODS**

10 11 **Study registration**

12
13 The protocol of the meta-analysis has been registered in International Prospective Register of
14 Systematic Reviews (PROSPERO), and CRD42019146439 is the registration number. This
15 systematic review protocol will be conducted and reported severely on the basis of Preferred
16 Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)[36] statement guidelines,
17 and the crucial protocol amendments will be documented in the full review.
18
19
20
21
22

23 24 25 **Eligibility criteria**

26 We will severely screen studies that have compliance with following inclusion criteria.

27 28 29 30 31 Type of studies

32 We will only cover randomized controlled trials (RCTs) in English or Chinese, excepting Quasi-
33 RCTs and cluster RCTs. Experiments on animals and nonrandomized clinical trials will be out of
34 our consideration. Articles with significant overlap with other articles through print or e-publishing
35 will fail to be included. Duplicate publications, that is results originated from a single experiment
36 but measured by other standards and then published as disparate papers, will also be excluded.
37 Publications with originality is our first choice. The publication status will not be restricted.
38
39
40
41
42
43
44

45 46 47 Type of Participants

48 Diabetic Patients with depression or anxiety symptoms, regardless of the type of diabetes and no
49 restriction on gender, age, race, economic ability and education level. Patients of other disease or
50 healthy persons will be excluded.
51
52
53
54
55
56
57
58
59
60

Type of interventions and controls

Interventions include music alone or music in combination with other interventions. No restriction on the type of music. Other interventions refer to: (1) routine treatment for diabetes which mainly includes pharmacologic approaches to glycemic treatment, lifestyle management, etc. according to the newest guideline of American Diabetes Association (ADA);[37,38] (2) Other therapies such as acupuncture therapy,[39] metabolic surgery[40] etc. When intervention is music alone, controls can be no treatment, placebo or other active therapies. Notably, when intervention is music combined with active therapies, the control must be the same active therapies.

Type of outcome measures

This review aims at systematically assessing the effect of music intervention on depression or anxiety of diabetic patients, so we choose depression score and anxiety score as the primary outcomes which can give an index to the degree of depression and anxiety by scales such as SDS, HAMD, SAS or HAMA.

Primary outcomes: Depression score and anxiety score. Depression score is measured using self-rating depression scale (SDS) or Hamilton depression scale (HAMD). Anxiety score is measured using self-rating anxiety scale (SAS) or Hamilton anxiety scale (HAMA).

Secondary outcomes: Safety. Measurement via incidence of adverse effects.

Search methods

Search resources

We will retrieve the following nine online electronic databases: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical Database (CBM), from their inception until August 2019. We also scrutinized the reference list of relevant publications for additional studies.

Search strategies

The search strategy will include the MeSH terms for diabetes mellitus, depression, anxiety and music therapy, combined with their respective free-text terms. The search details of the PubMed was simply shown in table 1.

Table 1 Example of PubMed search strategy.

Number	Search terms
#1	"Diabetes Mellitus"[Mesh]
#2	"Depression"[Mesh]) OR Depressions[Title/Abstract]) OR Depressive Symptoms[Title/Abstract] OR Depressive Symptom[Title/Abstract] OR Symptom, Depressive[Title/Abstract] OR Symptoms, Depressive[Title/Abstract] OR Emotional Depression[Title/Abstract] OR Depression, Emotional[Title/Abstract] OR Depressions, Emotional[Title/Abstract] OR Emotional Depressions[Title/Abstract]
#3	"Anxiety"[Mesh] OR Hypervigilance [Title/Abstract] OR Nervousness [Title/Abstract]
#4	#2 OR #3
#5	"Music Therapy"[Mesh] OR Therapy, Music [Title/Abstract]
#6	randomized controlled trial [Publication Type] OR randomized [Title/Abstract] OR placebo [Title/Abstract]
#7	#1 AND #4 AND #5 AND #6

Data collection and analysis

Studies selection

Firstly, two researchers (L-yZ and YT) will respectively screen the keywords, titles and abstracts of all obtained studies using endnote x9 software and determine whether trials have compliance with inclusion criteria. Secondly, we will also get access to the full text of all studies of possible relevance for further assessment. Each of excluded study will be noted with reasons. Any

1
2
3
4 disagreements will be settled through consensus or referral to the third researcher (YZ). The
5 specific selection steps were presented in the figure 1.
6
7

8 9 Data extraction

10 Two researchers (L-yZ and YT) will independently extract and tabulate the following data items
11 via Microsoft Excel:
12
13

- 14 1. General information (including publication year and name of the first author).
- 15 2. Study design (including random method, allocation concealment and blinding).
- 16 3. Participants (including sample size, gender, age, depression score, anxiety score and course of
17 disease).
- 18 4. Treatment (including interventions, controls, music type and the course of treatment).
- 19 5. Outcomes (SDS, HAMD, SAS or HAMA and safety).

20
21 If there is any different opinion, the final decision will go to the third researcher (X-xF) to be
22 made. If the required data are unavailable from literatures, one researcher (YZ) will send an e-mail
23 to the corresponding author and ask for further information
24
25

26 27 Assessment of risk of bias

28 The Cochrane Collaboration's Risk of Bias tool[41] will be applied to evaluate the risk of bias of
29 all included RCT studies by two authors (L-zW and YZ). The following domains will be assessed:
30 sequence generation, allocation concealment, blinding of participants and personnel, blinding of
31 outcome assessors, incomplete outcome data, selective reporting and other sources of bias. The
32 assessments will be classified into "low", "unclear" or "high" of risk.
33
34

35 36 Data analysis

37 RevMan 5.3. software provided by the Cochrane collaboration will be used to analyze the data.
38 Considering that all outcomes (depression score, anxiety score and safety) are continuous variables,
39 the weighted mean difference (WMD) and 95% confidence interval (CI) will be calculated when
40
41
42
43
44
45
46
47
48
49

1
2
3
4 the quantitative data are from the same measurement method or small data scale. On the contrary,
5 the standardized mean difference (SMD) and 95% CI will be calculated.
6

7 A meta-analysis will be implemented if the included studies are adequately homogenous. *P*-
8 value and *I*² will be used to assess the heterogeneity of statistic. When *P*-value >0.05 and *I*² ≤50%,
9 the fixed-effects model will be used for statistical combination. When *P*-value <0.05 or *I*² >50%,
10 subgroup analyses or sensitivity analyses will be performed. If a meta-analysis is unavailable,
11 descriptive summaries will be provided.
12
13
14
15
16

17 18 19 Assessment of reporting bias

20 Publication bias will be implemented, if there are more than 10 studies included, through a funnel
21 plot.
22
23
24

25 26 27 Patient and public involvement

28 Patients and public involvement will fail to be included in this study.
29
30
31

32 33 Evidence assessed

34 GRADE guideline will be used to evaluate the quality of evidence.[42] To fulfil transparency and
35 simplification, the quality of evidence can be assessed by 4 levels: “high”, “medium”, “low” and
36 “very low”. GRADE profiler 3.2 will be applied for analysis.
37
38
39
40
41

42 43 **DISCUSSION**

44 Depression and anxiety often occur independently or simultaneously during the development of
45 many chronic diseases. In recent years, the number of studies about comorbid depression or anxiety
46 in diabetics are continually increasing, not only because the two-way relationship between
47 depression/anxiety and diabetes can lead to deterioration of diseases, increased difficulty in
48 treatment, even as well as the terrible mortality,[43-46] but also because therapeutic methods for
49 treating the comorbidities is limited and unsatisfactory. On the one hand, some antidepressant drugs,
50
51
52
53
54
55
56

1
2
3
4 such as nortriptyline and sertraline, may complicate glycemic control.[47] On the other hand,
5
6 psychotherapy requires a professional psychiatrist and it is expensive.

7
8 Music intervention has been proved by many literatures to have a beneficial effect on depression
9
10 and anxiety, such as strengthening awareness and sensitiveness for positive emotions.[48]
11
12 Noticeably, a few studies also show that music intervention can reduce blood glucose and HbA1c
13
14 simultaneously.[20,49] Therefore, music intervention is an applicable method for treating
15
16 depression and anxiety with diabetes. Additionally, music intervention, almost no side effects, is
17
18 safer than pharmacotherapy, easier to be operated and costs less than psychotherapy, so it is
19
20 conducive to clinical promotion and application.

21
22 Since 2007,[50] music intervention has been applied in the treatment of depression and anxiety
23
24 with diabetes for 12 years. Nonetheless, no any systematic review or meta-analysis were conducted
25
26 to evaluate the potential benefits and harms of this therapy method. Thus, this study intends to
27
28 provide strong evidence for music intervention applied to treat mental disorders in diabetic patients
29
30 and synchronously furnish evidence reference for related guideline.

31
32 However, a language bias may exist in this meta-analysis because it will include studies only in
33
34 English and Chinese because of the language barrier. Therefore, some related studies from medical
35
36 databases in other languages might be missed.

37
38 In conclusion, this meta-analysis will be conducive to assess underlying benefits and harms of
39
40 music intervention on comorbid depression or anxiety with diabetes. Moreover, the results of this
41
42 study might not merely furnish reference basis for related guideline, but also may attract more
43
44 attention on music intervention, which can be a chance to facilitate its clinical application and
45
46 ultimately benefit more patients.

47 48 **ETHICS AND DISSEMINATION**

49
50 There is no required ethical approval because this is a protocol for meta-analysis. The results of
51
52 this study will be published to a peer reviewed journal.
53
54
55
56

Contributions L-yZ and C-gX contributed to the conception of the study. The manuscript of the protocol was drafted by L-yZ, and was revised by XxF and C-gX. L-yZ and YT will also independently screen the potential studies and extract data from included studies. L-zW and YZ will assess the risk of bias and accomplish data synthesis. XxF will arbitrate any disagreements. C-gX will ensure that no errors occur during the review. All authors have approved the publication of the protocol.

Funding This study was supported by grants from the Science and Technology Project of Sichuan Province (2019YFS002), the Special Scientific and Technological Research Project of Administration of TCM of Sichuan Province (2018KF003), and the Foundation of Development of Science and Technology of Chengdu University of TCM (2016-D-YY-61).

Competing interests None.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

References

1. International Diabetes Federation. IDF Diabetes Atlas (9th edition), 2019. Available: <http://www.diabetesatlas.org/>. Accessed November 9, 2019
2. World Health Organization. Global report on diabetes, 2014. Available: <https://www.who.int/news-room/fact-sheets/detail/diabetes>. Accessed November 9, 2019
3. Ali S, Stone MA, Peters JL, et al. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* 2006;23(11):1165–1173. doi:10.1111/j.1464-5491.2006.01943.x

- 1
2
3
4 4. Anderson RJ, Freedland KE, Clouse RE, et al. The prevalence of comorbid depression in
5 adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24(6):1069–1078.
6
7 doi:10.2337/diacare.24.6.1069
8
- 9
10 5. Knol MJ, Twisk JW, Beekman AT, et al. Depression as a risk factor for the onset of type 2
11 diabetes mellitus. A meta-analysis. *Diabetologia* 2006;49(5):837–845.doi:10.1007/s00125-
12 006-0159-x
13
- 14
15 6. Cosgrove MP, Sargeant LA, Griffin SJ. Does depression increase the risk of developing type
16 2 diabetes?. *Occup Med (Lond)* 2008;58(1):7–14.doi:10.1093/occmed/kqm105
17
- 18
19 7. Mezuk B, Eaton WW, Albrecht S, et al. Depression and type 2 diabetes over the lifespan: a
20 meta-analysis. *Diabetes Care* 2008;31(12):2383–2390.doi:10.2337/dc08-0985
21
- 22
23 8. Grigsby AB, Anderson RJ, Freedland KE, et al. Prevalence of anxiety in adults with diabetes:
24 a systematic review. *J Psychosom Res* 2002; 53: 1053–1060.doi:10.1016/s0022-
25 3999(02)00417-8
26
- 27
28 9. Smith KJ, Beland M, Clyde M, et al. Association of diabetes with anxiety: a systematic
29 review and metaanalysis. *J Psychosom Res* 2013; 74: 89–99.
30
31 doi:10.1016/j.jpsychores.2012.11.013
32
- 33
34 10. Smith KJ, Deschênes SS, Schmitz N. Investigating the longitudinal association between
35 diabetes and anxiety: a systematic review and meta-analysis. *Diabet Med* 2018;35(6):677–
36 693.doi:10.1111/dme.13606
37
- 38
39 11. Egede LE. Major depression in individuals with chronic medical disorders: prevalence,
40 correlates and association with health resource utilization, lost productivity and functional
41 disability. *Gen Hosp Psychiatry* 2007;29:409–416.doi:10.1016/j.genhosppsy.2007.06.002
42
- 43
44 12. Naicker K, Johnson JA, Skogen JC, et al. Type 2 Diabetes and Comorbid Symptoms of
45 Depression and Anxiety: Longitudinal Associations With Mortality Risk. *Diabetes Care*
46 2017;40(3):352–358.doi:10.2337/dc16-2018
47
- 48
49 13. van der Feltz-Cornelis CM, Nuyen J, Stoop C, et al. Effect of interventions for major
50 depressive disorder and significant depressive symptoms in patients with diabetes
51
52
53
54
55
56

- 1
2
3
4 mellitus: a systematic review and meta-analysis. *Gen Hosp Psychiatry* 2010;32(4):380–395.
5
6 doi:10.1016/j.genhosppsy.2010.03.011
- 7
8 14. Komsuoglu Celikyurt I, Mutlu O, Ulak G, et al. Exenatide treatment exerts anxiolytic- and
9
10 antidepressant-like effects and reverses neuropathy in a mouse model of type-2 diabetes. *Med*
11
12 *Sci Monit Basic Res* 2014;20:112–117.doi:10.12659/MSMBR.891168
- 13
14 15. Leubner D, Hinterberger T. Reviewing the Effectiveness of Music Interventions in Treating
15
16 Depression. *Front Psychol* 2017;8:1109.doi:10.3389/fpsyg.2017.01109
- 17
18 16. Thaut MH, Gardiner JC, Holmberg D, et al. Neurologic music therapy improves executive
19
20 function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*
21
22 2009;1169:406–416.doi:10.1111/j.1749-6632.2009.04585.x
- 23
24 17. Kim DS, Park YG, Choi JH, et al. Effects of music therapy on mood in stroke patients.
25
26 *Yonsei Med J* 2011;52(6):977–981.doi:10.3349/ymj.2011.52.6.977
- 27
28 18. Nwebube C, Glover V, Stewart L. Prenatal listening to songs composed for pregnancy and
29
30 symptoms of anxiety and depression: a pilot study. *BMC Complement Altern Med*
31
32 2017;17(1):256.doi:10.1186/s12906-017-1759-3
- 33
34 19. Lin XX, Ji JM, Zhou GY. Influence of listening to Jiao tune music on sleep quality and blood
35
36 glucose in the elderly patients with type 2 diabetes accompanied by insomnia. *Journal of*
37
38 *rehabilitation* 2017;27(01):44-48.
- 39
40 20. Mandel SE, Davis BA, Secic M. Effects of music therapy and music-assisted relaxation and
41
42 imagery on health-related outcomes in diabetes education: a feasibility study. *Diabetes Educ*
43
44 2013;39(4):568–581.doi:10.1177/0145721713492216
- 45
46 21. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2014.
47
48 Available: [https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf)
49
50 [burden-in-the-united-states.pdf](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf). Accessed November 9, 2019
- 51
52 22. Gabrielsson A, Wik SL. Strong experiences related to music: a descriptive system. *Music Sci*
53
54 2003;7(2):157–217.
- 55
56 23. Zentner M, Grandjean D, Scherer KR. Emotions evoked by the sound of music:

- 1
2
3
4 characterization, classification, and measurement. *Emotion* 2008;8(4):494–521.
5
6 doi:10.1037/1528-3542.8.4.494
- 7
8 24. Saarikallio S. Music as emotional self-regulation throughout adulthood. *Psychol Music*
9
10 2011;39(3):307–327.
- 11
12 25. Gök Ugur H, Yaman Aktaş Y, Orak OS, et al. The effect of music therapy on depression and
13
14 physiological parameters in elderly people living in a Turkish nursing home: a randomized-
15
16 controlled trial. *Aging Ment Health* 2017;21(12):1280–1286.
17
18 doi:10.1080/13607863.2016.1222348
- 19
20 26. Guétin S, Portet F, Picot MC, et al. Effect of music therapy on anxiety and depression in
21
22 patients with Alzheimer's type dementia: randomised, controlled study. *Dement Geriatr Cogn*
23
24 *Disord* 2009;28(1):36–46. doi:10.1159/000229024
- 25
26 27. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement
27
28 intervention on the cognitive functions of people with moderate dementia: a randomized
29
30 controlled trial. *Aging Ment Health* 2018;22(3):306–315.
31
32 doi:10.1080/13607863.2016.1251571
- 33
34 28. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in
35
36 elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging*
37
38 *Ment Health* 2017;21(2):156–162. doi:10.1080/13607863.2015.1093602
- 39
40 29. Ashida S. The effect of reminiscence music therapy sessions on changes in depressive
41
42 symptoms in elderly persons with dementia. *J Music Ther* 2000;37(3):170–182.
43
44 doi:10.1093/jmt/37.3.170
- 45
46 30. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a
47
48 controlled randomized study. *Neurol Sci* 2017;38(8):1485–1493. doi:10.1007/s10072-017-
49
50 3003-9
- 51
52 31. Deng HX, Gao GS, Li ZM, et al. Application of receiving music therapy in elderly patients
53
54 with diabetes. *Chinese journal of health and nutrition (1st Ten-days of month)*
55
56 2013;23(3):1186-1186.

- 1
2
3
4 32. Zhang ZW. Application of music therapy and psychotherapy adjuvant therapy in type 2
5 diabetes. *Chinese medical science* 2011,01(18):150-151.
6
7
8 33. Solanki MS, Zafar M, Rastogi R. Music as a therapy: role in psychiatry. *Asian J Psychiatr*
9 2013;6(3):193–199.doi:10.1016/j.ajp.2012.12.001
10
11 34. Petrovsky D, Cacchione PZ, George M. Review of the effect of music interventions on
12 symptoms of anxiety and depression in older adults with mild dementia. *Int Psychogeriatr*
13 2015;27(10):1661–1670.doi:10.1017/S1041610215000393
14
15 35. Aalbers S, Fusar-Poli L, Freeman RE, et al. Music therapy for depression. *Cochrane*
16 *Database Syst Rev* 2017;11(11):CD004517. Published 2017 Nov 16.
17 doi:10.1002/14651858.CD004517.pub3
18
19 36. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and
20 meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*
21 2015;350:g7647.doi:10.1136/bmj.g7647
22
23 37. American Diabetes Association. 9. Pharmacologic Approaches to Glycemic Treatment:
24 Standards of Medical Care in Diabetes-2019. *Diabetes Care* 2019;42(Suppl 1):S90–S102.
25 doi:10.2337/dc19-S009
26
27 38. American Diabetes Association. 5. Lifestyle Management: Standards of Medical Care in
28 Diabetes-2019. *Diabetes Care* 2019;42(Suppl 1):S46–S60.doi:10.2337/dc19-S005
29
30 39. Feng Y, Fang Y, Wang Y, et al. Acupoint Therapy on Diabetes Mellitus and Its Common
31 Chronic Complications: A Review of Its Mechanisms. *Biomed Res Int* 2018;2018:3128378.
32 doi:10.1155/2018/3128378
33
34 40. Pérez-Pevida B, Escalada J, Miras AD, et al. Mechanisms Underlying Type 2 Diabetes
35 Remission After Metabolic Surgery. *Front Endocrinol (Lausanne)* 2019;10:641.
36 doi:10.3389/fendo.2019.00641
37
38 41. Higgins JPT, Green S. *Cochrane Handbook for systematic reviews of interventions version*
39 5.1.0 (updated March 2011). The Cochrane collaboration, 2011. Available:
40 <http://www.handbook.cochrane.org/>
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

- 1
2
3
4 42. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of
5 evidence. *J Clin Epidemiol* 2011;64:401–406.doi:10.1016/j.jclinepi.2010.07.015
6
7
8 43. Gonzalez JS, Peyrot M, McCarl LA, et al. Depression and diabetes treatment nonadherence: a
9 meta-analysis. *Diabetes Care* 2008;31(12):2398–2403.doi:10.2337/dc08-1341
10
11 44. Trief PM, Foster NC, Chaytor N, et al. Longitudinal Changes in Depression Symptoms and
12 Glycemia in Adults With Type 1 Diabetes. *Diabetes Care* 2019;42(7):1194–1201.
13
14
15 doi:10.2337/dc18-2441
16
17 45. Novak M, Mucsi I, Rhee CM, et al. Increased Risk of Incident Chronic Kidney Disease,
18 Cardiovascular Disease, and Mortality in Patients With Diabetes With Comorbid Depression.
19
20
21
22
23
24 46. Lustman PJ, Anderson RJ, Freedland KE, et al. Depression and poor glycemic control: a
25 meta-analytic review of the literature. *Diabetes Care* 2000;23(7):934–942.
26
27
28 doi:10.2337/diacare.23.7.934
29
30 47. Roopan S, Larsen ER. Use of antidepressants in patients with depression and comorbid
31 diabetes mellitus: a systematic review. *Acta Neuropsychiatr* 2017;29(3):127–139.
32
33
34
35
36 48. Croom AM. Music, neuroscience, and the psychology of well-being: a précis. *Front Psychol*
37 2012;2:393.doi:10.3389/fpsyg.2011.00393
38
39 49. Abd-Elshafy SK, Khalaf GS, Abo-Kerisha MZ, et al. Not All Sounds Have Negative Effects
40 on Children Undergoing Cardiac Surgery. *J Cardiothorac Vasc Anesth* 2015;29(5):1277–
41
42
43 1284.doi:10.1053/j.jvca.2015.01.005
44
45 50. Yu SJ, Sun ZY, Zhou J, et al. Exploration of music behavioral intervention in the treatment of
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

Identification

Screening

Eligibility

Included

Records identified through Searching (n=?)
 Wanfang (n=?) CNKI (n=?)
 CBM (n=?) VIP (n=?)
 EMBASE (n=?) ChiCTR (n=?)
 PubMed (n=?) Cochrane Library (n=?)
 ClinicalTrials.gov (n=?)

Records identified through other sources (n=?)

Duplicate records excluded (n=?)


Titles and abstracts screened (n=?)

Records excluded with reasons (n=?)
 -Not RCTs (n=?)
 -With serious complications (n=?)
 -Improper Interventions (n=?)
 -Improper comparisons (n=?)
 ...

Full text articles assessed for eligibility (n=?)

Records excluded with reasons (n=?)
 -Not RCTs (n=?)
 -With serious complications (n=?)
 -Improper Interventions (n=?)
 -Improper comparisons (n=?)
 -No statistical indicator/not original data (n=?)
 ...

RCTs included (n=?)



PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification P₁	1a	Identify the report as a protocol of a systematic review
Update None	1b	If the protocol is for an update of a previous systematic review, identify as such
Registration P₂	2	If registered, provide the name of the registry (such as PROSPERO) and registration number
Authors:		
Contact P₁	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author
Contributions P₁₁	3b	Describe contributions of protocol authors and identify the guarantor of the review
Amendments None	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments
Support:		
Sources P₁₂	5a	Indicate sources of financial or other support for the review
Sponsor None	5b	Provide name for the review funder and/or sponsor
Role of sponsor or funder None	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol
INTRODUCTION		
Rationale P₄₋₅	6	Describe the rationale for the review in the context of what is already known
Objectives P₆₋₇	7	Provide an explicit statement of the question(s) the review will address with reference to participants P₆ , interventions P₇ , comparators P₇ , and outcomes P₇ (PICO)
METHODS		
Eligibility criteria P₅	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review
Information sources P₆, P₇, P₉	9	Describe all intended information sources (such as electronic databases P₇ , contact with study authors P₉ , trial registers P₆ or other grey literature sources P₇) with planned dates of coverage P₇
Search strategy P₈	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated
Study records:		

Data management P ₉	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review
Selection process P ₈₋₉	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)
Data collection process P ₉	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators
Data items P ₉	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications
Outcomes and prioritization P ₇	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale
Risk of bias in individual studies P ₉	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis
Data synthesis P ₉₋₁₀	15a	Describe criteria under which study data will be quantitatively synthesised P ₉₋₁₀
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I ² , Kendall's τ) P ₉₋₁₀
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) P ₁₀
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned P ₁₀
Meta-bias(es) P ₁₀	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)
Confidence in cumulative evidence P ₁₀	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.

BMJ Open

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-036268.R1
Article Type:	Protocol
Date Submitted by the Author:	28-Jan-2020
Complete List of Authors:	Zhou, Lin-yue ; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Zhang, Yuan ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Tian, Yuan; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Fu, Xiaoxu; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Wang, Li-zhen ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Xie, Chun-guang; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine
Primary Subject Heading:	Diabetes and endocrinology
Secondary Subject Heading:	Mental health, Complementary medicine
Keywords:	DIABETES & ENDOCRINOLOGY, MENTAL HEALTH, COMPLEMENTARY MEDICINE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Lin-yue Zhou,^{1,2} Yuan Zhang,² Yuan Tian,² Xiaoxu Fu,^{1,2} Li-zhen Wang,² Chun-guang Xie*^{1,2}

Author affiliations

¹Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Chengdu, Sichuan Province, China

²School of Clinical Medicine, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan Province, China

*Correspondence to:

Corresponding Author: Professor Chun-guang Xie

Address: Department of Endocrinology, Hospital of Chengdu University of Traditional Chinese Medicine, No 39 Shi-er-qiao Road, Chengdu, Sichuan Province, China, 610072

Tel: (86)13438826824

E-mail: killuya233@stu.cdutcm.edu.cn

Word count:2745 words (excluding title page, abstract, references, figures and tables)

ABSTRACT

Introduction About 463 million adults aged 20 to 79 have diabetes globally and mental disorders often exist in diabetic patients as comorbidities. The comorbidities can lead to deterioration of diseases, increased difficulty in treatment, even as well as the great mortality. Music intervention has been applied in the treatment of the comorbidities for 12 years, but there are not recommendations due to lacking of evidence. Thus, a meta-analysis is necessary to be conducted to evaluate the effect of music intervention.

Methods and analysis We will search the following nine online electronic databases: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical Database (CBM), from their inception until the present. We also plan to search other relevant resources, including grey literatures and the reference lists of relevant publications. Only randomized controlled trials (RCTs) of music intervention for depression or anxiety with diabetes will be involved. The primary outcomes include depression score and anxiety score measured by scales, and the secondary outcome is safety. Data extraction will be implemented by two researchers independently. Risk of bias will be evaluated through Cochrane Collaboration's Risk of Bias tool. Eventually, all the data will be analyzed via the software Review Manager V.5.3.

Ethics and dissemination This meta-analysis will provide information about music intervention applied to treat depression or anxiety in diabetic patients. There is no required ethical approval because this is a protocol for meta-analysis. The results of this study will be published to a peer reviewed journal.

Trial registration number CRD42019146439

Strengths and limitations of this study

· This meta-analysis will provide evidence of the effectiveness and safety of music intervention applied to treat depression or anxiety in diabetic patients for the first time.

· This systematic review will furnish evidence reference for guideline of relevant areas.

· This meta-analysis will be implemented on the basis of rigorous international gold-standard methodology and integrated search strategy.

· A language bias may exist in this meta-analysis because it will include studies only in English and Chinese on account of the language barrier.

INTRODUCTION

Description of the condition

According to the newest “IDF Diabetes Atlas (9th edition)” published by International Diabetes Federation (IDF), about 463 million adults aged 20 to 79 have diabetes globally (1 in 11).[1] It is well known that diabetes has a significant influence on life quality because it can cause a lot of complications such as retinopathy, renal disease, cardiovascular disease, neurological disease and foot ulcers which may result in lower extremity amputation.[2]

In addition to physical complications, some mental disorders often exist in diabetic patients as comorbidities. Some studies have indicated that depression is more prevalent in diabetic patients compared to those without this disease.[3,4] At the same time, depression could also increase the risk of incidence of diabetes,[5-7] which suggests that depression has a bidirectional association with diabetes. And the relation between anxiety and diabetes has been confirmed to be the same. People with diabetes are more susceptible to concurrent anxiety[8,9] while reversely anxiety is a risk factor of diabetic incidents.[10] Individuals with chronic diseases and also comorbid depression or anxiety have obviously higher health-care utilization, functional disability, work absence and mortality risk than those without them.[11,12]

Description of the intervention

When it comes to the treatment of diabetes with depression or anxiety, conventional glucose-lowering treatment combined with drug therapy or psychological intervention are the routine treatments.[13,14] Music intervention is also a successful therapy using music for treatment by professionals, which can be divided into two major categories, namely music therapy and music medicine.[15] The official definition of music therapy is the “...clinical and evidence-based use of music interventions to accomplish individualized goals within a therapeutic relationship by a credentialed professional who has completed an approved music therapy program”.[16] A trained music therapist leads the treatment process, choosing the type of music, the form of music presentation (pre-recorded or live), and the treatment setting (singing, playing instruments,

1
2
3
4 composing, etc.) based on the patient's condition, ultimately resulting in a therapeutic scheme that
5 is best for the patient. By comparison, music medicine is defined as the passive listening to pre-
6 recorded music, which can be implemented independently by medical or healthcare professionals
7 instead of music therapists.[17] In brief, music therapy focuses on active music therapy and
8 personalized treatments while music medicine belongs to passive music listening and non-
9 personalized therapy. It is significant to emphasize the distinction between music therapy and music
10 medicine because some literatures suggest that the effect of the former is better than the latter in a
11 great many outcomes.[18-19]

12
13
14
15
16
17
18
19 A lot of studies have suggested that music, no matter what type of it is, demonstrates a remarkable
20 effect in alleviating depression and anxiety among patients with various diseases.[20-22] As for
21 diabetes, recent studies have showed that music intervention could improve fasting blood-glucose
22 and systolic blood pressure in diabetic patients.[23,24] Good control of blood glucose, blood lipids
23 and blood pressure can effectively reduce the incidence of serious complications of diabetes.[25]

30 31 **How the intervention might work**

32
33 Researches about music psychology have shown the special potential of music. Music, as a stimulus,
34 can not only trigger a variety of emotional experiences,[26, 27] but also strengthen ability to deal
35 with negative emotions in everyday life.[28] The proposed mechanisms of music intervention on
36 depression and anxiety span several domains. Firstly, music intervention can exert
37 neurophysiological and neurochemical effects, such as triggering endorphin release[29] and
38 stimulating cognitive functioning.[30] Secondly, music intervention can promote positive self-
39 emotions including enhancing the sense of happiness[31] and activating self-efficacy and self-
40 confidence through experiences of success.[32] Lastly, music intervention can build up sociability,
41 such as interaction skills[33] and interpersonal relationship.[34] Furthermore, there are published
42 studies indicating that music intervention combined with diabetic routine treatments could improve
43 depression score, anxiety score, fasting glucose and glycosylated hemoglobin (HbA1c).[35,36]

Why it is important to do this review

From the above, many studies have indicated the bidirectional relationship between diabetes and mental disorders. Music intervention, as a therapeutic method, has been applied to alleviate depression and anxiety and improve related indicators of diabetes. By searching literatures, it can be found that some reviews evaluated the effect of music intervention on patients with mental disorders,[37-39] while no related systematic review and meta-analysis can be searched about the effect of music intervention on mental health of patients with diabetes. Therefore, if music intervention can be proved to be both valid and with few side effects through this study, it is worth being promoted and applicated in clinic all over the world to benefit more people since this method is easy to be operated.

Objectives

The purpose of this review is to systematically assess the effect of music intervention and compare the effects between music therapy and music medicine on depression and anxiety of diabetic patients. Additionally, it is also our expectation to furnish evidence reference for related guideline.

METHODS

Study registration

The protocol of the meta-analysis has been registered in International Prospective Register of Systematic Reviews (PROSPERO), and CRD42019146439 is the registration ID. This protocol has been planned severely on the basis of Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P)[40] statement guidelines. The results will be reported in conformity to the PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Healthcare Interventions.[41]

Eligibility criteria

We will severely screen studies that have compliance with following inclusion criteria.

Type of studies

We will only cover randomized controlled trials (RCTs) in English or Chinese, excepting Quasi-RCTs and cluster RCTs. Experiments on animals and nonrandomized clinical trials will be out of our consideration. Articles with significant overlap with other articles through print or e-publishing will fail to be included. Duplicate publications, that is results originated from a single experiment but measured by other standards and then published as disparate papers, will also be excluded. Publications with originality is our first choice. The publication status will not be restricted.

Type of Participants

Diabetic Patients with depression or anxiety symptoms, regardless of the type of diabetes and no restriction on gender, age, race, economic ability and education level. Patients of other disease or healthy persons will be excluded.

Type of interventions and controls

Interventions include music intervention (music therapy or music medicine) alone or music intervention in combination with other interventions. No restriction on the type of music. Other interventions refer to: (1) routine treatments for diabetes which mainly include pharmacologic approaches to glycemetic treatment, diabetes self-management education and support (DSMES),, etc. according to the newest guideline of American Diabetes Association (ADA);[42,43] (2) other therapies such as acupoint therapy,[44] metabolic surgery,[45] etc. When intervention is music intervention alone, controls can be no treatment, placebo or other active therapies. Notably, when intervention is music intervention combined with active therapies, the control must be the same active therapies.

Type of outcome measures

This review aims at systematically assessing the effect of music intervention on depression or

1
2
3
4 anxiety of diabetic patients, so we choose depression score and anxiety score as the primary
5
6 outcomes, which can give an index to the degree of depression and anxiety by scales.

7
8 Primary outcomes: Depression score and anxiety score. Depression score is measured using self-
9
10 rating depression scale (SDS), Hamilton depression scale (HAMD) or other validated scales for
11
12 depression. Anxiety score is measured using self-rating anxiety scale (SAS), Hamilton anxiety
13
14 scale (HAMA) or other validated scales for anxiety.

15
16 Secondary outcomes: Safety. Measurement via incidence of adverse effects.
17
18

19 **Search methods**

20 Search resources

21
22 We will search the following nine online electronic databases: PubMed, Web of Science, Embase,
23
24 EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang
25
26 Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical
27
28 Database (CBM), from their inception to the present. We also plan to search other relevant
29
30 resources as far as possible, including: (1) grey literatures (like conference literatures and
31
32 dissertations) from the following four sources: OpenGrey (www.opengrey.eu/), CNKI
33
34 (www.cnki.net), Open Access Theses and Dissertations (oatd.org) and British Library EThOS
35
36 (ethos.bl.uk/); (2) the reference lists of relevant publications for additional studies.
37
38
39

40 Search strategies

41
42 The search strategy will include the MeSH terms for diabetes mellitus, depression, anxiety, music
43
44 therapy, music and singing, combined with their respective free-text terms. The search details of
45
46 the PubMed are simply shown in table 1.
47
48
49
50
51
52
53
54
55
56

Table 1 Example of PubMed search strategy.

Number	Search terms
#1	"Diabetes Mellitus"[Mesh]
#2	"Depression"[Mesh] OR Depressions[Title/Abstract] OR Depressive Symptoms[Title/Abstract] OR Depressive Symptom[Title/Abstract] OR Symptom, Depressive[Title/Abstract] OR Symptoms, Depressive[Title/Abstract] OR Emotional Depression[Title/Abstract] OR Depression, Emotional[Title/Abstract] OR Depressions, Emotional[Title/Abstract] OR Emotional Depressions[Title/Abstract]
#3	"Anxiety"[Mesh] OR Hypervigilance [Title/Abstract] OR Nervousness [Title/Abstract]
#4	#2 OR #3
#5	"Music Therapy"[Mesh] OR Therapy, Music [Title/Abstract]
#6	"Music"[Mesh] OR music intervention[Title/Abstract] OR music medicine[Title/Abstract] OR music listening[Title/Abstract] OR audio recording[Title/Abstract] OR mp3 hearing[Title/Abstract]
#7	"Singing"[Mesh] OR sing[Title/Abstract] OR sings[Title/Abstract] OR song[Title/Abstract] OR compose[Title/Abstract] OR composing[Title/Abstract]
#8	#5 OR #6 OR #7
#9	randomized controlled trial [Publication Type] OR randomized [Title/Abstract] OR placebo [Title/Abstract]
#10	#1 AND #4 AND #8 AND #9

Data collection and analysis

Studies selection

Firstly, two researchers (L-yZ and YT) will respectively screen the keywords, titles and abstracts of all obtained studies using endnote x9 software and determine whether trials have compliance with inclusion criteria. Secondly, we will also get access to the full text of all studies of possible relevance for further assessment. Each of excluded study will be noted with reasons. Any disagreements will be settled through consensus or referral to the third researcher (YZ). The

1
2
3
4 specific selection steps are presented in the figure 1.
5
6

7 Data extraction

8
9 Two researchers (L-yZ and YT) will independently extract and tabulate the following data items
10 via Microsoft Excel:
11

- 12 1. General information (including publication year and name of the first author).
- 13 2. Study design (including random method, allocation concealment and blinding).
- 14 3. Participants (including sample size, gender, age, area, depression score, anxiety score and course
15 of disease).
- 16 4. Treatment (including interventions, controls, music type and the course of treatment).
- 17 5. Outcomes (including depression score, anxiety score and safety).

18
19 If there is any different opinion, the final decision will go to the third researcher (X-xF) to be
20 made.
21
22

23 Assessment of risk of bias

24
25 The Cochrane Collaboration's Risk of Bias tool[46] will be applied to evaluate the risk of bias of
26 all included RCT studies by two authors (L-zW and YZ). The following domains will be assessed:
27 sequence generation, allocation concealment, blinding of participants and personnel, blinding of
28 outcome assessors, incomplete outcome data, selective reporting and other sources of bias. The
29 assessments will be classified into "low", "unclear" or "high" of risk.
30
31

32 Measures of treatment effect

33
34 Considering that all outcomes (depression score, anxiety score and safety) are continuous variables,
35 the mean difference (MD) and 95% confidence interval (CI) will be calculated when the
36 quantitative data are from the same measurement method or small data scale. On the contrary, the
37 standardized mean difference (SMD) and 95% CI will be calculated.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Unit of analysis issues

For the standard research which has one experimental group with one control group, the unit of analysis is based on the individual. For the multi-arm study which has more than two intervention groups, the recommended method is to create a single pair-wise comparison through combining groups.

Dealing with missing data

If the required data are unavailable from literatures, one researcher (YZ) will send an e-mail to the corresponding author and ask for further information.

Assessment of heterogeneity

A meta-analysis will be implemented if the included studies are adequately homogenous. *P*-value and *I*² will be used to assess the heterogeneity of statistic. When *P*-value >0.05 and *I*² ≤50%, the heterogeneity is regarded as acceptable. When *P*-value <0.05 or *I*² >50%, the heterogeneity is considered as significant.

Assessment of reporting bias

Publication bias will be implemented, if there are more than 10 studies included, through a funnel plot.

Data synthesis

RevMan 5.3. software provided by the Cochrane collaboration will be used to analyze the data. If the included studies are adequately homogenous, fixed-effects model will be used for statistical combination. Since all outcomes are continuous variables, the mean difference (MD) and 95% confidence interval (CI) will be calculated when the outcomes are from the same measurement method or small data scale. On the contrary, the standardized mean difference (SMD) and 95% CI will be calculated. If there is significant heterogeneity among the included studies, subgroup

1
2
3 analyses or sensitivity analyses will be performed. If a meta-analysis is unavailable, descriptive
4 summaries will be provided.
5
6
7

8 9 Subgroup analysis

10 The subgroup analyses are planned as follows:

- 11 1. Type of music intervention (music therapy or music medicine).
 - 12 2. Type of scales (SDS, HAMD, SAS, HAMA or other validated scales for depression or anxiety).
 - 13 3. Different age groups.
- 14
15
16
17
18
19
20

21 Sensitivity analysis

22 Sensitivity analyses will be performed through including or excluding a specific study, such as a
23 study without a clear allocation concealment or randomization method, to observe whether the
24 results changed.
25
26
27
28
29
30

31 Patient and public involvement

32 Patients and public involvement will fail to be included in this study.
33
34
35
36

37 Evidence assessed

38 GRADE guideline will be used to evaluate the quality of evidence.[47] To fulfil transparency and
39 simplification, the quality of evidence can be assessed by 4 levels: “high”, “medium”, “low” and
40 “very low”. GRADE profiler 3.2 will be applied for analysis.
41
42
43
44
45

46 **DISCUSSION**

47 Depression and anxiety often occur independently or simultaneously during the development of
48 many chronic diseases. In recent years, the number of studies about comorbid depression or anxiety
49 in diabetics are continually increasing, not only because the two-way relationship between
50 depression/anxiety and diabetes can lead to deterioration of diseases, increased difficulty in
51
52
53
54
55
56

1
2
3
4 treatment, even as well as the great mortality,[48-51] but also because therapeutic methods for
5 treating the comorbidities are limited and unsatisfactory. On the one hand, some antidepressant
6 drugs, such as nortriptyline and sertraline, may complicate glycemic control.[52] On the other hand,
7 psychotherapy based on verbal communication requires a professional psychiatrist and it is
8 expensive.
9
10
11
12

13 Music intervention has been proved by many literatures to have a beneficial effect on depression
14 and anxiety, such as strengthening awareness and sensitiveness for positive emotions.[53]
15 Noticeably, a few studies also show that music intervention can reduce blood glucose and HbA1c
16 simultaneously.[24,54] Therefore, music intervention is an applicable method for treating
17 depression and anxiety with diabetes. Additionally, music intervention, almost no side effects, is
18 safer than pharmacotherapy, easier to be operated and costs less than psychotherapy based on verbal
19 communication, so it is conducive to clinical promotion and application.
20
21
22
23
24
25
26

27 Since 2007,[55] music intervention has been applied in the treatment of depression and anxiety
28 with diabetes for 12 years. Nonetheless, there are not systematic review or meta-analysis conducted
29 to evaluate the potential benefits and harms of this therapy method. Thus, this study intends to
30 provide strong evidence for music intervention applied to treat mental disorders in diabetic patients
31 and synchronously furnish evidence reference for related guideline.
32
33
34
35
36

37 However, a language bias may exist in this meta-analysis because it will include studies only in
38 English and Chinese because of the language barrier. Therefore, some related studies from
39 medical databases in other languages might be missed.
40
41
42

43 In conclusion, this meta-analysis will be conducive to assess underlying benefits and harms of
44 music intervention on comorbid depression or anxiety with diabetes. Moreover, the results of this
45 study might not merely furnish reference basis for related guideline, but also may attract more
46 attention on music intervention, which can be a chance to facilitate its clinical application and
47 ultimately benefit more patients.
48
49
50
51
52
53

54 **ETHICS AND DISSEMINATION**

55
56
57
58
59
60

1
2
3
4 There is no required ethical approval because this is a protocol for meta-analysis. The results of
5 this study will be published to a peer reviewed journal.
6
7

8 9 **Figure legend**

10
11 **Figure 1** Flowchart to present the search process.
12
13

14
15 **Contributions** L-yZ and C-gX contributed to the conception of the study. The manuscript of the
16 protocol was drafted by L-yZ, and was revised by XxF and C-gX. L-yZ and YT will also
17 independently screen the potential studies and extract data from included studies. L-zW and YZ
18 will assess the risk of bias and accomplish data synthesis. XxF will arbitrate any disagreements.
19 C-gX will ensure that no errors occur during the review. All authors have approved the
20 publication of the protocol.
21
22

23
24 **Funding** This study was supported by grants from the Science and Technology Project of
25 Sichuan Province (2019YFS002), the Special Scientific and Technological Research Project of
26 Administration of TCM of Sichuan Province (2018KF003), and the Foundation of Development
27 of Science and Technology of Chengdu University of TCM (2016-D-YY-61).
28
29

30
31 **Competing interests** None.
32
33

34
35 **Patient consent for publication** Not required.
36
37

38
39 **Provenance and peer review** Not commissioned; externally peer reviewed.
40
41

42
43 **Open access** This is an open access article distributed in accordance with the Creative Commons
44 Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix,
45 adapt, build upon this work non-commercially, and license their derivative works on different terms,
46 provided the original work is properly cited, appropriate credit is given, any changes made
47 indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.
48
49

50 51 **References**

52
53
54 1. International Diabetes Federation. IDF Diabetes Atlas (9th edition), 2019. Available:
55
56

- 1
2
3
4 <http://www.diabetesatlas.org/>. Accessed November 9, 2019
- 5
6 2. World Health Organization. Global report on diabetes, 2014. Available:
7 <https://www.who.int/news-room/fact-sheets/detail/diabetes>. Accessed November 9, 2019
- 8
9
10 3. Ali S, Stone MA, Peters JL, et al. The prevalence of co-morbid depression in adults with
11 Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* 2006;23(11):1165–
12 1173. doi:10.1111/j.1464-5491.2006.01943.x
- 13
14
15 4. Anderson RJ, Freedland KE, Clouse RE, et al. The prevalence of comorbid depression in
16 adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24(6):1069–1078.
17
18 doi:10.2337/diacare.24.6.1069
- 19
20
21 5. Knol MJ, Twisk JW, Beekman AT, et al. Depression as a risk factor for the onset of type 2
22 diabetes mellitus. A meta-analysis. *Diabetologia* 2006;49(5):837–845.doi:10.1007/s00125-
23 006-0159-x
- 24
25
26
27 6. Cosgrove MP, Sargeant LA, Griffin SJ. Does depression increase the risk of developing type
28 2 diabetes?. *Occup Med (Lond)* 2008;58(1):7–14.doi:10.1093/occmed/kqm105
- 29
30
31 7. Mezuk B, Eaton WW, Albrecht S, et al. Depression and type 2 diabetes over the lifespan: a
32 meta-analysis. *Diabetes Care* 2008;31(12):2383–2390.doi:10.2337/dc08-0985
- 33
34
35 8. Grigsby AB, Anderson RJ, Freedland KE, et al. Prevalence of anxiety in adults with diabetes:
36 a systematic review. *J Psychosom Res* 2002; 53: 1053–1060.doi:10.1016/s0022-
37 3999(02)00417-8
- 38
39
40 9. Smith KJ, Beland M, Clyde M, et al. Association of diabetes with anxiety: a systematic
41 review and metaanalysis. *J Psychosom Res* 2013; 74: 89–99.
42
43 doi:10.1016/j.jpsychores.2012.11.013
- 44
45
46 10. Smith KJ, Deschênes SS, Schmitz N. Investigating the longitudinal association between
47 diabetes and anxiety: a systematic review and meta-analysis. *Diabet Med* 2018;35(6):677–
48 693.doi:10.1111/dme.13606
- 49
50
51
52 11. Egede LE. Major depression in individuals with chronic medical disorders: prevalence,
53
54 correlates and association with health resource utilization, lost productivity and functional
55
56

- 1
2
3
4 disability. *Gen Hosp Psychiatry* 2007;29:409–416.doi:10.1016/j.genhosppsy.2007.06.002
- 5
6 12. Naicker K, Johnson JA, Skogen JC, et al. Type 2 Diabetes and Comorbid Symptoms of
7
8 Depression and Anxiety: Longitudinal Associations With Mortality Risk. *Diabetes Care*
9
10 2017;40(3):352–358.doi:10.2337/dc16-2018
- 11
12 13. van der Feltz-Cornelis CM, Nuyen J, Stoop C, et al. Effect of interventions for major
13
14 depressive disorder and significant depressive symptoms in patients with diabetes mellitus:
15
16 a systematic review and meta-analysis. *Gen Hosp Psychiatry* 2010;32(4):380–395.
17
18 doi:10.1016/j.genhosppsy.2010.03.011
- 19
20 14. Komsuoglu Celikyurt I, Mutlu O, Ulak G, et al. Exenatide treatment exerts anxiolytic- and
21
22 antidepressant-like effects and reverses neuropathy in a mouse model of type-2 diabetes. *Med*
23
24 *Sci Monit Basic Res* 2014;20:112–117.doi:10.12659/MSMBR.891168
- 25
26 15. Leubner D, Hinterberger T. Reviewing the Effectiveness of Music Interventions in Treating
27
28 Depression. *Front Psychol* 2017;8:1109.doi:10.3389/fpsyg.2017.01109
- 29
30 16. American Music Therapy Association. What is music therapy?, 2013.Available:
31
32 <https://www.musictherapy.org/about/musictherapy/>. Accessed January 9, 2020.
- 33
34 17. Dileo C. A classification model for music and medicine. National Association of Music
35
36 Therapy, Washington, DC, 1999: 1-6.
- 37
38 18. Raglio A, Bellelli G, Mazzola P, et al. Music, music therapy and dementia: a review of
39
40 literature and the recommendations of the Italian Psychogeriatric Association. *Maturitas*
41
42 2012;72(4):305–310.doi:10.1016/j.maturitas.2012.05.016
- 43
44 19. Dileo C. Effects of music and music therapy on medical patients: a meta-analysis of the
45
46 research and implications for the future. *J Soc Integr Oncol* 2006;4(2):67–70.
47
48 doi:10.2310/7200.2006.002
- 49
50 20. Thaut MH, Gardiner JC, Holmberg D, et al. Neurologic music therapy improves executive
51
52 function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*
53
54 2009;1169:406–416.doi:10.1111/j.1749-6632.2009.04585.x
- 55
56 21. Kim DS, Park YG, Choi JH, et al. Effects of music therapy on mood in stroke patients.

- 1
2
3
4 *Yonsei Med J* 2011;52(6):977–981.doi:10.3349/ymj.2011.52.6.977
- 5
6 22. Nwebube C, Glover V, Stewart L. Prenatal listening to songs composed for pregnancy and
7 symptoms of anxiety and depression: a pilot study. *BMC Complement Altern Med*
8 2017;17(1):256.doi:10.1186/s12906-017-1759-3
- 9
10
11 23. Lin XX, Ji JM, Zhou GY. Influence of listening to Jiao tune music on sleep quality and blood
12 glucose in the elderly patients with type 2 diabetes accompanied by insomnia. *Journal of*
13 *rehabilitation* 2017;27(01):44-48.
- 14
15
16 24. Mandel SE, Davis BA, Secic M. Effects of music therapy and music-assisted relaxation and
17 imagery on health-related outcomes in diabetes education: a feasibility study. *Diabetes Educ*
18 2013;39(4):568–581.doi:10.1177/0145721713492216
- 19
20
21 25. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2014.
22 Available: [https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf)
23 [burden-in-the-united-states.pdf](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf). Accessed November 9, 2019
- 24
25
26 26. Gabrielsson A, Wik SL. Strong experiences related to music: a descriptive system. *Music Sci*
27 2003;7(2):157–217.
- 28
29
30 27. Zentner M, Grandjean D, Scherer KR. Emotions evoked by the sound of music:
31 characterization, classification, and measurement. *Emotion* 2008;8(4):494–521.
32 doi:10.1037/1528-3542.8.4.494
- 33
34
35 28. Saarikallio S. Music as emotional self-regulation throughout adulthood. *Psychol Music*
36 2011;39(3):307–327.
- 37
38
39 29. Gök Ugur H, Yaman Aktaş Y, Orak OS, et al. The effect of music therapy on depression and
40 physiological parameters in elderly people living in a Turkish nursing home: a randomized-
41 controlled trial. *Aging Ment Health* 2017;21(12):1280–1286.
42 doi:10.1080/13607863.2016.1222348
- 43
44
45 30. Gómez Gallego M, Gómez García J. Music therapy and Alzheimer's disease: Cognitive,
46 psychological, and behavioural effects. *Neurologia* 2017;32(5):300–308.
47 doi:10.1016/j.nrl.2015.12.003
- 48
49
50
51
52
53
54
55
56
57
58
59
60


- 1
2
3
4 31. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement
5 intervention on the cognitive functions of people with moderate dementia: a randomized
6 controlled trial. *Aging Ment Health* 2018;22(3):306–315.
7
8 doi:10.1080/13607863.2016.1251571
9
10
11 32. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in
12 elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging*
13 *Ment Health* 2017;21(2):156–162.doi:10.1080/13607863.2015.1093602
14
15
16
17 33. Bieleninik L, Geretsegger M, Mössler K, et al. Effects of Improvisational Music Therapy vs
18 Enhanced Standard Care on Symptom Severity Among Children With Autism Spectrum
19 Disorder: The TIME-A Randomized Clinical Trial. *JAMA* 2017;318(6):525–535.
20
21
22 doi:10.1001/jama.2017.9478
23
24
25 34. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a
26 controlled randomized study. *Neurol Sci* 2017;38(8):1485–1493.doi:10.1007/s10072-017-
27 3003-9
28
29
30
31 35. Deng HX, Gao GS, Li ZM, et al. Application of receiving music therapy in elderly patients
32 with diabetes. *Chinese journal of health and nutrition (1st Ten-days of month)*
33 2013,23(3):1186-1186.
34
35
36
37 36. Zhang ZW. Application of music therapy and psychotherapy adjuvant therapy in type 2
38 diabetes. *Chinese medical science* 2011,01(18):150-151.
39
40
41 37. Solanki MS, Zafar M, Rastogi R. Music as a therapy: role in psychiatry. *Asian J Psychiatr*
42 2013;6(3):193–199.doi:10.1016/j.ajp.2012.12.001
43
44
45 38. Petrovsky D, Cacchione PZ, George M. Review of the effect of music interventions on
46 symptoms of anxiety and depression in older adults with mild dementia. *Int Psychogeriatr*
47 2015;27(10):1661–1670.doi:10.1017/S1041610215000393
48
49
50 39. Aalbers S, Fusar-Poli L, Freeman RE, et al. Music therapy for depression. *Cochrane*
51 *Database Syst Rev* 2017;11(11):CD004517. Published 2017 Nov 16.
52
53
54 doi:10.1002/14651858.CD004517.pub3
55
56
57
58
59
60

- 1
2
3
4 40. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and
5 meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*
6 2015;350:g7647.doi:10.1136/bmj.g7647
7
8
9
10 41. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic
11 reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and
12 elaboration. *BMJ* 2009;339:b2700.doi:10.1136/bmj.b2700
13
14
15 42. American Diabetes Association. 9. Pharmacologic Approaches to Glycemic Treatment:
16 Standards of Medical Care in Diabetes-2020. *Diabetes Care* 2020;43(Suppl 1):S98–S110.
17 doi:10.2337/dc20-S009
18
19
20
21 43. American Diabetes Association. 5. Facilitating Behavior Change and Well-being to Improve
22 Health Outcomes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*
23 2020;43(Suppl 1):S48–S65. doi:10.2337/dc20-S005
24
25
26
27 44. Feng Y, Fang Y, Wang Y, et al. Acupoint Therapy on Diabetes Mellitus and Its Common
28 Chronic Complications: A Review of Its Mechanisms. *Biomed Res Int* 2018;2018:3128378.
29 doi:10.1155/2018/3128378
30
31
32
33 45. Pérez-Pevida B, Escalada J, Miras AD, et al. Mechanisms Underlying Type 2 Diabetes
34 Remission After Metabolic Surgery. *Front Endocrinol (Lausanne)* 2019;10:641.
35 doi:10.3389/fendo.2019.00641
36
37
38
39 46. Higgins JPT, Thomas J, Chandler J, et al. Cochrane Handbook for Systematic Reviews of
40 Interventions version 6.0 (updated July 2019). Cochrane, 2019. Available:
41 www.training.cochrane.org/handbook.
42
43
44
45 47. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of
46 evidence. *J Clin Epidemiol* 2011;64:401–406.doi:10.1016/j.jclinepi.2010.07.015
47
48
49 48. Gonzalez JS, Peyrot M, McCarl LA, et al. Depression and diabetes treatment nonadherence: a
50 meta-analysis. *Diabetes Care* 2008;31(12):2398–2403.doi:10.2337/dc08-1341
51
52
53 49. Trief PM, Foster NC, Chaytor N, et al. Longitudinal Changes in Depression Symptoms and
54 Glycemia in Adults With Type 1 Diabetes. *Diabetes Care* 2019;42(7):1194–1201.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

doi:10.2337/dc18-2441

50. Novak M, Mucsi I, Rhee CM, et al. Increased Risk of Incident Chronic Kidney Disease, Cardiovascular Disease, and Mortality in Patients With Diabetes With Comorbid Depression. *Diabetes Care* 2016;39(11):1940–1947. doi:10.2337/dc16-0048
51. Lustman PJ, Anderson RJ, Freedland KE, et al. Depression and poor glycemic control: a meta-analytic review of the literature. *Diabetes Care* 2000;23(7):934–942. doi:10.2337/diacare.23.7.934
52. Roopan S, Larsen ER. Use of antidepressants in patients with depression and comorbid diabetes mellitus: a systematic review. *Acta Neuropsychiatr* 2017;29(3):127–139. doi:10.1017/neu.2016.54
53. Croom AM. Music, neuroscience, and the psychology of well-being: a précis. *Front Psychol* 2012;2:393. doi:10.3389/fpsyg.2011.00393
54. Abd-Elshafy SK, Khalaf GS, Abo-Kerisha MZ, et al. Not All Sounds Have Negative Effects on Children Undergoing Cardiac Surgery. *J Cardiothorac Vasc Anesth* 2015;29(5):1277–1284. doi:10.1053/j.jvca.2015.01.005
55. Yu SJ, Sun ZY, Zhou J, et al. Exploration of music behavioral intervention in the treatment of diabetes mellitus. *Hebei med* 2007;13(8):903-906.



PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification P₁	1a	Identify the report as a protocol of a systematic review
Update None	1b	If the protocol is for an update of a previous systematic review, identify as such
Registration P₂	2	If registered, provide the name of the registry (such as PROSPERO) and registration number
Authors:		
Contact P₁	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author
Contributions P₁₄	3b	Describe contributions of protocol authors and identify the guarantor of the review
Amendments None	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments
Support:		
Sources P₁₄	5a	Indicate sources of financial or other support for the review
Sponsor None	5b	Provide name for the review funder and/or sponsor
Role of sponsor or funder None	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol
INTRODUCTION		
Rationale P₄₋₆	6	Describe the rationale for the review in the context of what is already known
Objectives P₇₋₈	7	Provide an explicit statement of the question(s) the review will address with reference to participants P₇ , interventions P₇ , comparators P₇ , and outcomes P₇₋₈ (PICO)
METHODS		
Eligibility criteria P₆₋₇	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review
Information sources P₆, P₈, P₁₁	9	Describe all intended information sources (such as electronic databases P₈ , contact with study authors P₁₁ , trial registers P₆ or other grey literature sources P₈) with planned dates of coverage P₈
Search strategy P₉	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated
Study records:		

Data management P₁₀	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review
Selection process P₉₋₁₀	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)
Data collection process P₁₀	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators
Data items P₁₀	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications
Outcomes and prioritization P₇₋₈	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale
Risk of bias in individual studies P₁₀	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis
Data synthesis P₁₀₋₁₂	15a	Describe criteria under which study data will be quantitatively synthesised P₁₁
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I ² , Kendall's τ) P₁₀₋₁₁
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) P₁₂
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned P₁₂
Meta-bias(es) P₁₁	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)
Confidence in cumulative evidence P₁₂	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.

BMJ Open

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-036268.R2
Article Type:	Protocol
Date Submitted by the Author:	01-Apr-2020
Complete List of Authors:	Zhou, Lin-yue ; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Zhang, Yuan ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Tian, Yuan; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Fu, Xiaoxu; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Wang, Li-zhen ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Xie, Chun-guang; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine
Primary Subject Heading:	Diabetes and endocrinology
Secondary Subject Heading:	Mental health, Complementary medicine
Keywords:	DIABETES & ENDOCRINOLOGY, MENTAL HEALTH, COMPLEMENTARY MEDICINE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Effect of music intervention on mental health in patients with diabetes mellitus: a protocol for a systematic review and meta-analysis of randomized controlled trials

Lin-yue Zhou,^{1,2} Yuan Zhang,² Yuan Tian,² Xiaoxu Fu,^{1,2} Li-zhen Wang,² Chun-guang Xie*^{1,2}

Author affiliations

¹Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Chengdu, Sichuan Province, China

²School of Clinical Medicine, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan Province, China

*Correspondence to:

Corresponding Author: Professor Chun-guang Xie

Address: Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, No 39 Shi-er-qiao Road, Chengdu, Sichuan Province, China, 610075

Tel: (86)13438826824

E-mail: killuya233@stu.cdutcm.edu.cn

Word count:2771 words (excluding title page, abstract, references, figures and tables)

ABSTRACT

Introduction About 463 million adults aged 20 to 79 have diabetes globally and mental disorders often exist in diabetic patients as comorbidities. The comorbidities can lead to deterioration of diseases, increased difficulty in treatment, even as well as the great mortality. Music intervention has been applied in the treatment of the comorbidities for 12 years, but there are not recommendations due to lacking of evidence. Thus, a meta-analysis is necessary to be conducted to evaluate the effect of music intervention.

Methods and analysis We will search the following nine online electronic databases: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical Database (CBM), from their inception until March 2020. We also plan to search other relevant resources, including grey literatures and the reference lists of relevant publications. Only randomized controlled trials (RCTs) of music intervention for depression or anxiety with diabetes will be involved. The primary outcomes include depression score and anxiety score measured by scales, and the secondary outcome is safety. Data extraction will be implemented by two researchers independently. Risk of bias will be evaluated through Cochrane Collaboration's Risk of Bias tool. Eventually, all the data will be analyzed via the software Review Manager V.5.3.

Ethics and dissemination This meta-analysis will provide information about music intervention applied to treat depression or anxiety in diabetic patients. There is no required ethical approval because this meta-analysis is on the basis of published data. The results of this systematic review will be published to a peer reviewed journal.

Trial registration number CRD42019146439

Strengths and limitations of this study

· This meta-analysis will furnish evidence of the effectiveness and safety of music intervention applied to treat depression or anxiety in diabetic patients without precedent.

· On the basis of our comprehensive search strategy, the relevant original studies will fail to be omitted.

· Two authors will implement data extraction and estimation of the risk of bias separately.

· We will only cover randomized controlled trials (RCTs) in this study.

· A language bias may exist in this meta-analysis because it will include studies only in English and Chinese on account of the language barrier.

INTRODUCTION

Description of the condition

According to the newest “IDF Diabetes Atlas (9th edition)” published by International Diabetes Federation (IDF), about 463 million adults aged 20 to 79 have diabetes globally (1 in 11).[1] It is well known that diabetes has a significant influence on life quality because it can cause a lot of complications such as retinopathy, renal disease, cardiovascular disease, neurological disease and foot ulcers which may result in lower extremity amputation.[2]

In addition to physical complications, some mental disorders often exist in diabetic patients as comorbidities. Some studies have indicated that depression is more prevalent in diabetic patients compared to those without this disease.[3,4] At the same time, depression could also increase the risk of incidence of diabetes,[5-7] which suggests that depression has a bidirectional association with diabetes. And the relation between anxiety and diabetes has been confirmed to be the same. People with diabetes are more susceptible to concurrent anxiety[8,9] while reversely anxiety is a risk factor of diabetic incidents.[10] Individuals with chronic diseases and also comorbid depression or anxiety have obviously higher health-care utilization, functional disability, work absence and mortality risk than those without them.[11,12]

Description of the intervention

When it comes to the treatment of diabetes with depression or anxiety, conventional glucose-lowering treatment combined with drug therapy or psychological intervention are the routine treatments.[13,14] Music intervention is also a successful therapy using music for treatment by professionals, which can be divided into two major categories, namely music therapy and music medicine.[15] The official definition of music therapy is the “...clinical and evidence-based use of music interventions to accomplish individualized goals within a therapeutic relationship by a credentialed professional who has completed an approved music therapy program”.[16] A trained music therapist leads the treatment process, choosing the type of music, the form of music presentation (pre-recorded or live), and the treatment setting (singing, playing instruments,

1
2
3
4 composing, etc.) based on the patient's condition, ultimately resulting in a therapeutic scheme that
5 is best for the patient. By comparison, music medicine is defined as the passive listening to pre-
6 recorded music, which can be implemented independently by medical or healthcare professionals
7 instead of music therapists.[17] In brief, music therapy focuses on active music therapy and
8 personalized treatments while music medicine belongs to passive music listening and non-
9 personalized therapy. It is significant to emphasize the distinction between music therapy and music
10 medicine because some literatures suggest that the effect of the former is better than the latter in a
11 great many outcomes.[18-19]

12
13
14
15
16
17
18
19 A lot of studies have suggested that music, no matter what type of it is, demonstrates a
20 remarkable effect in alleviating depression and anxiety among patients with various diseases.[20-
21 22] As for diabetes, recent studies have showed that music intervention could improve fasting
22 blood-glucose and systolic blood pressure in diabetic patients.[23,24] Good control of blood
23 glucose, blood lipids and blood pressure can effectively reduce the incidence of serious
24 complications of diabetes.[25]

31 32 33 **How the intervention might work**

34
35 Researches about music psychology have shown the special potential of music. Music, as a stimulus,
36 can not only trigger a variety of emotional experiences,[26, 27] but also strengthen ability to deal
37 with negative emotions in everyday life.[28] The proposed mechanisms of music intervention on
38 depression and anxiety span several domains. Firstly, music intervention can exert
39 neurophysiological and neurochemical effects, such as triggering endorphin release[29] and
40 stimulating cognitive functioning.[30] Secondly, music intervention can promote positive self-
41 emotions including enhancing the sense of happiness[31] and activating self-efficacy and self-
42 confidence through experiences of success.[32] Lastly, music intervention can build up sociability,
43 such as interaction skills[33] and interpersonal relationship.[34] Furthermore, there are published
44 studies indicating that music intervention combined with diabetic routine treatments could improve
45 depression score, anxiety score, fasting glucose and glycosylated hemoglobin (HbA1c).[35,36]

Why it is important to do this review

From the above, many studies have indicated the bidirectional relationship between diabetes and mental disorders. Music intervention, as a therapeutic method, has been applied to alleviate depression and anxiety and improve related indicators of diabetes. By searching literatures, it can be found that some reviews evaluated the effect of music intervention on patients with mental disorders,[37-39] while no related systematic review and meta-analysis can be searched about the effect of music intervention on mental health of patients with diabetes. Therefore, if music intervention can be proved to be both valid and with few side effects through this study, it is worth being promoted and applicated in clinic all over the world to benefit more people since this method is easy to be operated.

Objectives

The purpose of this review is to systematically assess the effect of music intervention and compare the effects between music therapy and music medicine on depression and anxiety of diabetic patients. Additionally, it is also our expectation to furnish evidence reference for related guideline.

METHODS

Study registration

Our protocol for the meta-analysis has been registered on the International Prospective Register of Systematic Reviews (PROSPERO), and CRD42019146439 is the registration ID. This protocol has been planned severely on the basis of Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P)[40] statement guidelines. The results will be reported in conformity to the PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Healthcare Interventions.[41]

Eligibility criteria

1
2
3
4 Studies that have compliance with the following inclusion criteria will be rigorously screened.

5
6 Type of studies

7 We will merely cover randomized controlled trials (RCTs) in English or Chinese, excepting Quasi-
8 RCTs and cluster RCTs. Experiments on animals and nonrandomized clinical trials will be out of
9 our consideration. Articles with significant overlap with other articles through print or e-publishing
10 will fail to be included. Duplicate publications, that is results originated from a single experiment
11 but measured by other standards and then published as disparate papers, will also be excluded.
12 Publications with originality is our first choice. The publication status will not be restricted.
13
14
15
16
17
18
19
20

21 Type of Participants

22 Diabetic Patients with depression or anxiety symptoms, regardless of the type of diabetes and no
23 restriction on gender, age, race, economic ability and education level. Patients of other disease or
24 healthy persons will be excluded.
25
26
27
28
29
30

31 Type of interventions and controls

32 Interventions include music intervention (music therapy or music medicine) alone or music
33 intervention in combination with other interventions. No restriction on the type of music. Other
34 interventions refer to: (1) routine treatments for diabetes which mainly include pharmacologic
35 approaches to glycemic treatment, diabetes self-management education and support (DSMES), etc.
36 according to the newest guideline of American Diabetes Association (ADA);[42,43] (2) other
37 therapies such as acupoint therapy,[44] metabolic surgery,[45] etc. When intervention is music
38 intervention alone, controls can be no treatment, placebo or other active therapies. Notably, when
39 intervention is music intervention combined with active therapies, the control must be the same
40 active therapies.
41
42
43
44
45
46
47
48
49
50

51 Type of outcome measures

52 This review aims at systematically assessing the effect of music intervention on depression or
53
54
55
56

1
2
3
4 anxiety of diabetic patients, so we choose depression score and anxiety score as the primary
5
6 outcomes, which can give an index to the degree of depression and anxiety by scales.

7
8 Primary outcomes: Depression score and anxiety score. Depression score is measured using self-
9
10 rating depression scale (SDS), Hamilton depression scale (HAMD) or other validated scales for
11
12 depression. Anxiety score is measured using self-rating anxiety scale (SAS), Hamilton anxiety
13
14 scale (HAMA) or other validated scales for anxiety.

15
16 Secondary outcomes: Safety. Measurement via incidence of adverse effects.
17
18

19 **Search methods**

20 Search resources

21
22 We will search the following nine online electronic databases: PubMed, Web of Science, Embase,
23
24 EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang
25
26 Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical
27
28 Database (CBM), from their inception to March 2020. We also plan to search other relevant
29
30 resources as far as possible, including: (1) grey literatures (like conference literatures and
31
32 dissertations) from the following four sources: OpenGrey (www.opengrey.eu/), CNKI
33
34 (www.cnki.net), Open Access Theses and Dissertations (oatd.org) and British Library EThOS
35
36 (ethos.bl.uk/); (2) the reference lists of relevant publications for additional studies.
37
38
39

40 Search strategies

41
42 The search strategy will include the MeSH terms for diabetes mellitus, depression, anxiety, music
43
44 therapy, music and singing, combined with their respective free-text terms. The search details of
45
46 the PubMed are simply shown in table 1.
47
48
49
50
51
52
53
54
55
56
57

Table 1 Example of PubMed search strategy.

Number	Search terms
#1	"Diabetes Mellitus"[Mesh]
#2	"Depression"[Mesh] OR Depressions[Title/Abstract] OR Depressive Symptoms[Title/Abstract] OR Depressive Symptom[Title/Abstract] OR Symptom, Depressive[Title/Abstract] OR Symptoms, Depressive[Title/Abstract] OR Emotional Depression[Title/Abstract] OR Depression, Emotional[Title/Abstract] OR Depressions, Emotional[Title/Abstract] OR Emotional Depressions[Title/Abstract]
#3	"Anxiety"[Mesh] OR Hypervigilance [Title/Abstract] OR Nervousness [Title/Abstract]
#4	#2 OR #3
#5	"Music Therapy"[Mesh] OR Therapy, Music [Title/Abstract]
#6	"Music"[Mesh] OR music intervention[Title/Abstract] OR music medicine[Title/Abstract] OR music listening[Title/Abstract] OR audio recording[Title/Abstract] OR mp3 hearing[Title/Abstract]
#7	"Singing"[Mesh] OR sing[Title/Abstract] OR sings[Title/Abstract] OR song[Title/Abstract] OR compose[Title/Abstract] OR composing[Title/Abstract]
#8	#5 OR #6 OR #7
#9	randomized controlled trial [Publication Type] OR randomized [Title/Abstract] OR placebo [Title/Abstract]
#10	#1 AND #4 AND #8 AND #9

Data collection and analysis

Studies selection

Firstly, two researchers (L-yZ and YT) will respectively screen the keywords, titles and abstracts of all obtained studies using endnote x9 software and determine whether trials have compliance with inclusion criteria. Secondly, we will also get access to the full text of all studies of possible relevance for further assessment. Each of excluded study will be noted with reasons. Any disagreements will be settled through consensus or referral to the third researcher (XxF). The

1
2
3
4 specific selection steps are presented in the figure 1.
5
6

7 Data extraction

8
9 Two researchers (L-yZ and YT) will independently extract and tabulate the following data items
10 via Microsoft Excel:
11

- 12 1. General information (including publication year and name of the first author).
- 13 2. Study design (including random method, allocation concealment and blinding).
- 14 3. Participants (including sample size, gender, age, area, depression score, anxiety score and course
15 of disease).
- 16 4. Treatment (including interventions, controls, music type and the course of treatment).
- 17 5. Outcomes (including depression score, anxiety score and safety).

18
19 If there is any different opinion, the final decision will go to the third researcher (XxF) to be
20 made.
21
22

23 Assessment of risk of bias

24
25 The Cochrane Collaboration's Risk of Bias tool[46] will be applied to evaluate the risk of bias of
26 all included RCT studies by two authors (L-zW and YZ). The following domains will be assessed:
27 sequence generation, allocation concealment, blinding of participants and personnel, blinding of
28 outcome assessors, incomplete outcome data, selective reporting and other sources of bias. The
29 assessments will be classified into "low", "unclear" or "high" of risk.
30
31

32 Measures of treatment effect

33
34 Considering that all outcomes (depression score, anxiety score and safety) are continuous variables,
35 the mean difference (MD) and 95% confidence interval (CI) will be calculated when the
36 quantitative data are from the same measurement method or small data scale. On the contrary, the
37 standardized mean difference (SMD) and 95% CI will be calculated.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Unit of analysis issues

For the standard research which has one experimental group with one control group, the unit of analysis is based on the individual. For the multi-arm study which has more than two intervention groups, the recommended method is to create a single pair-wise comparison through combining groups.

Dealing with missing data

If the required data are unavailable from literatures, one researcher (YZ) will send an e-mail to the corresponding author and ask for further information.

Assessment of heterogeneity

A meta-analysis will be implemented if the included studies are adequately homogenous. *P*-value and *I*² will be used to assess the heterogeneity of statistic. When *P*-value >0.05 and *I*² ≤50%, the heterogeneity is regarded as acceptable. When *P*-value <0.05 or *I*² >50%, the heterogeneity is considered as significant.

Assessment of reporting bias

Publication bias will be implemented, if there are more than 10 studies included, through a funnel plot.

Data synthesis

RevMan 5.3. software provided by the Cochrane collaboration will be used to analyze the data. If the included studies are adequately homogenous, fixed-effects model will be used for statistical combination. Since all outcomes are continuous variables, the mean difference (MD) and 95% confidence interval (CI) will be calculated when the outcomes are from the same measurement method or small data scale. On the contrary, the standardized mean difference (SMD) and 95% CI will be calculated. If there is significant heterogeneity among the included studies, subgroup

1
2
3 analyses or sensitivity analyses will be performed. If a meta-analysis is unavailable, descriptive
4 summaries will be provided.
5
6
7

8 9 Subgroup analysis

10 The subgroup analyses are planned as follows:

- 11 1. Type of music intervention (music therapy or music medicine).
 - 12 2. Type of scales (SDS, HAMD, SAS, HAMA or other validated scales for depression or anxiety).
 - 13 3. Different age groups.
- 14
15
16
17
18
19
20

21 Sensitivity analysis

22 Sensitivity analyses will be performed through including or excluding a specific study, such as a
23 study without a clear allocation concealment or randomization method, to observe whether the
24 results changed.
25
26
27
28
29
30

31 Evidence assessed

32 GRADE guideline will be used to evaluate the quality of evidence.[47] In the light of the GRADE
33 method, the quality of evidence can be assessed by 4 levels: “high”, “medium”, “low” and “very
34 low”. GRADE profiler 3.2 will be applied for analysis.
35
36
37
38
39
40

41 Patient and public involvement

42 Since this study is a systematic review, patients or the public will fail to participate.
43
44
45

46 **DISCUSSION**

47 Depression and anxiety often occur independently or simultaneously during the development of
48 many chronic diseases. In recent years, the number of studies about comorbid depression or anxiety
49 in diabetics are continually increasing, not only because the two-way relationship between
50 depression/anxiety and diabetes can lead to deterioration of diseases, increased difficulty in
51
52
53
54
55
56

1
2
3
4 treatment, even as well as the great mortality,[48-51] but also because therapeutic methods for
5 treating the comorbidities are limited and unsatisfactory. On the one hand, some antidepressant
6 drugs, such as nortriptyline and sertraline, may complicate glycemic control.[52] On the other hand,
7 psychotherapy based on verbal communication requires a professional psychiatrist and it is
8 expensive.
9

10
11
12
13 Music intervention has been proved by many literatures to have a beneficial effect on depression
14 and anxiety, such as strengthening awareness and sensitiveness for positive emotions.[53]
15 Noticeably, a few studies also show that music intervention can reduce blood glucose and HbA1c
16 simultaneously.[24,54] Therefore, music intervention is an applicable method for treating
17 depression and anxiety with diabetes. Additionally, music intervention, almost no side effects, is
18 safer than pharmacotherapy, easier to be operated and costs less than psychotherapy based on verbal
19 communication, so it is conducive to clinical promotion and application.
20
21
22
23
24
25
26

27 Since 2007,[55] music intervention has been applied in the treatment of depression and anxiety
28 with diabetes for 12 years. Nonetheless, there are not systematic review or meta-analysis conducted
29 to evaluate the potential pros and cons of this therapy method. Thus, this study intends to provide
30 strong evidence for music intervention applied to treat mental disorders in diabetic patients and
31 synchronously furnish evidence reference for related guideline.
32
33
34
35
36

37 However, a language bias may exist in this meta-analysis because it will include studies only in
38 English and Chinese because of the language barrier. Therefore, some related studies from
39 medical databases in other languages might be missed.
40
41
42

43 In conclusion, this meta-analysis will be conducive to assess underlying benefits and harms of
44 music intervention on comorbid depression or anxiety with diabetes. Moreover, the results of this
45 study might not merely furnish evidence reference for guideline of relevant areas, but also may
46 attract more attention on music intervention, which can be a chance to facilitate its clinical
47 application and ultimately benefit more patients.
48
49
50
51
52
53

54 **ETHICS AND DISSEMINATION**

55
56
57
58
59
60

1
2
3
4 There is no required ethical approval because this meta-analysis is on the basis of published data.
5
6 The results of this systematic review will be published to a peer reviewed journal.
7
8

9 **Figure legend**

10
11 **Figure 1** Flowchart to present the search process.
12
13

14
15 **Contributions** L-yZ and C-gX contributed to the conception of the study. The manuscript of the
16 protocol was drafted by L-yZ, and was revised by XxF and C-gX. L-yZ and YT will also
17 independently screen the potential studies and extract data from included studies. L-zW and YZ
18 will assess the risk of bias and accomplish data synthesis. XxF will arbitrate any disagreements.
19 C-gX will ensure that no errors occur during the review. All authors have approved the
20 publication of the protocol.
21
22

23
24 **Funding** This study was supported by grants from the Science and Technology Project of
25 Sichuan Province (2019YFS002), the Special Scientific and Technological Research Project of
26 Administration of TCM of Sichuan Province (2018KF003), and the Foundation of Development
27 of Science and Technology of Chengdu University of TCM (2016-D-YY-61).
28
29

30
31 **Competing interests** None declared.
32
33

34
35 **Patient consent for publication** Not required.
36
37

38
39 **Provenance and peer review** Not commissioned; externally peer reviewed.
40
41

42
43 **Open access** This is an open access article distributed in accordance with the Creative Commons
44 Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix,
45 adapt, build upon this work non-commercially, and license their derivative works on different terms,
46 provided the original work is properly cited, appropriate credit is given, any changes made
47 indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.
48
49

50 **REFERENCES**

51
52
53
54 1. International Diabetes Federation. IDF Diabetes Atlas (9th edition), 2019. Available:
55
56

- 1
2
3
4 <http://www.diabetesatlas.org/>. Accessed November 9, 2019
- 5
6 2. World Health Organization. Global report on diabetes, 2014. Available:
7 <https://www.who.int/news-room/fact-sheets/detail/diabetes>. Accessed November 9, 2019
- 8
9
10 3. Ali S, Stone MA, Peters JL, et al. The prevalence of co-morbid depression in adults with
11 Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* 2006;23(11):1165–
12 1173. doi:10.1111/j.1464-5491.2006.01943.x
- 13
14
15 4. Anderson RJ, Freedland KE, Clouse RE, et al. The prevalence of comorbid depression in
16 adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24(6):1069–1078.
17
18 doi:10.2337/diacare.24.6.1069
- 19
20
21 5. Knol MJ, Twisk JW, Beekman AT, et al. Depression as a risk factor for the onset of type 2
22 diabetes mellitus. A meta-analysis. *Diabetologia* 2006;49(5):837–845.doi:10.1007/s00125-
23 006-0159-x
- 24
25
26
27 6. Cosgrove MP, Sargeant LA, Griffin SJ. Does depression increase the risk of developing type
28 2 diabetes?. *Occup Med (Lond)* 2008;58(1):7–14.doi:10.1093/occmed/kqm105
- 29
30
31 7. Mezuk B, Eaton WW, Albrecht S, et al. Depression and type 2 diabetes over the lifespan: a
32 meta-analysis. *Diabetes Care* 2008;31(12):2383–2390.doi:10.2337/dc08-0985
- 33
34
35 8. Grigsby AB, Anderson RJ, Freedland KE, et al. Prevalence of anxiety in adults with diabetes:
36 a systematic review. *J Psychosom Res* 2002; 53: 1053–1060.doi:10.1016/s0022-
37 3999(02)00417-8
- 38
39
40 9. Smith KJ, Beland M, Clyde M, et al. Association of diabetes with anxiety: a systematic
41 review and metaanalysis. *J Psychosom Res* 2013; 74: 89–99.
42
43 doi:10.1016/j.jpsychores.2012.11.013
- 44
45
46 10. Smith KJ, Deschênes SS, Schmitz N. Investigating the longitudinal association between
47 diabetes and anxiety: a systematic review and meta-analysis. *Diabet Med* 2018;35(6):677–
48 693.doi:10.1111/dme.13606
- 49
50
51
52 11. Egede LE. Major depression in individuals with chronic medical disorders: prevalence,
53 correlates and association with health resource utilization, lost productivity and functional
54
55
56

- 1
2
3
4 disability. *Gen Hosp Psychiatry* 2007;29:409–416.doi:10.1016/j.genhosppsy.2007.06.002
- 5
6 12. Naicker K, Johnson JA, Skogen JC, et al. Type 2 Diabetes and Comorbid Symptoms of
7
8 Depression and Anxiety: Longitudinal Associations With Mortality Risk. *Diabetes Care*
9
10 2017;40(3):352–358.doi:10.2337/dc16-2018
- 11
12 13. van der Feltz-Cornelis CM, Nuyen J, Stoop C, et al. Effect of interventions for major
13
14 depressive disorder and significant depressive symptoms in patients with diabetes mellitus:
15
16 a systematic review and meta-analysis. *Gen Hosp Psychiatry* 2010;32(4):380–395.
17
18 doi:10.1016/j.genhosppsy.2010.03.011
- 19
20 14. Komsuoglu Celikyurt I, Mutlu O, Ulak G, et al. Exenatide treatment exerts anxiolytic- and
21
22 antidepressant-like effects and reverses neuropathy in a mouse model of type-2 diabetes. *Med*
23
24 *Sci Monit Basic Res* 2014;20:112–117.doi:10.12659/MSMBR.891168
- 25
26 15. Leubner D, Hinterberger T. Reviewing the Effectiveness of Music Interventions in Treating
27
28 Depression. *Front Psychol* 2017;8:1109.doi:10.3389/fpsyg.2017.01109
- 29
30 16. American Music Therapy Association. What is music therapy?, 2013.Available:
31
32 <https://www.musictherapy.org/about/musictherapy/>. Accessed January 9, 2020.
- 33
34 17. Dileo C. A classification model for music and medicine. National Association of Music
35
36 Therapy, Washington, DC, 1999: 1-6.
- 37
38 18. Raglio A, Bellelli G, Mazzola P, et al. Music, music therapy and dementia: a review of
39
40 literature and the recommendations of the Italian Psychogeriatric Association. *Maturitas*
41
42 2012;72(4):305–310.doi:10.1016/j.maturitas.2012.05.016
- 43
44 19. Dileo C. Effects of music and music therapy on medical patients: a meta-analysis of the
45
46 research and implications for the future. *J Soc Integr Oncol* 2006;4(2):67–70.
47
48 doi:10.2310/7200.2006.002
- 49
50 20. Thaut MH, Gardiner JC, Holmberg D, et al. Neurologic music therapy improves executive
51
52 function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*
53
54 2009;1169:406–416.doi:10.1111/j.1749-6632.2009.04585.x
- 55
56 21. Kim DS, Park YG, Choi JH, et al. Effects of music therapy on mood in stroke patients.

- 1
2
3
4 *Yonsei Med J* 2011;52(6):977–981.doi:10.3349/ymj.2011.52.6.977
- 5
6 22. Nwebube C, Glover V, Stewart L. Prenatal listening to songs composed for pregnancy and
7 symptoms of anxiety and depression: a pilot study. *BMC Complement Altern Med*
8 2017;17(1):256.doi:10.1186/s12906-017-1759-3
- 9
10
11 23. Lin XX, Ji JM, Zhou GY. Influence of listening to Jiao tune music on sleep quality and blood
12 glucose in the elderly patients with type 2 diabetes accompanied by insomnia. *Journal of*
13 *rehabilitation* 2017;27(01):44-48.
- 14
15
16 24. Mandel SE, Davis BA, Secic M. Effects of music therapy and music-assisted relaxation and
17 imagery on health-related outcomes in diabetes education: a feasibility study. *Diabetes Educ*
18 2013;39(4):568–581.doi:10.1177/0145721713492216
- 19
20
21 25. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2014.
22 Available: [https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf)
23 [burden-in-the-united-states.pdf](https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf). Accessed November 9, 2019
- 24
25
26 26. Gabrielsson A, Wik SL. Strong experiences related to music: a descriptive system. *Music Sci*
27 2003;7(2):157–217.
- 28
29
30 27. Zentner M, Grandjean D, Scherer KR. Emotions evoked by the sound of music:
31 characterization, classification, and measurement. *Emotion* 2008;8(4):494–521.
32 doi:10.1037/1528-3542.8.4.494
- 33
34
35 28. Saarikallio S. Music as emotional self-regulation throughout adulthood. *Psychol Music*
36 2011;39(3):307–327.
- 37
38
39 29. Gök Ugur H, Yaman Aktaş Y, Orak OS, et al. The effect of music therapy on depression and
40 physiological parameters in elderly people living in a Turkish nursing home: a randomized-
41 controlled trial. *Aging Ment Health* 2017;21(12):1280–1286.
42 doi:10.1080/13607863.2016.1222348
- 43
44
45 30. Gómez Gallego M, Gómez García J. Music therapy and Alzheimer's disease: Cognitive,
46 psychological, and behavioural effects. *Neurologia* 2017;32(5):300–308.
47 doi:10.1016/j.nrl.2015.12.003
- 48
49
50
51
52
53
54
55
56
57
58
59
60


- 1
2
3
4 31. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement
5 intervention on the cognitive functions of people with moderate dementia: a randomized
6 controlled trial. *Aging Ment Health* 2018;22(3):306–315.
7
8 doi:10.1080/13607863.2016.1251571
9
10
11 32. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in
12 elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging*
13 *Ment Health* 2017;21(2):156–162.doi:10.1080/13607863.2015.1093602
14
15
16
17 33. Bieleninik L, Geretsegger M, Mössler K, et al. Effects of Improvisational Music Therapy vs
18 Enhanced Standard Care on Symptom Severity Among Children With Autism Spectrum
19 Disorder: The TIME-A Randomized Clinical Trial. *JAMA* 2017;318(6):525–535.
20
21
22
23
24
25
26
27
28
29
30
31 34. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a
32 controlled randomized study. *Neurol Sci* 2017;38(8):1485–1493.doi:10.1007/s10072-017-
33 3003-9
34
35
36
37 35. Deng HX, Gao GS, Li ZM, et al. Application of receiving music therapy in elderly patients
38 with diabetes. *Chinese journal of health and nutrition (1st Ten-days of month)*
39
40
41
42
43
44
45
46
47
48
49
50
51 36. Zhang ZW. Application of music therapy and psychotherapy adjuvant therapy in type 2
52 diabetes. *Chinese medical science* 2011,01(18):150-151.
53
54
55
56
57 37. Solanki MS, Zafar M, Rastogi R. Music as a therapy: role in psychiatry. *Asian J Psychiatr*
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
31. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement intervention on the cognitive functions of people with moderate dementia: a randomized controlled trial. *Aging Ment Health* 2018;22(3):306–315. doi:10.1080/13607863.2016.1251571
32. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging Ment Health* 2017;21(2):156–162.doi:10.1080/13607863.2015.1093602
33. Bieleninik L, Geretsegger M, Mössler K, et al. Effects of Improvisational Music Therapy vs Enhanced Standard Care on Symptom Severity Among Children With Autism Spectrum Disorder: The TIME-A Randomized Clinical Trial. *JAMA* 2017;318(6):525–535. doi:10.1001/jama.2017.9478
34. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a controlled randomized study. *Neurol Sci* 2017;38(8):1485–1493.doi:10.1007/s10072-017-3003-9
35. Deng HX, Gao GS, Li ZM, et al. Application of receiving music therapy in elderly patients with diabetes. *Chinese journal of health and nutrition (1st Ten-days of month)* 2013,23(3):1186-1186.
36. Zhang ZW. Application of music therapy and psychotherapy adjuvant therapy in type 2 diabetes. *Chinese medical science* 2011,01(18):150-151.
37. Solanki MS, Zafar M, Rastogi R. Music as a therapy: role in psychiatry. *Asian J Psychiatr* 2013;6(3):193–199.doi:10.1016/j.ajp.2012.12.001
38. Petrovsky D, Cacchione PZ, George M. Review of the effect of music interventions on symptoms of anxiety and depression in older adults with mild dementia. *Int Psychogeriatr* 2015;27(10):1661–1670.doi:10.1017/S1041610215000393
39. Aalbers S, Fusar-Poli L, Freeman RE, et al. Music therapy for depression. *Cochrane Database Syst Rev* 2017;11(11):CD004517. Published 2017 Nov 16. doi:10.1002/14651858.CD004517.pub3

- 1
2
3
4 40. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and
5 meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*
6 2015;350:g7647.doi:10.1136/bmj.g7647
7
8
9
10 41. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic
11 reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and
12 elaboration. *BMJ* 2009;339:b2700.doi:10.1136/bmj.b2700
13
14
15 42. American Diabetes Association. 9. Pharmacologic Approaches to Glycemic Treatment:
16 Standards of Medical Care in Diabetes-2020. *Diabetes Care* 2020;43(Suppl 1):S98–S110.
17 doi:10.2337/dc20-S009
18
19
20
21 43. American Diabetes Association. 5. Facilitating Behavior Change and Well-being to Improve
22 Health Outcomes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*
23 2020;43(Suppl 1):S48–S65. doi:10.2337/dc20-S005
24
25
26
27 44. Feng Y, Fang Y, Wang Y, et al. Acupoint Therapy on Diabetes Mellitus and Its Common
28 Chronic Complications: A Review of Its Mechanisms. *Biomed Res Int* 2018;2018:3128378.
29 doi:10.1155/2018/3128378
30
31
32
33 45. Pérez-Pevida B, Escalada J, Miras AD, et al. Mechanisms Underlying Type 2 Diabetes
34 Remission After Metabolic Surgery. *Front Endocrinol (Lausanne)* 2019;10:641.
35 doi:10.3389/fendo.2019.00641
36
37
38
39 46. Higgins JPT, Thomas J, Chandler J, et al. Cochrane Handbook for Systematic Reviews of
40 Interventions version 6.0 (updated July 2019). Cochrane, 2019. Available:
41 www.training.cochrane.org/handbook.
42
43
44
45 47. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of
46 evidence. *J Clin Epidemiol* 2011;64:401–406.doi:10.1016/j.jclinepi.2010.07.015
47
48
49 48. Gonzalez JS, Peyrot M, McCarl LA, et al. Depression and diabetes treatment nonadherence: a
50 meta-analysis. *Diabetes Care* 2008;31(12):2398–2403.doi:10.2337/dc08-1341
51
52
53 49. Trief PM, Foster NC, Chaytor N, et al. Longitudinal Changes in Depression Symptoms and
54 Glycemia in Adults With Type 1 Diabetes. *Diabetes Care* 2019;42(7):1194–1201.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

doi:10.2337/dc18-2441

50. Novak M, Mucsi I, Rhee CM, et al. Increased Risk of Incident Chronic Kidney Disease, Cardiovascular Disease, and Mortality in Patients With Diabetes With Comorbid Depression. *Diabetes Care* 2016;39(11):1940–1947. doi:10.2337/dc16-0048
51. Lustman PJ, Anderson RJ, Freedland KE, et al. Depression and poor glycemic control: a meta-analytic review of the literature. *Diabetes Care* 2000;23(7):934–942. doi:10.2337/diacare.23.7.934
52. Roopan S, Larsen ER. Use of antidepressants in patients with depression and comorbid diabetes mellitus: a systematic review. *Acta Neuropsychiatr* 2017;29(3):127–139. doi:10.1017/neu.2016.54
53. Croom AM. Music, neuroscience, and the psychology of well-being: a précis. *Front Psychol* 2012;2:393. doi:10.3389/fpsyg.2011.00393
54. Abd-Elshafy SK, Khalaf GS, Abo-Kerisha MZ, et al. Not All Sounds Have Negative Effects on Children Undergoing Cardiac Surgery. *J Cardiothorac Vasc Anesth* 2015;29(5):1277–1284. doi:10.1053/j.jvca.2015.01.005
55. Yu SJ, Sun ZY, Zhou J, et al. Exploration of music behavioral intervention in the treatment of diabetes mellitus. *Hebei med* 2007;13(8):903-906.



PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification P₁	1a	Identify the report as a protocol of a systematic review
Update None	1b	If the protocol is for an update of a previous systematic review, identify as such
Registration P₂	2	If registered, provide the name of the registry (such as PROSPERO) and registration number
Authors:		
Contact P₁	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author
Contributions P₁₄	3b	Describe contributions of protocol authors and identify the guarantor of the review
Amendments None	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments
Support:		
Sources P₁₄	5a	Indicate sources of financial or other support for the review
Sponsor None	5b	Provide name for the review funder and/or sponsor
Role of sponsor or funder None	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol
INTRODUCTION		
Rationale P₄₋₆	6	Describe the rationale for the review in the context of what is already known
Objectives P₇₋₈	7	Provide an explicit statement of the question(s) the review will address with reference to participants P₇ , interventions P₇ , comparators P₇ , and outcomes P₇₋₈ (PICO)
METHODS		
Eligibility criteria P₆₋₇	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review
Information sources P₆, P₈, P₁₁	9	Describe all intended information sources (such as electronic databases P₈ , contact with study authors P₁₁ , trial registers P₆ or other grey literature sources P₈) with planned dates of coverage P₈
Search strategy P₉	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated
Study records:		

Data management P₁₀	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review
Selection process P₉₋₁₀	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)
Data collection process P₁₀	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators
Data items P₁₀	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications
Outcomes and prioritization P₇₋₈	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale
Risk of bias in individual studies P₁₀	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis
Data synthesis P₁₀₋₁₂	15a	Describe criteria under which study data will be quantitatively synthesised P₁₁
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I ² , Kendall's τ) P₁₀₋₁₁
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) P₁₂
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned P₁₂
Meta-bias(es) P₁₁	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)
Confidence in cumulative evidence P₁₂	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.

BMJ Open

Effect of music intervention on mental health in patients with diabetes mellitus: protocol for a systematic review and meta-analysis of randomized controlled trials

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-036268.R3
Article Type:	Protocol
Date Submitted by the Author:	07-Jul-2020
Complete List of Authors:	Zhou, Lin-yue ; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Zhang, Yuan ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Tian, Yuan; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Fu, Xiaoxu; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Wang, Li-zhen ; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine Xie, Chun-guang; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Endocrinology; Chengdu University of Traditional Chinese Medicine, School of Clinical Medicine
Primary Subject Heading:	Diabetes and endocrinology
Secondary Subject Heading:	Mental health, Complementary medicine
Keywords:	DIABETES & ENDOCRINOLOGY, MENTAL HEALTH, COMPLEMENTARY MEDICINE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Effect of music intervention on mental health in patients with diabetes mellitus: protocol for a systematic review and meta-analysis of randomized controlled trials

Lin-yue Zhou,^{1,2} Yuan Zhang,² Yuan Tian,² Xiaoxu Fu,^{1,2} Li-zhen Wang,² Chun-guang Xie*^{1,2}

Author affiliations

¹Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Chengdu, Sichuan Province, China

²School of Clinical Medicine, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan Province, China

*Correspondence to:

Corresponding Author: Professor Chun-guang Xie

Address: Department of Endocrinology, Chengdu University of Traditional Chinese Medicine Affiliated Hospital, No 39 Shi-er-qiao Road, Chengdu, Sichuan Province, China, 610075

Tel: (86)13438826824

E-mail: killuya233@stu.cdutcm.edu.cn

Word count:2725 words (excluding title page, abstract, references, figures and tables)

ABSTRACT

Introduction About 463 million adults aged 20 to 79 have diabetes globally. Mental disorders often exist in diabetic patients as comorbidities, which can lead to aggravation of the diseases, increased difficulties in treatment, as well as elevated mortality rates. Music intervention has been applied in the treatment of comorbidities for 12 years now, but there are still no recommendations due to the lack of evidence. Thus, a meta-analysis is necessary to evaluate the effect of music intervention in treating mental disorders of diabetic patients.

Methods and analysis We will search the following nine online electronic databases from their inception until March 2020: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP) and Chinese Biomedical and Medical Database (CBM). We also plan to search other relevant resources, including grey literature and the reference lists of relevant publications. Only randomized controlled trials (RCTs) of music intervention to treat depression or anxiety in diabetic patients will be involved. The primary outcomes include the depression score and anxiety score measured on certain scales, and the secondary outcome is safety. Data extraction will be independently implemented by two researchers. The risk of bias will be evaluated through the Cochrane Collaboration's Risk of Bias tool. Eventually, all the data will be analyzed via the Review Manager 5.3 software.

Ethics and dissemination This meta-analysis will provide information about applying music intervention to treat depression or anxiety in diabetic patients. No ethical approval is required because this meta-analysis is based on published data. The results of this systematic review will be published in a peer-reviewed journal.

PROSPERO registration number CRD42019146439

Strengths and limitations of this study

· This meta-analysis will be the first to provide evidence on the effectiveness and safety of applying music intervention to treat depression or anxiety in diabetic patients.

· Based on our comprehensive search strategy, the relevant original studies will be included.

· Two authors will separately implement data extraction and estimation of the risk of bias.

· We will only cover randomized controlled trials (RCTs) in this study.

· A language bias may exist in this meta-analysis because it will include only the studies in English and Chinese due to the language barrier.

INTRODUCTION

According to the newest “IDF Diabetes Atlas (9th edition)” published by the International Diabetes Federation (IDF), about 463 million adults aged 20 to 79 have diabetes mellitus globally (1 in 11).[1] Diabetes has a well-known significant influence on the patient’s quality of life (QoL) due to its complications, such as retinopathy, cardiovascular, neurological or renal diseases, as well as foot ulcers, which may result in the amputation of lower extremity.[2]

In addition to the physical complications, diabetic patients may also develop mental disorders as comorbidities. Some studies have indicated that depression is more prevalent in diabetic patients compared with non-diabetics.[3,4] Meanwhile, depression could also increase the risk of diabetes,[5-7] which suggests a bidirectional diabetes-depression relation; a similar relation has been confirmed between anxiety and diabetes: people with diabetes are more susceptible to concurrent anxiety[8,9] while anxiety is a risk factor of diabetes.[10] Individuals with chronic diseases who have comorbid depression or anxiety have obviously higher healthcare utilization, functional disability, work absence, and mortality risk than healthy ones.[11,12]

When it comes to the treatment of diabetes with comorbid depression or anxiety, conventional glucose-lowering treatments combined with antidepressants, anti-anxiety medications, or psychological intervention are the routine treatments.[13,14] Music intervention is also a successful therapy, which can be divided into two major categories: music therapy and music medicine.[15] The official definition of music therapy is the “clinical and evidence-based use of music interventions to accomplish individualized goals within a therapeutic relationship by a credentialed professional who has completed an approved music therapy program”.[16] Based on the patient’s condition, a trained music therapist conducts the treatment by choosing the type of music, the form of music presentation (pre-recorded or live), and the treatment setting (singing, playing instruments, composing, etc.), ultimately resulting in a therapeutic scheme that is best suited for the patient. On the other hand, music medicine is defined as passive listening to pre-recorded music, which can be independently implemented by medical or healthcare professionals instead of music therapists.[17] In brief, music therapy focuses on active music therapy and

1
2
3
4 personalized treatments, while music medicine belongs to passive music listening and non-
5
6 personalized therapy. It is important to emphasize the distinction between music therapy and
7
8 music medicine because some studies suggest that the effect of the former is better than the latter
9
10 with regard to many outcomes.[18-19]

11 A lot of studies suggested that music, with all its types, demonstrates a remarkable effect in
12
13 alleviating depression and anxiety among patients with various diseases.[20-22] As for diabetes,
14
15 recent studies have shown that music intervention could improve fasting blood glucose and
16
17 systolic blood pressure in diabetic patients.[23,24] Good control of blood glucose and blood
18
19 pressure can effectively reduce the incidence of serious complications of diabetes.[25]

21 Previous research on music psychology showed the special potential of music. As a stimulus,
22
23 music can not only trigger a variety of emotional experiences,[26, 27] but it can also strengthen
24
25 the ability to deal with negative emotions in everyday life.[28] The proposed mechanisms behind
26
27 the effects of music intervention on depression and anxiety span across several domains. Firstly,
28
29 music intervention can exert neurophysiological and neurochemical effects, such as triggering the
30
31 release of endorphin[29] and stimulating cognitive functioning.[30] Secondly, music intervention
32
33 can promote positive self-conscious emotions including enhancing the sense of happiness[31] and
34
35 activating self-efficacy and self-confidence through experiencing success.[32] Lastly, music
36
37 intervention can enhance the sociable aspects, such as interaction skills[33] and interpersonal
38
39 relationships.[34] Furthermore, many published studies indicated that music intervention
40
41 combined with the diabetic routine treatments could improve the depression score, anxiety score,
42
43 and the values of fasting glucose and glycosylated hemoglobin (HbA1c).[35,36]

44 By searching the literature, it can be found that some reviews evaluated the effect of music
45
46 intervention on the patients with mental disorders,[37-39] while no related systematic reviews or
47
48 meta-analyses can be found about the effect of music intervention on the mental health of diabetic
49
50 patients. Therefore, a review that can show music intervention to be both valid and with few side
51
52 effects would be useful to indicate that this intervention is worth being promoted and clinically
53
54 applied to benefit more people with this easy-operated method.
55

OBJECTIVE

In this review, we present a meta-analysis to systematically assess the effect of music intervention and compare the effects of music therapy and those of music medicine on depression and anxiety of diabetic patients. Additionally, it serves to establish an evidence reference for related guidelines.

METHODS AND ANALYSIS

Study registration

Our protocol for the meta-analysis has been registered in the International Prospective Register of Systematic Reviews (PROSPERO) with the registration ID of CRD42019146439. This protocol has been strictly planned based on the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P)[40] statement guidelines. The results will be reported in accordance with the PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Healthcare Interventions.[41]

Eligibility criteria

Type of studies

This analysis will merely cover the randomized controlled trials (RCTs) in English or Chinese, except for Quasi-RCTs and cluster RCTs. The experiments on animals and nonrandomized clinical trials will not be considered. The articles with a significant overlap with other articles through printing or e-publishing will not be included. Duplicate publications, in which the results originated from a single experiment but were measured by other standards and then published as disparate papers, will also be excluded. The main choice is original publications, regardless of the publication status.

Type of participants

1
2
3
4 Diabetic patients with depression or anxiety symptoms, regardless of the type of diabetes, gender,
5 age, race, economic ability, and education level. Patients with other diseases or healthy persons
6 will be excluded.
7
8
9

10 11 Type of interventions and controls 12

13 The interventions include music intervention (music therapy or music medicine, no restriction on
14 the type of music) alone or combined with other interventions that refer to the following therapies:
15 (1) routine treatments for diabetes, which mainly include pharmacologic approaches to glycemic
16 treatment, diabetes self-management education and support (DSMES), and others, according to
17 the newest guideline of the American Diabetes Association (ADA);[42,43] (2) other therapies
18 such as acupoint therapy,[44] metabolic surgery,[45] etc. When the used intervention is the music
19 intervention alone, the controls can be people with no treatment, placebo, or other active therapies,
20 while the same therapies should be used with the controls when the music intervention is used
21 combined with active therapies.
22
23
24
25
26
27
28
29
30
31
32

33 Type of outcome measures 34

35 Since this review aims at the systematic assessment of the effect of music intervention on
36 depression or anxiety in diabetic patients, we will choose the depression and anxiety scores as the
37 primary outcomes, which can indicate the degree of depression and anxiety on certain scales.
38
39

40 The depression score is measured using the self-rating depression scale (SDS), Hamilton
41 depression scale (HAMD), or other validated scales for depression. The anxiety score is measured
42 using the self-rating anxiety scale (SAS), Hamilton anxiety scale (HAMA), or other validated
43 scales for anxiety.
44
45
46
47

48 The secondary outcome is the safety, measured via the incidence of adverse effects.
49
50
51

52 **Search methods** 53

54 Search resources 55 56 57 58 59 60

1
2
3
4 The following nine online electronic databases will be searched from their inception to March
5 2020: PubMed, Web of Science, Embase, EBSCO, Cochrane Library, Chinese National
6 Knowledge Infrastructure (CNKI), Wanfang Database, Chinese Scientific Journal Database (VIP)
7 and Chinese Biomedical and Medical Database (CBM). We also plan to search other relevant
8 resources as far as possible, including: (1) grey literature (like conference literature and
9 dissertations) from the following four sources: OpenGrey (www.opengrey.eu/), CNKI
10 (www.cnki.net), Open Access Theses and Dissertations (oatd.org), and the British Library
11 EThOS (ethos.bl.uk/); (2) the reference lists of relevant publications for additional studies.
12
13
14
15
16
17
18
19
20

21 Search strategies

22
23 The search strategy will include the MeSH terms of diabetes mellitus, depression, anxiety, music
24 therapy, music and singing, combined with their respective free-text terms. The search details of
25 PubMed are simplified in table 1.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Example of PubMed search strategy

Number	Search terms
#1	"Diabetes Mellitus"[Mesh]
#2	"Depression"[Mesh] OR Depressions[Title/Abstract] OR Depressive Symptoms[Title/Abstract] OR Depressive Symptom[Title/Abstract] OR Symptom, Depressive[Title/Abstract] OR Symptoms, Depressive[Title/Abstract] OR Emotional Depression[Title/Abstract] OR Depression, Emotional[Title/Abstract] OR Depressions, Emotional[Title/Abstract] OR Emotional Depressions[Title/Abstract]
#3	"Anxiety"[Mesh] OR Hypervigilance [Title/Abstract] OR Nervousness [Title/Abstract]
#4	#2 OR #3
#5	"Music Therapy"[Mesh] OR Therapy, Music [Title/Abstract]
#6	"Music"[Mesh] OR music intervention[Title/Abstract] OR music medicine[Title/Abstract] OR music listening[Title/Abstract] OR audio recording[Title/Abstract] OR mp3 hearing[Title/Abstract]
#7	"Singing"[Mesh] OR sing[Title/Abstract] OR sings[Title/Abstract] OR song[Title/Abstract] OR compose[Title/Abstract] OR composing[Title/Abstract]
#8	#5 OR #6 OR #7
#9	randomized controlled trial [Publication Type] OR randomized [Title/Abstract] OR placebo [Title/Abstract]
#10	#1 AND #4 AND #8AND #9

Data collection and analysis

Studies selection

First, two researchers (L-yZ and YT) will respectively screen the keywords, titles, and abstracts of all the obtained studies using the EndNote X9 software and determine whether the trials comply with the inclusion criteria. Then, we will also get access to the full text of all the studies with possible relevance for further assessment. Each of the excluded studies will be noted with the exclusion reasons. Any disagreements will be settled through consensus or referral to a third researcher (XxF). The specific selection steps are presented in figure 1.

Data extraction

Two researchers (L-yZ and YT) will independently extract and tabulate the following data items using the Microsoft Excel software:

1. General information (including the publication year and name of the first author).
2. Study design (including the random method, allocation concealment, and blinding).
3. Participants (including the sample size, gender, age, area, depression score, anxiety score, and course of disease).
4. Treatment (including the interventions, controls, music type, and the course of treatment).
5. Outcomes (including the depression score, anxiety score, and safety).

In the case of different opinions, the final decision will be made by the third researcher (XxF).

Assessment of risk of bias

In order to evaluate the risk of bias of all the included RCT studies, the Cochrane Collaboration's Risk of Bias tool[46] will be applied by two authors (L-zW and YZ). The assessed domains will be the following: sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessors, incomplete outcome data, selective reporting, and other sources of bias. The assessments will be classified into "low", "unclear" or "high" risk.

Measures of treatment effect

Considering that all the outcomes (depression score, anxiety score, and safety) are continuous variables, when the quantitative data are from the same measurement method or a small data scale, the mean difference (MD) and 95% confidence interval (CI) will be calculated. Otherwise, the standardized mean difference (SMD) and 95% CI will be calculated.

Unit of analysis issues

For the standard research that has one experimental group with one control group, the unit of

1
2
3
4 analysis is the individual. For the multi-arm study with more than two intervention groups, the
5 recommended method is to create a single pair-wise comparison by combining the groups.
6
7

8 9 Dealing with missing data

10
11 If the required data are unavailable from the literature, one researcher (YZ) will contact the
12 corresponding author via email and ask for further information.
13
14
15

16 17 Assessment of heterogeneity

18
19 A meta-analysis will be implemented if the included studies are adequately homogenous. The p
20 value and I^2 are used to assess the heterogeneity of the statistics. The heterogeneity is considered
21 to be acceptable when p value > 0.05 and $I^2 \leq 50\%$ and significant when p value < 0.05 or $I^2 >$
22 50%.
23
24
25
26
27
28

29 30 Assessment of reporting bias

31 The publication bias will be implemented if more than 10 studies are included through a funnel
32 plot.
33
34
35
36

37 38 Data synthesis

39 The Review Manager 5.3 software, provided by the Cochrane collaboration, will be used to
40 analyze the data. If the included studies are adequately homogenous, then a fixed-effects model
41 will be used for the statistical combination. Since all the outcomes are continuous variables in this
42 meta-analysis, we will calculate the mean difference (MD) and 95% confidence interval (CI)
43 when the outcomes are from the same measurement method or a small data scale. Otherwise, we
44 will calculate the standardized mean difference (SMD) and 95% CI. If there is significant
45 heterogeneity among the included studies, subgroup or sensitivity analyses will be performed. If a
46 meta-analysis is unavailable, descriptive summaries will be provided.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Subgroup analysis

The subgroup analyses are planned as follows:

1. The type of music intervention (music therapy or music medicine).
2. The type of scales (SDS, HAMD, SAS, HAMA, or other validated depression or anxiety scales).
3. The different age groups.

Sensitivity analysis

The sensitivity analyses will be performed by including or excluding a specific study, such as a study without clear allocation concealment or a randomization method, to observe whether the results change.

Evaluating the evidence

The GRADE guideline will be used to evaluate the quality of evidence.[47] Based on the GRADE method, the quality of evidence can be assessed by 4 levels: “high”, “medium”, “low” and “very low”. The GRADE profiler 3.2 software will be applied for the analysis.

Patient and public involvement

No patients or volunteers from the public will be involved in this study.

DISCUSSION

Depression and anxiety often occur independently or simultaneously during the development of many chronic diseases. In recent years, an increasing number of studies have been investigating the comorbid depression or anxiety in diabetics. This increased interest is not only because the bidirectional relationship between depression/anxiety and diabetes can lead to deterioration of the diseases, increased difficulties in treatment, as well as elevated mortality rates,[48-51] but also because the therapeutic methods for treating the comorbidities are limited and unsatisfactory. On

1
2
3
4 the one hand, some antidepressant drugs, such as nortriptyline and sertraline, may complicate the
5 glycaemic control.[52] On the other hand, psychotherapy that is based on verbal communication
6 requires a professional psychiatrist and is expensive.
7

8
9 Music intervention has been proved by many studies to have a beneficial effect on depression
10 and anxiety, as it strengthens the awareness and sensitiveness for positive emotions.[53]
11 Noticeably, a few studies also showed that music intervention can simultaneously reduce blood
12 glucose and HbA1c levels.[24,54] Therefore, music intervention is an applicable method to treat
13 depression and anxiety in diabetic patients. Additionally, this almost side-effect-free intervention
14 is safer than pharmacotherapy, easier to be operated and costs less than psychotherapy based on
15 verbal communication; thus, it is conducive to clinical promotion and application.
16
17

18
19 Although music intervention has been applied in the treatment of depression and anxiety with
20 diabetes for 12 years now,[55] there is still no systematic review or meta-analysis to evaluate the
21 potential pros and cons of this therapeutic method. Thus, this study intends to provide strong
22 evidence on the benefits of applying music intervention to treat mental disorders in diabetic
23 patients and synchronously provide evidence to which related guidelines can refer.
24
25

26
27 However, a language bias may exist in this meta-analysis because it includes only the studies in
28 English and Chinese due to the language barrier. Therefore, some related studies from the
29 medical databases in other languages might be missed.
30
31

32
33 In conclusion, this meta-analysis will help to assess the underlying benefits and harms of music
34 intervention on comorbid depression or anxiety in diabetic patients. Moreover, the results of this
35 study will represent a reference for the guidelines of relevant areas, attracting more attention to
36 the music intervention, and facilitating its clinical application to benefit more patients.
37
38

39 **ETHICS AND DISSEMINATION**

40
41 Since this meta-analysis used published data, no ethical approval was required. The results of this
42 systematic review will be published in a peer-reviewed journal.
43
44
45
46
47

Figure legend

Figure 1 Flowchart to present the search process.

Contributions L-yZ and C-gX contributed to the conception of the study. The manuscript of the protocol was drafted by L-yZ and revised by XxF and C-gX. L-yZ and YT will independently screen the potential studies and extract the data from the included studies. L-zW and YZ will assess the risk of bias and accomplish data synthesis. XxF will arbitrate any disagreements. C-gX will ensure that no errors occur during the review. All the authors approved the publication of the protocol.

Funding This study was supported by grants from the Science and Technology Project of Sichuan Province (2019YFS002), the Special Scientific and Technological Research Project of Administration of TCM of Sichuan Province (2018KF003), and the Foundation of Development of Science and Technology of Chengdu University of TCM (2016-D-YY-61).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. International Diabetes Federation. IDF Diabetes Atlas (9th edition), 2019. Available: <http://www.diabetesatlas.org/> [Accessed 9 Nov 2019].
2. World Health Organization. Global report on diabetes, 2014. Available: <https://www.who.int/news-room/fact-sheets/detail/diabetes> [Accessed 9 Nov 2019].

3. Ali S, Stone MA, Peters JL, et al. The prevalence of co-morbid depression in adults with type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* 2006;23:1165–73.doi:10.1111/j.1464-5491.2006.01943.x
4. Anderson RJ, Freedland KE, Clouse RE, et al. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24:1069–78.doi:10.2337/diacare.24.6.1069
5. Knol MJ, Twisk JW, Beekman AT, et al. Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. *Diabetologia* 2006;49:837–45.doi:10.1007/s00125-006-0159-x
6. Cosgrove MP, Sargeant LA, Griffin SJ. Does depression increase the risk of developing type 2 diabetes?. *Occup Med (Lond)* 2008;58:7–14.doi:10.1093/occmed/kqm105
7. Mezuk B, Eaton WW, Albrecht S, et al. Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care* 2008;31:2383–90.doi:10.2337/dc08-0985
8. Grigsby AB, Anderson RJ, Freedland KE, et al. Prevalence of anxiety in adults with diabetes: a systematic review. *J Psychosom Res* 2002;53:1053–60.doi:10.1016/s0022-3999(02)00417-8
9. Smith KJ, Beland M, Clyde M, et al. Association of diabetes with anxiety: a systematic review and meta-analysis. *J Psychosom Res* 2013;74:89–99.doi:10.1016/j.jpsychores.2012.11.013
10. Smith KJ, Deschênes SS, Schmitz N. Investigating the longitudinal association between diabetes and anxiety: a systematic review and meta-analysis. *Diabet Med* 2018;35:677–93.doi:10.1111/dme.13606
11. Egede LE. Major depression in individuals with chronic medical disorders: prevalence, correlates and association with health resource utilization, lost productivity and functional disability. *Gen Hosp Psychiatry* 2007;29:409–16.doi:10.1016/j.genhosppsy.2007.06.002
12. Naicker K, Johnson JA, Skogen JC, et al. Type 2 diabetes and comorbid symptoms of depression and anxiety: longitudinal associations with mortality risk. *Diabetes Care* 2017;40:352–8.doi:10.2337/dc16-2018


- 1
2
3
4 13. van der Feltz-Cornelis CM, Nuyen J, Stoop C, et al. Effect of interventions for major
5 depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a
6 systematic review and meta-analysis. *Gen Hosp Psychiatry* 2010;32:380–
7
8 95.doi:10.1016/j.genhosppsy.2010.03.011
9
- 10
11 14. Komsuoglu Celikyurt I, Mutlu O, Ulak G, et al. Exenatide treatment exerts anxiolytic- and
12 antidepressant-like effects and reverses neuropathy in a mouse model of type-2 diabetes. *Med*
13
14 *Sci Monit Basic Res* 2014;20:112–7.doi:10.12659/MSMBR.891168
15
- 16
17 15. Leubner D, Hinterberger T. Reviewing the effectiveness of music interventions in treating
18 depression. *Front Psychol* 2017;8:1109.doi:10.3389/fpsyg.2017.01109
19
- 20
21 16. American Music Therapy Association. What is music therapy?, 2013.Available:
22
23 <https://www.musictherapy.org/about/musictherapy/> [Accessed 9 Jan 2020].
24
- 25
26 17. Dileo C. A classification model for music and medicine. National Association of Music
27
28 Therapy. Washington, DC:1999:1-6.
- 29
30 18. Raglio A, Bellelli G, Mazzola P, et al. Music, music therapy and dementia: a review of
31
32 literature and the recommendations of the Italian Psychogeriatric Association. *Maturitas*
33
34 2012;72:305–10.doi:10.1016/j.maturitas.2012.05.016
- 35
36 19. Dileo C. Effects of music and music therapy on medical patients: a meta-analysis of the
37
38 research and implications for the future. *J Soc Integr Oncol* 2006;4:67–
39
40 70.doi:10.2310/7200.2006.002
- 41
42 20. Thaut MH, Gardiner JC, Holmberg D, et al. Neurologic music therapy improves executive
43
44 function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*
45
46 2009;1169:406–16.doi:10.1111/j.1749-6632.2009.04585.x
- 47
48 21. Kim DS, Park YG, Choi JH, et al. Effects of music therapy on mood in stroke patients.
49
50 *Yonsei Med J* 2011;52:977–81.doi:10.3349/ymj.2011.52.6.977
- 51
52 22. Nwebube C, Glover V, Stewart L. Prenatal listening to songs composed for pregnancy and
53
54 symptoms of anxiety and depression: a pilot study. *BMC Complement Altern Med*
55
56 2017;17:256.doi:10.1186/s12906-017-1759-3

- 1
2
3
4 23. Lin XX, Ji JM, Zhou GY. Influence of listening to Jiao tune music on sleep quality and blood
5 glucose in elderly patients with type 2 diabetes accompanied by insomnia. *Journal of Fujian*
6 *University of Traditional Chinese Medicine* 2017;27:44-8.doi:CNKI:SUN:FYXB.0.2017-01-
7 011
8
9
10
11 24. Mandel SE, Davis BA, Secic M. Effects of music therapy and music-assisted relaxation and
12 imagery on health-related outcomes in diabetes education: a feasibility study. *Diabetes Educ*
13 2013;39:568–81.doi:10.1177/0145721713492216
14
15
16
17 25. Centers for Disease Control and Prevention. National diabetes statistics report, 2020.
18 Available: <https://www.cdc.gov/diabetes/data/statistics/statistics-report.html> [Accessed 3 Jul
19 2020].
20
21
22
23 26. Gabrielsson A, Wik SL. Strong experiences related to music: a descriptive system. *Music Sci*
24 2003;7:157–217.doi:0.1177/102986490300700201
25
26
27 27. Zentner M, Grandjean D, Scherer KR. Emotions evoked by the sound of music:
28 characterization, classification, and measurement. *Emotion* 2008;8:494–
29 521.doi:10.1037/1528-3542.8.4.494
30
31
32
33 28. Saarikallio S. Music as emotional self-regulation throughout adulthood. *Psychol Music*
34 2011;39:307–27.doi:10.1177/0305735610374894
35
36
37 29. Gök Ugur H, Yaman Aktaş Y, Orak OS, et al. The effect of music therapy on depression and
38 physiological parameters in elderly people living in a Turkish nursing home: a randomized-
39 controlled trial. *Aging Ment Health* 2017;21:1280–6.doi:10.1080/13607863.2016.1222348
40
41
42
43 30. Gómez Gallego M, Gómez García J. Music therapy and Alzheimer's disease: Cognitive,
44 psychological, and behavioural effects. *Neurologia* 2017;32:300–
45 8.doi:10.1016/j.nrl.2015.12.003
46
47
48 31. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement
49 intervention on the cognitive functions of people with moderate dementia: a randomized
50 controlled trial. *Aging Ment Health* 2018;22:306–15.doi:10.1080/13607863.2016.1251571
51
52
53
54 32. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in
55
56
57
58
59
60

- 1
2
3
4 elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging*
5 *Ment Health* 2017;21:156–62.doi:10.1080/13607863.2015.1093602
6
7
8 33. Bieleninik L, Geretsegger M, Mössler K, et al. Effects of improvisational music therapy vs
9 enhanced standard care on symptom severity among children with autism spectrum disorder:
10 the TIME-A randomized clinical trial. *JAMA* 2017;318(6):525–
11 535.doi:10.1001/jama.2017.9478
12
13
14
15 34. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a
16 controlled randomized study. *Neurol Sci* 2017;38:1485–93.doi:10.1007/s10072-017-3003-9
17
18
19 35. Deng HX, Gao GS, Li ZM, et al. Study of the effect of music therapy in the treatment of
20 elderly patients with diabetes. *China Health Care and Nutrition*
21 2013;23:1186.doi:10.3969/j.issn.1004-7484(s).2013.03.206
22
23
24
25 36. Zhang ZW. Study of the effect of music therapy and psychotherapy adjuvant therapy in the
26 treatment of type 2 diabetes. *China Medicine and Pharmacy* 2011,01:150-
27 1.doi:CNKI:SUN:GYKX.0.2011-18-103
28
29
30
31 37. Solanki MS, Zafar M, Rastogi R. Music as a therapy: role in psychiatry. *Asian J Psychiatr*
32 2013;6:193–9.doi:10.1016/j.ajp.2012.12.001
33
34
35 38. Petrovsky D, Cacchione PZ, George M. Review of the effect of music interventions on
36 symptoms of anxiety and depression in older adults with mild dementia. *Int Psychogeriatr*
37 2015;27:1661–70.doi:10.1017/S1041610215000393
38
39
40
41 39. Aalbers S, Fusar-Poli L, Freeman RE, et al. Music therapy for depression. *Cochrane*
42 *Database Syst Rev* 2017;11:CD004517.doi:10.1002/14651858.CD004517.pub3
43
44
45 40. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and
46 meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*
47 2015;350:g7647.doi:10.1136/bmj.g7647
48
49
50
51 41. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic
52 reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and
53 elaboration. *BMJ* 2009;339:b2700.doi:10.1136/bmj.b2700
54
55
56
57
58
59
60

- 1
2
3
4 42. American Diabetes Association. 9. Pharmacologic approaches to glycemic treatment:
5 standards of medical care in diabetes-2020. *Diabetes Care* 2020;43(Suppl 1):S98–
6 110.doi:10.2337/dc20-S009
7
8
9
10 43. American Diabetes Association. 5. Facilitating behavior change and well-being to improve
11 health outcomes: standards of medical care in diabetes-2020. *Diabetes Care* 2020;43(Suppl
12 1):S48–65.doi:10.2337/dc20-S005
13
14
15 44. Feng Y, Fang Y, Wang Y, et al. Acupoint therapy on diabetes mellitus and its common
16 chronic complications: a review of its mechanisms. *Biomed Res Int*
17 2018;2018:3128378.doi:10.1155/2018/3128378
18
19
20 45. Pérez-Pevida B, Escalada J, Miras AD, et al. Mechanisms underlying type 2 diabetes
21 remission after metabolic surgery. *Front Endocrinol (Lausanne)*
22 2019;10:641.doi:10.3389/fendo.2019.00641
23
24
25 46. Higgins JPT, Thomas J, Chandler J, et al. Cochrane Handbook for Systematic Reviews of
26 Interventions version 6.0 (updated July 2019). Cochrane, 2019. Available:
27 www.training.cochrane.org/handbook [Accessed 3 Jul 2020].
28
29
30
31 47. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of
32 evidence. *J Clin Epidemiol* 2011;64:401–6.doi:10.1016/j.jclinepi.2010.07.015
33
34
35 48. Gonzalez JS, Peyrot M, McCarl LA, et al. Depression and diabetes treatment nonadherence: a
36 meta-analysis. *Diabetes Care* 2008;31:2398–403.doi:10.2337/dc08-1341
37
38
39 49. Trief PM, Foster NC, Chaytor N, et al. Longitudinal changes in depression symptoms and
40 glycemia in adults with type 1 diabetes. *Diabetes Care* 2019;42:1194–201.doi:10.2337/dc18-
41 2441
42
43
44
45 50. Novak M, Mucsi I, Rhee CM, et al. Increased risk of incident chronic kidney disease,
46 cardiovascular disease, and mortality in patients with diabetes with comorbid depression.
47
48
49
50
51
52 51. Lustman PJ, Anderson RJ, Freedland KE, et al. Depression and poor glycemic control: a
53 meta-analytic review of the literature. *Diabetes Care* 2000;23:934–
54
55
56
57
58
59
60

- 1
2
3
4 42.doi:10.2337/diacare.23.7.934
5
6 52. Roopan S, Larsen ER. Use of antidepressants in patients with depression and comorbid
7
8 diabetes mellitus: a systematic review. *Acta Neuropsychiatr* 2017;29:127–
9
10 39.doi:10.1017/neu.2016.54
11
12 53. Croom AM. Music, neuroscience, and the psychology of well-being: a précis. *Front Psychol*
13
14 2012;2:393.doi:10.3389/fpsyg.2011.00393
15
16 54. Abd-Elshafy SK, Khalaf GS, Abo-Kerisha MZ, et al. Not all sounds have negative effects on
17
18 children undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 2015;29:1277–
19
20 84.doi:10.1053/j.jvca.2015.01.005
21
22 55. Yu SJ, Sun ZY, Zhou J, et al. Study of the effect of music therapy and psychological
23
24 intervention in the treatment of diabetic patients. *Hebei med* 2007;13:903–
25
26 6.doi:10.3969/j.issn.1006-6233.2007.08.010
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification P₁	1a	Identify the report as a protocol of a systematic review
Update None	1b	If the protocol is for an update of a previous systematic review, identify as such
Registration P₂	2	If registered, provide the name of the registry (such as PROSPERO) and registration number
Authors:		
Contact P₁	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author
Contributions P₁₄	3b	Describe contributions of protocol authors and identify the guarantor of the review
Amendments None	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments
Support:		
Sources P₁₄	5a	Indicate sources of financial or other support for the review
Sponsor None	5b	Provide name for the review funder and/or sponsor
Role of sponsor or funder None	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol
INTRODUCTION		
Rationale P₄₋₅	6	Describe the rationale for the review in the context of what is already known
Objectives P₆₋₇	7	Provide an explicit statement of the question(s) the review will address with reference to participants P₆₋₇ , interventions P₇ , comparators P₇ , and outcomes P₇ (PICO)
METHODS		
Eligibility criteria P₆₋₇	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review
Information sources P₆₋₈, P₁₁	9	Describe all intended information sources (such as electronic databases P₇₋₈ , contact with study authors P₁₁ , trial registers P₆ or other grey literature sources P₈) with planned dates of coverage P₈
Search strategy P₉	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated

Study records:		
Data management P₁₀	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review
Selection process P₉	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)
Data collection process P₁₀	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators
Data items P₁₀	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications
Outcomes and prioritization P₇	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale
Risk of bias in individual studies P₁₀	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis
Data synthesis P₁₀₋₁₂	15a	Describe criteria under which study data will be quantitatively synthesised P₁₀₋₁₁
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I ² , Kendall's τ) P₁₁
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) P₁₂
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned P₁₁
Meta-bias(es) P₁₁	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)
Confidence in cumulative evidence P₁₂	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.