

**SARS-CoV-2-specific antibody detection in healthcare workers in a UK maternity  
Hospital: Correlation with SARS-CoV-2 RT-PCR results**

Asma Khalil<sup>1,2</sup>, Robert Hill<sup>3,4</sup>, Alison Wright<sup>3,5</sup>, Shamez Ladhani<sup>6,7</sup>, Pat O'Brien<sup>8,9</sup>

<sup>1</sup>Fetal Medicine Unit, St George's Hospital, St George's University of London, Cranmer Terrace, London SW17 0RE, UK

<sup>2</sup>Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute, St George's University of London, Cranmer Terrace, London SW17 0RE, UK

<sup>3</sup>The Portland Hospital for Women and Children, 205-209 Great Portland St, London W1W 5AH, UK

<sup>4</sup>Great Ormond Street Hospital, Great Ormond Street, London WC1N 3JH, UK

<sup>5</sup>Royal free Hospital, London NW3 2QG, UK

<sup>6</sup>Immunisation and Countermeasures Division, Public Health England, London, 61 Colindale Avenue London NW9 5EQ, UK

<sup>7</sup>Paediatric Infectious Diseases Research Group, St. George's University of London, UK

<sup>8</sup>The Royal College of Obstetricians and Gynaecologists, London, 10-18, Union St, London SE1 1SZ, UK

<sup>9</sup>University College London Hospitals NHS Foundation Trust, London, 235 Euston Rd, Bloomsbury, London NW1 2BU, UK

**Corresponding author:**

Professor Asma Khalil MRCOG MD MSc (Epi) DFRS Dip (GUM)

Address:

Fetal Medicine Unit

Department of Obstetrics and Gynaecology

St. George's University Hospitals NHS Foundation Trust

Blackshaw Road

London SW17 0QT

United Kingdom.

Email address: [akhalil@sgul.ac.uk](mailto:akhalil@sgul.ac.uk)

Accepted Manuscript

Dear Editor,

During the ongoing COVID-19 pandemic, staff shortages due to illness, self-isolation and redeployment have been a major challenge. Universal healthcare worker (HCW) testing is potentially useful in ameliorating workforce depletion and reducing asymptomatic spread of SARS-CoV-2. Nasopharyngeal swab polymerase chain reaction (RT-PCR) can diagnose only current or recent infection; testing for antibody responses against SARS-CoV-2 could enhance the ability to expedite reinstatement of services, while ensuring patient and staff safety. Tests are now available for immunoglobulin (Ig) G against the SARS-CoV-2 nucleocapsid protein; the Abbott SARS-CoV-2 IgG ELISA is reported to have high specificity (99.9%) and sensitivity (96.9%).<sup>1</sup> In London, England, HCW at the Portland Hospital for Women and Children are routinely swabbed for SARS-CoV-2 as part of hospital surveillance policy and asked to self-isolate if infected. Between May 15-28, 2020, 190 HCW who had previously had a nasopharyngeal swab for SARS-CoV-2, were screened for SARS-CoV-2 IgG antibodies. Informed consent included acknowledgement that a positive result should not be considered an 'immunity certificate'.

SARS-CoV-2 IgG antibodies were detected in 41 (22%) HCWs, including all 25 [19 (76%) symptomatic, 6 (24%) asymptomatic] previously testing positive for SARS-CoV-2 on nasopharyngeal swab RT-PCR. At the same time, 16/165 (10%) HCWs who tested negative for SARS-CoV-2 on nasopharyngeal swab, of whom 2 (12.5%) had reported COVID-19-like symptoms, were positive for SARS-CoV-2 IgG. Of those positive for IgG, 39% had previously tested negative on nasopharyngeal swab (Figure). Risk factors associated with an increased risk of severe COVID-19 are included in the Figure.

We previously reported that 32% of HCW testing positive for SARS-CoV-2 on nasopharyngeal swab were asymptomatic at the time.<sup>2</sup> Symptomatic and asymptomatic SARS-CoV-2 positive adults have similar viral loads and infectious virus isolation.<sup>3</sup> Our finding that both of these groups developed SARS-CoV-2 IgG antibodies is reassuring.

Of those testing positive for SARS-CoV-2 IgG, 39% had an earlier negative nasopharyngeal swab. Possible explanations are that either infection occurred at an interval before or after the swab test, or the swab RT-PCR gave a false negative result (due to poor swabbing technique, suboptimal storage conditions, delay in testing, or poor sensitivity of nasopharyngeal swabs, reported to be as low as 70%).<sup>4</sup>

The overall prevalence of SARS-CoV-2 IgG (22%) among HCW was higher than in the general population in London (17%) or across the UK (5%).<sup>5</sup> Both symptomatic and asymptomatic infections were associated with SARS-COV-2 IgG antibodies, as were 10% of

HCW with negative nasopharyngeal swabs, despite the majority remaining asymptomatic. These findings are important for modelling SARS-CoV-2 transmission among HCW. Further studies are needed to establish whether these antibodies protect against re-infection and the duration of protection.

### **Potential Conflicts of Interest**

R.H. reports personal fees as Medical Director from The Portland Hospital, outside the submitted work.

### **References**

1. Bryan A, Pepper G, Wener MH, et al. Performance Characteristics of the Abbott Architect SARS-CoV-2 IgG Assay and Seroprevalence in Boise, Idaho. *J Clin Microbiol* 2020 doi: 10.1128/JCM.00941-20. [Epub ahead of print].
2. Khalil A, Hill R, Ladhani S, et al. COVID-19 screening of health-care workers in a London maternity hospital. *Lancet Infect Dis* 2020 doi: 10.1016/S1473-3099(20)30403-5 [Epub ahead of print].
3. Gandhi M, Yokoe DS, Havlir DV. Asymptomatic Transmission, the Achilles' Heel of Current Strategies to Control Covid-19. *N Engl J Med* 2020; 382: 2158-60. doi:10.1056/NEJMe2009758.
4. Yang Y, Yang M, Shen C, et al. Evaluating the accuracy of different respiratory specimens in the laboratory diagnosis and monitoring the viral shedding of 2019-nCoV infections. *MedRxiv* 2020 doi: 10.1101/2020.02.11.20021493. 2020.02.11.20021493.
5. <https://www.livemint.com/news/world/-17-londoners-5-of-uk-residents-have-covid-19-antibodies-11590148840735.html>

## **Acknowledgment**

We would like to thank Ms Jane Whitney-Smith (Chief Executive Officer, The Portland Hospital for Women and Children) and Gregory Bale (Acting Chief Operating Officer, The Portland Hospital for Women and Children) for their invaluable support.

Accepted Manuscript

Figure 1




Figure . SARS-CoV-19 RT-PCR and IgG test results among 190 healthcare workers and their characteristics

Accepted