# **Supplementary Online Content**

Palumbo SA, Adamson KM, Krishnamurthy S, et al. Assessment of probable opioid use disorder using electronic health record documentation. *JAMA Netw Open.* 2020;3(9):e2015909. doi:10.1001/jamanetworkopen.2020.15909

eAppendix. Supplemental Methods

eTable 1. ICD 9/10 Codes Used for Each Phenotype

**eTable 2.** Demographic and Health Record Summary Data for GMMP-M and GMMP-V Subgroups

eTable 3. EHR Search Categories and Number of Patients With Each Category

eTable 4. Specific Toxicology Screen Violations by Subtype

eTable 5. Frequency of Positive Search Criteria

This supplementary material has been provided by the authors to give readers additional information about their work.

#### **eAppendix.** Supplemental Methods

# eMethods 1: Agreement terms of the Patient Drug Monitoring Program contracts.

Patients that sign this contract agree to the following terms:

- (1) To attend regularly scheduled appointments.
- (2) To take medication only as prescribed.
- (3) To agree to one provider and fills at one specified pharmacy.
- (4) To pick up own prescription at pharmacy or by authorized persons (named in agreement).
- (5) To not to obtain other controlled substances, including opioid pain medicines, controlled stimulants or anti-anxiety medicines from any other provider.
- (6) To not borrow or accept medication from friends, use street or illegal drugs, and/or sell/give medication to another person.
- (7) To not make any attempt to forge or alter any prescriptions.
- (8) To safeguard medications to prevent lost or theft.
- (9) To submit to random drug screens and pill counts.

Patients who violate the terms of this agreement are denoted with a separate EPIC code.

Examples of violations of the above are as follows:

- (1) Cancellation or no show for two or more appointments (considered a cancellation if less than one working days' notice).
- (2) Increasing or decreasing dose without discussing with provider.
- (3) Seeking and/or filling prescription(s) from another provider or multiple pharmacies.
- (4) Allowing someone other than self or authorized person to pick up prescription.
- (5) Obtaining other controlled substances, including those legally prescribed by other physicians and/or illegal acquisition of prescription or street drugs.
- (6) To forge or alter prescriptions.
- (7) Repeated reports of lost or stolen medication.
- (8) Refusal to submit to urine toxicology screen and/or pill counts.
- (9) Positive urine toxicology screen for substances not prescribed.

### eMethods 2: Opioid drugs included and equivalent dose determination:

All opioid prescription records were also drawn from the EHR, and morphine milligram equivalent (MME) dose was calculated using the global rPH calculator (http://www.globalrph.com/narcotic.cgi). Opioid drugs included were hydromorphone, methadone, oxymorphone, oxycodone, morphine sulfate, hydrocodone, tapentadol, tramadol, codeine, and fentanyl.

#### **eMethods 3: Chart Review Variables:**

- (A) Demographic and health record information relevant to the GMMP program for each group, included:
- (A1) Length of Electronic Health Record (EHR),
  - The duration of the patient's health record was quantified in number of days. This was determined by calculating the number of days between the patient's very first encounter with the health system and the most recent visit, as of May 31, 2018.
- (A2) Current opioid prescription with Morphine Milligram Equivalence (MME),

Current opiate prescription and dosage, and other controlled substances were recorded. Morphine milligram equivalence (MME) was calculated based on current prescription. Current prescriptions are located under the medications tabs. To identify current opioid prescriptions, the medication list was filtered by pharmaceutical class to only show analgesic narcotics. This same process was conducted to identify current controlled substance. To do so, the medication list was filtered by class and then by class II, III, and IV. MME was calculated using standard conversion procedure.

- (A3) Number of Emergency Room (ER) visits over the course of the patient's record: Emergency rooms visits were searched and number summed from the beginning of the patient's health record until their most recent visit.
  - (B) Criteria related to an OUD DSM diagnosis:
- (B1) Vocational interference due to drug use or pain

A positive vocational interference may indicate that the patient is unable to work due to pain, or that their prescription opioid use is impacting their work. Health records were searched for terms that indicated if a patient's occupation was impacted by their use of opioids or pain. Terms that were searched included: occupation, work, job, boss, manager, too much pain, can't work.

• (B2) Patient Disability Status

Health records, including notes and encounters were searched for terms that indicated whether the patient was disabled. A positive disability status may indicate that the patient's pain was so great that it interfered with their occupation. If it was mentioned that a patient was planning to file for disability or was waiting to hear if they were approved, they were categorized as a no. Terms that were searched included: disability, too much pain, unemployment, can't work, paperwork.

- -Yes- disability indicated in the EHR
- -No- Explicitly states the patient is not disabled
- -No Record- No mention of ability status
- (B3) Whether weaning was described as unsuccessful or difficult

If it was indicated that the patient had been weaned off an opioid, a further search was conducted to determine if the weaning was difficult or had complications.

- -Yes- the record indicates that weaning was difficult or unsuccessful for the individual
- -No- weaning was mentioned for the individual but was successful and/or not difficult and complicated
- -No record- weaning was mentioned though the success and difficulty of the action was not recorded
- N/A- weaning was not mentioned for this individual thus cannot be difficult or unsuccessful
- (B4) Positive toxicology screen for opioids other than prescribed

In order to verify that the patient was following the conditions of the GMMP agreement, health records were searched for urine toxicology screens to ensure that medications that the patient was prescribed were present in their urine and non-prescribed medications were not present. Number of toxicology screens varied between patients ranging from no toxicology screens to 50+. Toxicology screens were identified under the lab tests section. Because there are several tests, a filter was used to identify toxicology screens. Tests that were searched were: toxicology screens, urine toxicology screens, and opiates

confirmatory. The only toxicology screens that were reviewed were those that were collected during the GMMP agreement period. After the toxicology screen result was identified, prescribed medications at the time of the collection were verified. To do so, the medication list was opened and filtered by pharmaceutical class then by analgesics-narcotics. The date of the toxicology screen was identified and then matched to the closest date of prescription. The same process was conducted for controlled substances. The date of the toxicology screen was also identified if a medication was not present in the system in order to review the medications that we prescribed at that time.

# • (B5) Lost pills mentioned

Several scenarios consistently occurred in the chart review where patients reported missing pills such as having them stolen, misplaced, lost, or ruined due to accidents such as dropping them in water. Terms that were searched included: lost, missing, stolen, misplace, robbed, police report, and gone.

- -Yes- the record mentions lost pills
- -No- there was no mention of lost pills

# • (B6) Multiple Opioid Prescribers Mentioned

To search if a patient was receiving more than one opioid prescription, the medication list was searched for more than one providers or prescription during the same time period. If a patient received an opioid prescription from the emergency room while actively being prescribed an opioid, that was identified as multiple prescribers. If it was not evident in the medication list that the patient was receiving multiple prescriptions, terms were searched in the notes to identify if a prescriber identified that the patient was receiving more than one prescription or going to the more than one pharmacy. Terms that were searched included: multiple, various, pharmacy, and prescribers.

- -Yes- Multiple opioid prescribers were mentioned in the record
- -No- Multiple opioid prescribers were not mentioned in the record

# • (B7) Multiple Pharmacies Mentioned

To search if a patient was receiving more than one opioid prescription, the medication list was searched for more than one provider or prescription during the same time period. If a patient received an opioid prescription from the emergency room while actively being prescribed an opioid, that was identified as multiple prescribers. If it was not evident in the medication list that the patient was receiving multiple prescriptions, terms were searched in the notes to identify if a prescriber identified that the patient was receiving more than one prescription or going to the more than one pharmacy. Terms that were searched included: multiple, various, pharmacy, and prescribers.

- -Yes- Multiple Pharmacies were mentioned
- -No- Multiple Pharmacies were not mentioned

# • (B8) Early Prescription Refills Mentioned

Was it ever mentioned in the patient's health record if the patient requested an early an medication refill? Terms that were searched included: refill, too soon, early, and ran out.

- -Yes- Early refills were mentioned
- -No- Early refills were not mentioned

### • (B9) Opioid overdose mentioned

Terms that were searched included: overdose, intoxicated, unconscious, and intravenous.

- -Yes- Opioid overdose was mentioned
- -No- Opioid overdose was not mentioned

#### • (B10) Substance abuse mentioned

The patient was classified as "yes" for substance if illicit drugs that were not separately screened for (i.e. alcohol, marijuana, tobacco), were in the health record. Terms that were searched included: abuse, substance abuse, heroin, street drugs, detox, selling, dealing, and high.

- -Yes- substance abuse was mentioned
- -No- substance abuse was not mentioned
- (B11) Hazardous situation as a result of opioid mentioned

An instance was considered to be hazardous if there was a suicide attempts via overdose, selling prescription, or ingesting prescription in conjunction with illicit drugs such as cocaine.

- -Yes- such a situation was noted in the record
- -No- a hazardous situation due to the opioid was not mentioned
- (B12) Interpersonal or legal issues as a result of opioid mentioned

An instance was considered to be an interpersonal or legal issue if the police were involved in any capacity, if lawyers were mentioned, if jail or prison occurred, or if children services were mentioned.

- -Yes- issues were mentioned
- -No- issues were not mentioned
- (B13) Medical issues as a result of opioid mentioned

Medical issues that were considered included narcotic abuse or dependence, withdrawal symptoms, constipation secondary to opioid use, or diseases contracted through drug use such as hepatitis C.

- -Yes- medical issues due to the opioid were mentioned
- -No- medical issues due to the opioid were not mentioned
- (B14) Craving mentioned

Was it ever mentioned in the health record if the patient exhibited craving? The health record was first searched for the SOAPP-R (Screener and Opioid Assessment for Patients with Pain-Revised) which provides a risk score for opioid abuse. A patient was considered to exhibit craving symptoms if it was mentioned in their record that they were experiencing withdrawal symptoms or if it was specifically mentioned that they were craving the drug.

- -Yes- craving was mentioned
- -No- craving was not mentioned
- (B15) Provider mentioned drug-seeking behavior

Was it ever mentioned in the health record if the patient exhibited drug-seeking behavior? The health record was first searched for the SOAPP-R (Screener and Opioid Assessment for Patients with Pain-Revised) which provides a risk score for opioid abuse. A patient was considered to exhibit drug-seeking behavior if it was mentioned in the physician's notes as so. Terms that were searched from drug-seeking behavior included: seeking, narcotic seeking, and drug seeking.

- -Yes- drug-seeking behavior was mentioned
- -No- drug-seeking behavior was not mentioned

eTable 1. ICD 9/10 Codes Used for Each Phenotype

ICD 9	ICD 10	Diagnosis	
Anxiety			
300	F41.9	Anxiety disorder, unspecified	
300.01	F41.0	Panic disorder [episodic paroxysmal anxiety] without agoraphobia	
300.02	F41.1	Generalized anxiety disorder	
300.09	F41.3	Other mixed anxiety disorders	
300.09	F41.8	Other specified anxiety disorders	
Depression			
296.3	F33.40	Major depressive disorder, recurrent, in remission, unspecified	
296.3	F33.9	Major depressive disorder, recurrent, unspecified	
296.31	F33.0	Major depressive disorder, recurrent, mild	
296.32	F33.1	Major depressive disorder, recurrent, moderate	
296.33	F33.2	Major depressive disorder, recurrent severe without psychotic features	
296.34 or 298.0	F33.3	Major depressive disorder, recurrent, severe with psychotic symptoms	
296.35	F33.41	Major depressive disorder, recurrent, in partial remission	
296.36	F33.42	Major depressive disorder, recurrent, in full remission	
296.99	F33.8	Other recurrent depressive disorders	
311	F32.9	Depressive disorder, not elsewhere classified	
Alcohol			
291	F10.121	Alcohol abuse with intoxication delirium	
291	F10.231	Alcohol dependence with withdrawal delirium	
291.3	F10.251	Alcohol dependence with alcohol-induced psychotic disorder with hallucinations	
291.4	F10.929	Alcohol use, unspecified with intoxication, unspecified	
291.81	F10.230	Alcohol dependence with withdrawal, uncomplicated	
291.81	F10.239	Alcohol dependence with withdrawal, unspecified	
291.89	F10.159	Alcohol abuse with alcohol-induced psychotic disorder, unspecified	
291.89	F10.288	Alcohol dependence with other alcohol-induced disorder	
291.9	F10.19	Alcohol abuse with unspecified alcohol-induced disorder	
291.9	F10.29	Alcohol dependence with unspecified alcohol-induced disorder	
291.9	F10.99	Alcohol use, unspecified with unspecified alcohol-induced disorder	
303	F10.229	Acute alcoholic intoxication in alcoholism, unspecified	
303.01	F10.229	Acute alcoholic intoxication in alcoholism, continuous	
303.9	F10.20	Other and unspecified alcohol dependence	
303.9	F10.20	Other and unspecified alcohol dependence, unspecified	
303.9	1.10.20	and the second and th	
303.9	F10.20	Alcohol dependence, uncomplicated	

303.92	F10.20	Other and unspecified alcohol dependence, episodic			
303.93	F10.21	Other and unspecified alcohol dependence, in remission			
303.93	F10.21	Alcohol dependence, in remission			
305	F10.10	Alcohol abuse, unspecified			
305.01	F10.10	Alcohol abuse, continuous			
305.02	F10.10	Alcohol abuse, episodic			
305.03	F10.11	Alcohol abuse, in remission			
303 or 303.01	F10.229	Alcohol dependence with intoxication, unspecified			
303.00 or 303.01 or 303.02 or 303.03	F10.220	Alcohol dependence with intoxication, uncomplicated			
305 or 305.01 or 305.02	F10.10	Alcohol abuse, uncomplicated			
305.00 or 305.01 or 305.02	F10.120	Alcohol abuse with intoxication, uncomplicated			
305.00 or 305.01 or 305.02	F10.129	Alcohol abuse with intoxication, unspecified			
Nicotine					
292.89	F17.218	Nicotine dependence, cigarettes, with other nicotine-induced disorders			
292.89	F17.228	Nicotine dependence, chewing tobacco, with other nicotine- induced disorders			
292.9	F17.209	Nicotine dependence, unspecified, with unspecified nicotine-induced disorders			
292.9	F17.219	Nicotine dependence, cigarettes, with unspecified nicotine- induced disorders			
292.9	F17.229	Nicotine dependence, chewing tobacco, with unspecified nicotine-induced disorders			
305.1	F17.200	Tobacco use disorder			
305.1	F17.200	Nicotine dependence, unspecified, uncomplicated			
305.1	F17.201	Nicotine dependence, unspecified, in remission			
305.1	F17.210	Nicotine dependence, cigarettes, uncomplicated			
305.1	F17.220	Nicotine dependence, chewing tobacco, uncomplicated			
305.1	F17.290	Nicotine dependence, other tobacco product, uncomplicated			
*	F17.21	Nicotine dependence, cigarettes			
Opioid					
304.7	*	Combinations of opioid type drug with any other drug dependence, unspecified			
304.71	*	Combinations of opioid type drug with any other drug dependence, continuous			
304.73	*	Combinations of opioid type drug with any other drug dependence, in remission			
304.8	*	Combinations of drug dependence excluding opioid type drug, unspecified			
304.81	*	Combinations of drug dependence excluding opioid type drug, continuous			
304.82	*	Combinations of drug dependence excluding opioid type drug, episodic			

304.83	*	Combinations of drug dependence excluding opioid type drug, in remission			
305.5	F11.10	Opioid abuse, unspecified			
305.51	F11.10	Opioid abuse, continuous			
305.52	F11.10	Opioid abuse, episodic			
305.53	F11.11	Opioid abuse, in remission			
Other Substance Abuse					
292	F15.93	Other stimulant use, unspecified with withdrawal			
292	F19.930	Other psychoactive substance use, unspecified with withdrawal, uncomplicated			
292	F19.939	Other psychoactive substance use, unspecified with withdrawal, unspecified			
292.11	F19.950	Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder with delusions			
292.12	F19.951	Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder with hallucinations			
292.2	F12.120	Cannabis abuse with intoxication, uncomplicated			
292.2	F15.920	Other stimulant use, unspecified with intoxication, uncomplicated			
292.84	F19.94	Other psychoactive substance use, unspecified with psychoactive substance-induced mood disorder			
292.85	F19.982	Other psychoactive substance use, unspecified with psychoactive substance-induced sleep disorder			
292.89	F15.959	Other stimulant use, unspecified with stimulant-induced psychotic disorder, unspecified			
292.89	F19.959	Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder, unspecified			
292.9	F12.19	Cannabis abuse with unspecified cannabis-induced disorder			
292.9	F19.99	Other psychoactive substance use, unspecified with unspecified psychoactive substance-induced disorder			
304.1	F13.20	Sedative, hypnotic or anxiolytic dependence, unspecified			
304.11	F13.20	Sedative, hypnotic or anxiolytic dependence, continuous			
304.13	F13.21	Sedative, hypnotic or anxiolytic dependence, in remission			
304.13	F13.21	Sedative, hypnotic or anxiolytic dependence, in remission			
304.2	F14.20	Cocaine dependence, unspecified			
304.2	F14.20	Cocaine dependence, uncomplicated			
304.23	F14.21	Cocaine dependence, in remission			
304.23	F14.21	Cocaine dependence, in remission			
304.3	F12.20	Cannabis dependence, unspecified			
304.31	F12.20	Cannabis dependence, continuous			
304.33	F12.21	Cannabis dependence, in remission			
304.4	F15.20	Amphetamine and other psychostimulant dependence, unspecified			

304.41	F15.20	Amphetamine and other psychostimulant dependence, continuous		
304.43	F15.21	Amphetamine and other psychostimulant dependence, in remission		
304.6	F19.20	Other specified drug dependence, unspecified		
304.61	F19.20	Other specified drug dependence, continuous		
304.63	F19.21	Other specified drug dependence, in remission		
304.63	F19.21	Other psychoactive substance dependence, in remission		
304.9	*	Unspecified drug dependence, unspecified		
304.91	*	Unspecified drug dependence, continuous		
304.92	*	Unspecified drug dependence, episodic		
304.93	*	Unspecified drug dependence, in remission		
305.2	F12.10	Cannabis abuse, unspecified		
305.21	F12.10 or F12.90	Cannabis abuse, continuous		
305.22	F12.10 or F12.90	Cannabis abuse, episodic		
305.23	F12.11	Cannabis abuse, in remission		
305.33	F16.11	Hallucinogen abuse, in remission		
305.4	F13.10	Sedative, hypnotic or anxiolytic abuse, unspecified		
305.41	F13.10	Sedative, hypnotic or anxiolytic abuse, continuous		
305.42	F13.10	Sedative, hypnotic or anxiolytic abuse, episodic		
305.6	F14.10	Cocaine abuse, unspecified		
305.61	F14.10	Cocaine abuse, continuous		
305.63	F14.11	Cocaine abuse, in remission		
305.7	F15.10	Amphetamine or related acting sympathomimetic abuse, unspecified		
305.7	F15.10	Other stimulant abuse, uncomplicated		
305.73	F15.11	Amphetamine or related acting sympathomimetic abuse, in remission		
305.8	F19.10	Antidepressant type abuse, unspecified		
305.8	F19.10	Other psychoactive substance abuse, uncomplicated		
305.8	F19.90	Other psychoactive substance use, unspecified, uncomplicated		
305.9	F19.10 or F18.10	Other, mixed, or unspecified drug abuse, unspecified		
305.91	F19.10 or F18.10	Other, mixed, or unspecified drug abuse, continuous		
305.92	F19.10 or F18.10	Other, mixed, or unspecified drug abuse, episodic		
305.93	F19.11 or F18.11	Other, mixed, or unspecified drug abuse, in remission		
292.0 or 304.10	F13.239	Sedative, hypnotic or anxiolytic dependence with withdrawal, unspecified		
292.0 or 304.60	F19.230	Other psychoactive substance dependence with withdrawal, uncomplicated		

304.1 or 304.11	F13.20	Sedative, hypnotic or anxiolytic dependence, uncomplicated
304.3 or 304.31	F12.20	Cannabis dependence, uncomplicated
304.4 or 304.41	F15.20	Other stimulant dependence, uncomplicated
304.6 or 304.61	F19.20	Other psychoactive substance dependence, uncomplicated
305.2 or 305.21 or 305.22	F12.10	Cannabis abuse, uncomplicated
305.21 or 305.2 or 305.22	F12.90	Cannabis use, unspecified, uncomplicated
305.4 or 305.41 or 305.42	F13.10	Sedative, hypnotic or anxiolytic abuse, uncomplicated
305.6 or 305.51 or 305.62	F14.10	Cocaine abuse, uncomplicated

eTable 2. Demographic and Health Record Summary Data for GMMP-M and GMMP-V Subgroups						
	GMMP-M N= 14,866	GMMP-V N=1,387	<i>P</i> Value	Test Statistic	Effect Size	95% CI
Sample Demographics						
Male	6,327	620				
Female	8,539	767	0.10	1.64	0.02	(0.0045, 0.0494)
Age	52.3	47.2	<0.001	15.60	5.15	(4.504, 5.799)
ВМІ	31.8	30.5	<0.001	5.75	1.27	(0.8374, 1.7040)
Health Record Data						(
EHR length in days	4231	4004	<0.001	4.28	226	(122.8, 330.6)
ER visits	8.3	11.1	<0.001	-4.34	2.77	(1.518, 4.022)
Daily MME	52.0	48.8	0.01	2.44	3.20	(0.6336, 5.776)
Psychiatric Codes						
Depression	4902	591	<0.001	35.86	0.08	(0.0513, 0.1048)
Anxiety	5896	702	<0.001	42.09	0.09	(0.0606, 0.1150)
Addiction Codes						
Alcohol	424	74	<0.001	21.97	0.02	(0.0105, 0.0345)
Nicotine	4178	622	<0.001	139	0.15	(0.1211, 0.1747)
Opioids	243	51	<0.001	25.5	0.02	(0.0087, 0.0288)
Other Substance Abuse	451	130	<0.001	134	0.06	(0.0439, 0.0745)

<sup>\*\*\*\*</sup>GMMP-M and GMMP-V groups were based on all available people with each designation and therefore do not represent matched groups.

eTable 3. EHR Search Categories and Number of Patients With Each Category

	Patient Group		
	Controls	GMMP- M	GMMP- V
Vocational Interference Due to Drug Use or Pain	16	21	7
Disabled	34	58	44
Weaning Described as Unsuccessful or Difficult	1	10	13
Positive Tox Screen for Opioids other than Prescribed	6*	7*	30*
Lost Pills	3	23	31
Multiple Opioid Prescribers	11	11	47
Multiple Pharmacies	2	6	10
Early Prescription Refills	9	52	64
Opioid Overdose	4	7	6
Substance Abuse	18	28	51
Hazardous Situation as Result of Opioid	4	20	23
Interpersonal or Legal Issues as Result of Opioid	4	19	14
Medical Issues as Result of Opioid	15	41	29
Craving	0	2	4
Provider Mentioned Drug-seeking Behavior	5	10	21

<sup>\*</sup>These numbers correspond to individuals that tested positive for an unprescribed opioid. Toxicology screens that are negative for a prescribed opioid and/or positive for unprescribed controlled substances are also considered violations. For complete breakdown of other toxicology screen violations, see eTable 4. Gray boxes indicate other categories that would violate the terms of the GMMP contract.

eTable 4. Specific Toxicology Screen Violations by Subtype

	Controls	GMMP- M	GMMP- V
Positive for unprescribed opioid	6	7	30
Positive for unprescribed non-opioid	15	16	52
Negative for prescribed opioid	0	13	48
Non-compliance with urine tox screen	0	0	8
Positive for unprescribed medication*	17	20	64
Negative for prescribed medication*	0	17	51
Both (+) for unprescribed and (-) for prescribed	0	10	35

<sup>\*</sup>Patients sometimes tested positive for more than one substance, thus values for all patients that test positive for unprescribed opioids/non-opioids, for example, do not add up to total # in 'positive for unprescribed medication'.

eTable 5. Frequency of Positive Search Criteria

# of Positive	Patient Group			
Categories	Controls	GMMP-M	GMMP-V	
0	124	11	0	
1	50	18	13	
2	11	19	17	
3	6	18	22	
4	4	12	13	
5	4	2	13	
6	1	5	6	
7	0	9	6	
8	0	3	7	
9	0	1	1	
10	0	1	1	
11	0	1	1	
12	0	0	0	
13	0	0	0	
14	0	0	0	
15	0	0	0	