

Supplementary Online Content

Desai AS, Maclean T, Blood AJ, et al. Remote optimization of guideline-directed medical therapy in patients with heart failure with reduced ejection fraction. *JAMA Cardiol*. Published online September 16, 2020. doi:10.1001/jamacardio.2020.3757


eFigure. Patient Selection

eTable. Adverse Events, by Study Group

eMethods. Physician-LIP Collaborative Drug Therapy Management Plan

This supplementary material has been provided by the authors to give readers additional information about their work.

eFigure. Patient Selection for the Remote Management Intervention



eTable. Adverse Events of Interest, by study group

	REMOTE GDMT OPTIMIZATION (N=197)	USUAL CARE (N=831)	P-VALUE
Symptomatic Hypotension* – n (%)	7 (3.6)	34 (4.2)	0.84
Hyperkalemia (K\geq5.5 meq/L) – n (%)	5 (2.5)	59 (9.9)	0.01
Hyperkalemia (K\geq6.0 meq/L) – n (%)	4 (2.0)	35 (4.2)	0.21
Rise in Serum Cr > 0.3 mg/dL – n (%)	18 (9.1)	3 (0.4)	<0.001
Cardiovascular Hospitalization – n (%)	18 (9.1)	92 (11.1)	0.52
Death – n (%)	1 (0.5)	8 (0.9)	0.84

*Symptomatic hypotension defined as SBP<100 mm Hg in association with dizziness/lightheadedness