

Electronic Supplementary Material

Prebiotic effect of inulin-type fructans on fecal microbiota and short-chain fatty acids in type 2 diabetes: A randomized controlled trial

European Journal of Nutrition

Eline Birkeland^{1,2}, Sedegheh Gharagozlian¹, Kåre I. Birkeland^{2,3}, Jørgen Valeur^{4,5}, Ingrid Måge⁶, Ida Rud⁶, Anne-Marie Aas^{1,2}

Ida Rud and Anne-Marie Aas share last authorship

¹Section of Nutrition and Dietetics, Department of Clinical Service, Division of Medicine, Oslo University Hospital, Norway, ²Institute of Clinical Medicine, University of Oslo, Norway ³Department of Transplantation Medicine, Oslo University Hospital, Norway, ⁴Department of Gastroenterology, Oslo University Hospital, Oslo, Norway, ⁵Unger-Vetlesen Institute, Lovisenberg Diaconal Hospital, Oslo, Norway, ⁶Nofima - Norwegian Institute of Food, Fisheries and Aquaculture Research, Ås, Norway.

Corresponding author: Eline Birkeland, eline.birkeland@ous-hf.no

Online Resource 6 SCFA baseline concentrations before and after washout¹

	Baseline concentrations		
	Before washout	After washout	<i>p</i> value
Total SCFA (mmol/kg)	62.70 (51.91-78.62)	62.67 (49.17-82.95)	0.91
Acetic acid (mmol/kg)	36.00 (30.59-43.23)	36.71 (26.47-41.15)	0.39
Propionic acid (mmol/kg)	11.02 (8.15-15.69)	11.06 (6.89-15.99)	0.98
Butyric acid (mmol/kg)	9.21 (7.29-16.20)	11.95 (8.49-15.03)	0.44

¹Data are median (25th-75th percentils). Wilcoxon Signed Rank Test. Significant differences in bold