

Serotonin Transporter Protein in Autopsied Brain of Chronic Users of Cocaine

Junchao Tong^{1,2,*}, Jeffrey H. Meyer³, Isabelle Boileau⁴, Lee-Cyn Ang⁵, Paul J. Fletcher⁶, Yoshiaki Furukawa⁷, and Stephen J. Kish²

¹ Preclinical Imaging, Brain Health Imaging Centre, Centre for Addiction and Mental Health, Toronto, ON, Canada

² Human Brain Laboratory, Brain Health Imaging Centre, Centre for Addiction and Mental Health, Toronto, ON, Canada

³ Brain Health Imaging Centre and Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, ON, Canada

⁴ Addiction Imaging Research Group, Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, ON, Canada

⁵ Division of Neuropathology, London Health Sciences Centre, University of Western Ontario, London, ON, Canada

⁶ Section of Biopsychology, Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health; Department of Psychology, University of Toronto, Toronto, ON, Canada

⁷ Department of Neurology, Juntendo Tokyo Koto Geriatric Medical Center, and Faculty of Medicine, University & Post Graduate University of Juntendo, Tokyo, Japan

***Corresponding author at:** Preclinical Imaging, Brain Health Imaging Centre, Centre for Addiction and Mental Health, 250 College Street, Toronto, ON, Canada M5T 1R8. Telephone: 416-535-8501 ext. 36422; E-mail: Junchao.tong@camh.ca

Supplementary Figure: Scatter plots of levels of serotonin transporter (SERT) in brain of control subjects and users of cocaine and heroin who were tested negative (-alcohol) or positive (+alcohol) for blood alcohol at the time of autopsy. *P < 0.05, +alcohol vs. -alcohol (Mann-Whitney U-test).

