

S10 Table. The pharmacokinetic parameters of praziquantel (PZQ), its enantiomers, and the major metabolite in Healthy Normal Volunteers (HNV) after the administration of racemic PZQ with drug ‘B’ to evaluate potential drug-drug interactions (DDIs). The brands, when named, were listed in subscript; BILT: Biltricide, DIS: Distocide. The superscript descriptions show the additional information of the study: (a): Egyptian Population, (b): CHQ dosed 2h before PZQ, (c): CIM dosed 1h prior to PZQ, (d): Without H₂O. Unless otherwise stated the following acronyms represent; AUC: Area under the plasma concentration-time curve, B (AUC_{0-inf}): Area under the plasma concentration-time curve extrapolated from time zero to infinity, t_{1/2}: Elimination half-life, C_{max}: Peak plasma concentration, T_{max}: Time to reach peak plasma concentration following drug administration.

Drug Administered (Model)	Drug Measured	PZQ Dose (mg/kg)	B Dose (mg/kg)	N	t _{1/2} (h)	T _{max} (h)	C _{max} (ng/ml)	AUC (ng/ml/h)
PZQ [1]	PZQ	20						
	HNV _(BILT) ^(d)			10	0.97	1.65	846	1303
	HNV _(DIS) ^(d)			10	1.24	2.60	558	2133
PZQ + BIC [2]	PZQ	20	2(g)					
	HNV ^(a)			10	2.15	1.85	563	951
	HNV _(DIS) ^(a)			10	4.11	1.20	384	956
PZQ + CIM [2, 3]	PZQ	20	800					
	HNV ^(a)			10	1.50	1.70	770	2190
	HNV _(DIS) ^(a)			10	2.66	1.65	835	2428
	HNV ^(c)	(3x25)	400	1	1.80			15770
	HNV ^(c)			1	2.40			21690
	HNV ^(c)			1	1.80			15770
	HNV ^(c)			1	2.30			8340
	HNV ^(c)			1	2.40			8950
	HNV ^(c)			1	1.90			10370
	HNV ^(c)			1	2.20			16180
	HNV ^(c)			1	2.40			16580
PZQ + GLUC [2]	PZQ	20	50(g)					
	HNV ^(a)			10	2.70	2.10	373	1130
	HNV _(DIS) ^(a)			10	2.46	2.11	449	1094
PZQ + KTZ [4]	PZQ	20	500(mg)					
	HNV			29	5.16	2.13	2266	6172 (ng*h/mL) ^B
PZQ _(BILT) + CHQ ^[5]	PZQ	40	600					
	HNV ^(b)			1	1.41	3.00	430	1620
	HNV ^(b)			1	1.54	1.75	2100	9420
	HNV ^(b)			1	0.99	2.00	330	1420

<i>HNV</i> ^(b)	1	2.57	1.50	400	2200
<i>HNV</i> ^(b)	1	2.82	2.00	1890	10080
<i>HNV</i> ^(b)	1	1.39	1.25	78	360
<i>HNV</i> ^(b)	1	2.57	2.00	4380	35920
<i>HNV</i> ^(b)	1	1.90	2.00	950	4060
<i>HNV</i> ^(b)	8	1.81	1.93	880	4170
PZQ + ABZ [6]	1500	400			
<i>HNV</i>	(S)-PZQ	9	1.67	2.76	450
<i>HNV</i>	(R)-PZQ	9	1.49	2.73	240
<i>HNV</i>	(S)-4-OH-PZQ	9	2.14	3.27	1070
<i>HNV</i>	(R)-4-OH-PZQ	9	1.84	2.77	790
					5380 ^B

References

1. Metwally A, Bennett J, Botros S, Ebeid F, el attar Gel D. Impact of drug dosage and brand on bioavailability and efficacy of praziquantel. Pharmacological research. 1995;31(1):53-9. Epub 1995/01/01. PubMed PMID: 7784306.
2. Metwally A, Bennett JL, Botros S, Ebeid F. Effect of cimetidine, bicarbonate and glucose on the bioavailability of different formulations of praziquantel. Arzneimittel-Forschung/Drug Research. 1995;45-1(4):516-8. PubMed PMID: WOS:A1995QU77700014.
3. Jung H, Medina R, Castro N, Corona T, Sotelo J. Pharmacokinetic study of praziquantel administered alone and in combination with cimetidine in a single-day therapeutic regimen. Antimicrobial agents and chemotherapy. 1997;41(6):1256-9. doi: 10.1128/aac.41.6.1256. PubMed PMID: WOS:A1997XB71200009.
4. Nleya L, Thelingwani R, Li XQ, Cavallin E, Isin E, Nhachi C, et al. The effect of ketoconazole on praziquantel pharmacokinetics and the role of CYP3A4 in the formation of X-OH-praziquantel and not 4-OH-praziquantel. European journal of clinical pharmacology. 2019;75(8):1077-87. doi: 10.1007/s00228-019-02663-8. PubMed PMID: WOS:000475794400006.
5. Masimirembwa CM, Naik YS, Hasler JA. The effect of chloroquine on the pharmacokinetics and metabolism of praziquantel in rats and in humans. Biopharmaceutics & Drug Disposition. 1994;15(1):33-43. doi: 10.1002/bdd.2510150103. PubMed PMID: WOS:A1994MT3510002.
6. Lima RM, Ferreira MAD, Carvalho T, Fernandes BJD, Takayanagui OM, Garcia HH, et al. Albendazole-praziquantel interaction in healthy volunteers: kinetic disposition, metabolism and enantioselectivity. British journal of clinical pharmacology. 2011;71(4):528-35. doi: 10.1111/j.1365-2125.2010.03874.x. PubMed PMID: WOS:000288249500008.