Data collection tool

D. High (≥5001)

Data conection tool				
I. S	Socio-demographic characteristics of the pa	tients		
Patient Study Number: Age in year:		Age in year:		
Se	Sex			
A.	. Male			
B.	. Female			
Ma	Iarital status			
A.	. Single			
B.	. Married			
C.	. Divorced			
D.	. Widowed			
Occupational status				
A.	. Farmer			
B.	. Employed			
C.	. Merchant			
D.	. Housewife			
E.	. Retired			
Educational Status				
A.	. No formal education			
B.	. Primary (1–8)			
C.	. Secondary (9–12)			
D.	. College/University			
Monthly Family Income (in Ethiopian Birr)				
A	A. Very low (≤860)			
В	B. Low (861-3000)			
C	C. Average (3001-5000)			

History of smoking status			
	A. Smoker		
	B. Non-smoker		
History of alcohol consumption			
	A. Yes		
	B. No		
II.	Clinical characteristics of the study participants		
1.	Duration of the disease (years):		
2.	Prior history of hospitalization: Yes No		
3.	Presence of concurrent comorbidities other than hypertension: Yes		
	No If yes, (specify):		
	Fasting blood sugar level of the patient in the last visit (mg/dL):		
	Blood pressure of the patient in the last visit (mmHg):		
	Treatment regimens of the patient (drug, dose, frequency and duration of treatment)		
7.	Number of medications per patient:		
8.	Is there any drug therapy problem present? YesNo If yes, specify the number		
9.	Is there any need for additional drug therapy? Yes No If yes, what is the cause of this problem?		
	A. Preventive drug therapy required		
	B. A medical condition that requires treatment		

C. To attain synergistic effect

10.	. Is there any unnecessary drug therapy in the treatment regimen? Yes No If yes, what
	is the cause of this problem?
	A. Duplication drug therapy
	B. No valid medical indication for the drug therapy at this time
11.	. Is there any ineffective drug therapy in the treatment regimen? Yes NoIf yes, what is
	the cause of this problem?
12.	. Is the dose too low? YesNO If yes, what is the cause of this problem?
	A. The dose is too low to produce the desired effect
	B. The dosing is too infrequent to produce the desired effect
13.	. Is the dose too high? YesNO
14.	. Is there any adverse drug reaction? Yes No If yes, what is the cause of this
	problem?
	A. Undesirable drug effect
	B. Unsafe or the patient/contraindication
15.	. Is there any non-compliance to their treatment regimen by the patient? Yes No If
	yes to this questions, what are the main reasons of non-compliance?
	A. Forgets to take medications
	B. Drug product too expensive for the patient
	C. Does not understand the instruction
	D. Cannot swallow or self-administer the drug product appropriately
	E. The drug product is not available for the patient