

Figure 1 Observed vs. simulated C_{\max} plot of oral TAF at various doses. The orange markers indicate 5 mg, black - 8 mg, blue - 10 mg and green - 25 mg. The triangles indicate tenofovir, diamonds – TFV-DP and circles – TAF plasma C_{\max} . Solid fill indicates single dose and dotted fill – multiple dose (day 8 or 10). The grey lines represent 0.8 and 1.25-fold range from the mean and the outer black lines represent 0.5 and 2-fold range




Figure 2 Observed vs. simulated AUC plot of oral TAF at various doses. The orange markers indicate 5 mg, black - 8 mg, blue - 10 mg, green - 25 mg and violet – 40 mg. The triangles indicate tenofovir, diamonds – TFV-DP and circles – TAF plasma AUC. Solid fill indicates single dose and dotted fill – multiple dose (day 8 or 10). The grey lines represent 0.8 and 1.25-fold range from the mean and the outer black lines represent 0.5 and 2-fold range

Table 1 PBPK qualification of TFV and TFV-DP at a single dose of 5 mg TAF [1]

Study	Clinical		Simulated		% difference	
	AUC _{last}	C _{max}	AUC _{last}	C _{max}	AUC _{last}	C _{max}
² TFV, plasma	52.8 (49.5-59.6)	0.8 (0.6-1)	59.8 (31-120)	1.55 (0.6-2.99)	13.3%	93.8%
TAF 5 mg day 1						
² TFV-DP, PBMC*	2268 (1519-4090)	33.7 (26.4-50.6)	2092 (1303-3981)	22.3 (14.2-42.6)	-7.8%	-33.8%

AUC is AUC_{0-14 days}. Values are represented as mean (CV). AUC is represented in ng.h/ml, C_{max} in ng/ml. *TFV-DP values are in fmol.h/10⁶ cells, C_{max} is in fmol/10⁶ cells and are represented as median (range).

Table 2 PBPK qualification of TAF and TFV on day 14 with once daily 8 mg TAF [2]

Study	Clinical		Simulated		% difference	
	AUC _{0-24h}	C _{max}	AUC _{0-24h}	C _{max}	AUC _{0-24h}	C _{max}
³ TAF, plasma	81.2 (43.9)	71.0 (72.9)	78.7 (19.3)	43.0 (21.9)	-3.1%	-39.4%
TAF 8 mg day 14						
³ TFV, plasma*	86.1 (19.4)	5.8 (19.5)	97.6 (37.5)	5.14 (32.9)	13.4%	-11.4%

*AUC is AUC_{0-24 h} on day 14. Values are represented as mean (CV). AUC is represented in ng.h/ml, C_{max} in ng/ml.

Table 3 PBPK qualification of TFV and TFV-DP at a single dose of 10 mg TAF [1]

Study	Clinical		Simulated		% difference	
	AUC _{last}	C _{max}	AUC _{last}	C _{max}	AUC _{last}	C _{max}
² TFV, plasma	78.1 (68.2-86.9)	1.5(1.4-1.7)	121 (61.5-239)	3.1 (1.8-5.9)	55.2%	106.7%
TAF 10 mg day 1						
² TFV-DP, PBMC*	4584 (3113-5734)	68 (56.9-88.8)	4121 (2598-7961)	44.4 (29.8-85.2)	-10.1%	-34.7%

AUC is AUC_{0-14 days}. Values are represented as mean (CV). AUC is represented in ng.h/ml, C_{max} in ng/ml. *TFV-DP values are in fmol.h/10⁶ cells, C_{max} is in fmol/10⁶ cells and are represented as median (range).

Table 4 PBPK qualification of TFV and TFV-DP at single and multiple doses of 25 mg TAF [1-3]

Study	Clinical		Simulated		% difference	
	AUC _{last}	C _{max}	AUC _{last}	C _{max}	AUC _{last}	C _{max}
TAF, plasma [3]	266 (49.5)	199 (62.1)	244 (17.9)	135 (21.5)	-8.3	-32.2
TAF 25 mg day 1	TFV, plasma [3]	298 (26.1)	9.50 (36.5)	287 (36.6)	7.54 (30.1)	-3.7
TFV-DP, PBMC* [1]		9306 (6891-10785)	113.6 (80.9-150.2)	9991 (6539-16590)	107.0 (66.8-191.6)	7.4 -5.8
TAF, plasma [2]	306 (18.4)	238 (25.5)	243 (19.7)	137 (20.7)	-20.6	-42.4
TAF 25 mg day 14	TFV plasma [2]	238 (14.0)	14.4 (12.9)	310 (40.3)	16.5 (36.4)	30.3
						14.6

Values are represented as mean (CV). AUC is represented in ng.h/ml, C_{max} in ng/ml. *AUC is AUC_{0-14 days} and are in fmol.h/10⁶ cells, C_{max} is in fmol/10⁶ cells. TFV-DP values are represented as median (range).

Table 5 Computation of TFV-DP cervical and rectal tissue ratios for TAF [1]**TAF single dose**

TFV-DP PBMCs		5 mg	10 mg	25 mg		5 mg	10 mg	25 mg
C _{max}	fmol/10 ⁶ cells	33.7	68	113.6	fmol/g	1.11E+05	2.23E+05	3.73E+05
AUC _{0-24 h}	fmol.h/10 ⁶ cells	657	1086	2094	fmol.h/g	2.16E+06	3.56E+06	6.87E+06
AUC _{0-14 days}	fmol.h/10 ⁶ cells	2268	4584	9306	fmol.h/g	7.44E+06	1.50E+07	3.05E+07
TFV-DP female genital tissue					TFV-DP female genital tissue/TFV-DP PBMC			
C _{max}	fmol/g	BLQ	4499	9552			0.020	0.026
AUC _{0-24 h}	fmol.h/g	BLQ	63703	104086	Average – 0.031		0.018	0.015
AUC _{0-14 days}	fmol.h/g	BLQ	986421	1309305			0.066	0.043
TFV-DP rectal tissue					TFV-DP rectal tissue/TFV-DP PBMC			
C _{max}	fmol/g	BLQ	3456	11163			0.015	0.030
AUC _{0-24 h}	fmol.h/g	BLQ	45103	39162	Average – 0.02		0.013	0.006
AUC _{0-14 days}	fmol.h/g	BLQ	315145	1041352			0.021	0.034

Unit conversion from fmol/ 10^6 cells to fmol/g

Volume of each PBMC = 282.9 fl [4]

Volume of each PBMC in litres = 2.829×10^{-13} L

Volume million PBMC in litres = 2.829×10^{-7} L

Volume million PBMC in mL = 2.829×10^{-4} mL

Density of PBMCs = 1.077 g/ml [5]

Weight of million PBMC in g = **3.05×10^{-4} g**

References

1. Cottrell, M.L., et al., *Single-dose pharmacokinetics of tenofovir alafenamide and its active metabolite in the mucosal tissues*. Journal of Antimicrobial Chemotherapy, 2017. **72**(6): p. 1731-1740.
2. Begley, R., et al., *Pharmacokinetics of Tenofovir Alafenamide When Coadministered With Other HIV Antiretrovirals*. JAIDS Journal of Acquired Immune Deficiency Syndromes, 2018. **78**(4): p. 465-472.
3. Custodio, J.M., et al., *Pharmacokinetics and Safety of Tenofovir Alafenamide in HIV-Uninfected Subjects with Severe Renal Impairment*. Antimicrobial Agents and Chemotherapy, 2016. **60**(9): p. 5135-5140.
4. Simiele, M., et al., *Evaluation of the Mean Corpuscular Volume of Peripheral Blood Mononuclear Cells of HIV Patients by a Coulter Counter To Determine Intracellular Drug Concentrations*. Antimicrobial Agents and Chemotherapy, 2011. **55**(6): p. 2976-2978.
5. Norouzi, N., H.C. Bhakta, and W.H. Grover, *Sorting cells by their density*. PLOS ONE, 2017. **12**(7): p. e0180520.