




SHAREABLE PDF

Patient-reported outcome measures after COVID-19: a prospective cohort study

Alyson W. Wong^{1,2,5}, Aditi S. Shah^{1,5}, James C. Johnston^{1,3,4,6},
Christopher Carlsten ^{1,3,6} and Christopher J. Ryerson^{1,2,6}

Affiliations: ¹Dept of Medicine, University of British Columbia, Vancouver, BC, Canada. ²Centre for Heart Lung Innovation, University of British Columbia, Vancouver, BC, Canada. ³UBC School of Population and Public Health, Vancouver, BC, Canada. ⁴BC Centre for Disease Control, Vancouver, BC, Canada. ⁵Contributed equally to this manuscript. ⁶Contributed equally to this manuscript.

Correspondence: Christopher J. Ryerson, St. Paul's Hospital, Ward 8B, 1081 Burrard St., Vancouver, Canada V6Z 1Y6. E-mail: chris.ryerson@hli.ubc.ca 1-604-806-8818

 @ERSpublications

Over 75% of patients admitted to hospital with COVID-19 have abnormal patient-reported outcome measures 3 months after symptom onset, with a third of patients reporting at least moderate impairment in major dimensions of quality of life <https://bit.ly/32QMMgw>

Cite this article as: Wong AW, Shah AS, Johnston JC, *et al.* Patient-reported outcome measures after COVID-19: a prospective cohort study. *Eur Respir J* 2020; 56: 2003276 [<https://doi.org/10.1183/13993003.03276-2020>].

This single-page version can be shared freely online.

To the Editor:

Patient reported outcome measures (PROMs) are standardised validated questionnaires completed by patients to provide information on their perceived functional well-being and health status [1]. These questionnaires can address various aspects of health including symptoms, quality of life, functionality, and physical, mental and social well-being. PROMs play an important role in increasing patient engagement, improving health systems, and ensuring that clinical care and research is person-centred.