Supplemental Table 1. Characteristics of enrolled patients in this study

| | | • |
|-----------------------|--------------|--------------|
| Parameters | MPT positive | MPT negative |
| Total (n) | 67 (23.5%) | 218 (76.4%) |
| Gender (n) | | |
| Male | 31 | 141 |
| Female | 36* | 77 |
| Age [mean (SD)] | 69.7 (10.9) | 69.7 (10.8) |
| Smoking status (n) | | |
| Former, Current | 27 | 164 |
| Never | 40* | 54 |
| Pack-year [mean (SD)] | 23.5 (39.6) | 25.8 (36.6) |
| HRCT pattern (n) | | |
| UIP | 18 | 75 |
| f-NSIP | 31 | 81 |
| c-NSIP | 9 | 15 |
| AIP | 6 | 8 |
| COP | 3 | 29 |
| PPFE | 0 | 4 |
| RB-ILD | 0 | 3 |
| HP | 0 | 3 |
| | | |

Data are presented as mean ± SD and number of patients as n. MPT, myositis panel test. HRCT classification are indicated usual interstitial pneumonia (UIP), fibrotic non-specific interstitial pneumonia (f-NSIP), cellular non-specific interstitial pneumonia (c-NSIP), acute interstitial pneumonia (AIP), cryptogenic organizing pneumonia (OP), pleuroparenchimal fibroelastosis (PPFE), respiratory bronchiolitis-associated interstitial lung disease (RB-ILD), and hypersensitivity pneumonia (HP). *p<0.05.