

Supplementary Table 1: Summary of Commonly Used Frailty Screening Tests

Test	Acronym	Range of Possible Scores	Frailty Indicator Threshold	Purpose
Eastern Collaborative Oncology Group Performance Status ^{1,2}	ECOG PS	0–4	≥1	Evaluation of cancer burden on functional status
Katz Activities of Daily Living ³	ADL	0–6	<5	Evaluation of functional independence
Mini-Cog ^{4,5}	Mini-Cog	0–5	≤2	Detection of cognitive impairment in older adults therefore suitable for a more thorough evaluation.
Flemish version of the Triage Risk Screening Test ^{6,7}	fTRST	0–6	≥2	Detection of hospitalized geriatric patients at risk for frailty
Timed Up & Go Test ^{8,9}	TUG	Not Applicable	≥20 s	10 feet (3 meters) walking test to determine gait speed
G8 ¹⁰	G8	0–17	≤14	Detection of onco-geriatric patients who may benefit from comprehensive geriatric assessment

Nutritional Risk Screening¹¹	NRS	Normal to severely impaired nutritional status	Moderately to severely impaired	Evaluation of nutritional status taking into account BMI, weight loss and food intake
American Society of Anesthesiology score¹²	ASA	1–5	Not applicable	Evaluation of preoperative general clinical condition and estimation of anesthesiologic risk
Charlson Age Comorbidity Index^{13,14}	CACI	0–42	≥6	Evaluation of cumulative burden of patient's comorbidities

References for Supplementary Table 1

1. Oken MM, Creech RH, Tormey DC, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol.* 1982;5:649–655.
2. Ethun CG, Bilen MA, Jani AB, Maithel SK, Ogan K, Master VA. Frailty and cancer: implications for oncology surgery, medical oncology, and radiation oncology. *CA Cancer J Clin.* 2017;67:362–377.
3. Katz S, Akpom CA. A measure of primary sociobiological functions. *Int J Health Serv.* 1976;6:493–508.
4. Ketelaars L, Pottel L, Lycke M, et al. Use of the Freund clock drawing test within the Mini-Cog as a screening tool for cognitive impairment in elderly patients with or without cancer. *J Geriatr Oncol.* 2013;4:174–182.
5. Robinson TN, Wu DS, Pointer LF, Dunn CL, Moss M. Preoperative cognitive dysfunction is related to adverse postoperative outcomes in the elderly. *J Am Coll Surg.* 2012;215:12–17.
6. Deschodt M, Wellens NI, Braes T, et al. Prediction of functional decline in older hospitalized patients: a comparative multicenter study of three screening tools. *Aging Clin Exp Res.* 2011;23:421–426.

7. Zattoni D, Montroni I, Saur NM, et al. A simple screening tool to predict outcomes in older adults undergoing emergency general surgery. *J Am Geriatr Soc.* 2019;67:309–316.
8. Huisman MG, van Leeuwen BL, Ugolini G, et al. “Timed Up & Go”: a screening tool for predicting 30-day morbidity in onco-geriatric surgical patients? A multicenter cohort study. *PLoS One.* 2014;9:e86863.
9. Podsiadlo D, Richardson S. The timed “Up & Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc.* 1991;39:142–148.
10. Bellera CA, Rainfray M, Mathoulin-Pélissier S, et al. Screening older cancer patients: first evaluation of the G-8 geriatric screening tool. *Ann Oncol.* 2012;23:2166–2172.
11. Kondrup J, Rasmussen HH, Hamberg O, Stanga Z; Ad Hoc ESPEN Working Group. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr.* 2003;22:321–336.
12. Owens WD, Felts JA, Spitznagel EL Jr. ASA physical status classifications: a study of consistency of ratings. *Anesthesiology.* 1978;49:239–243.
13. Chang CM, Yin WY, Wei CK, et al. Adjusted age-adjusted Charlson comorbidity index score as a risk measure of perioperative mortality before cancer surgery. *PLoS One.* 2016;11:e0148076.
14. Tian Y, Jian Z, Xu B, Liu H. Age-adjusted Charlson comorbidity index score as predictor of survival of patients with digestive system cancer who have undergone surgical resection. *Oncotarget.* 2017;8:79453–79461.