**Total patients Survivors** Non survivors p-N = 79N = 51N = 28value Sex Male.% 50 (63%) 36 (71%) 14 (50%) 0.069 29 (37%) Female,% 15 (29%) 14 (50%) Age, median [IQR] 61 [51-67] 61 [48-67] 63.5 [56-71.5] 0.073 Race Hispanic 43 (54%) 27 (53%) 16 (57%) 0.26 African American 27 (34%) 20 (39%) 7 (25%) Other 9 (11%) 4 (8%) 5 (18%) **Type of transplant** Deceased donor 61 (77%) 23 (82%) 38 (75%) 0.44 Living donor 18 (23%) 13 (26%) 5 (18%) **Time after** 58.2 [25.4-57.7 [27.3-65.2 [16.3transplantation, median 0.22 127.6] 123.7] 134.1] [IQR] months **Transplant** < 6 months 6 (8%) 5 (10%) 1 (4%) 0.32 **Transplant** < 12 months 9 (11%) 8 (16%) 1 (4%) 0.22 **Immunosuppressive** regimen Calcineurin inhibitors 74 (71/3) 49 (48/1) 25 (23/2) 0.34 (Tacrolimus/Cylosporin) (94%) (96%) (89%) MMF/MPA 45 (88%) 0.89 70 (89%) (89%) 0.12 Corticosteroid 75(95%) 50 (98%) 25 (89%) Sirolimus 7 (3%) 5 (4%) 2 (3%) 0.28 **Etiology of ESRD** Diabetes mellitus 42 (53%) 25 (49%) 17 (61%) Hypertension 13 (17%) 10 (20%) 3 (11%) Glomerulonephritis 15 (19%) 9 (18%) 6 (21%) 0.66 Polycystic kidney disease 3 (4%) 2 (4%) 1 (4%) Others 6 (8%) 5 (10%) 1 (4%)

Supplemental Table 1: Clinical characteristics of the patients hospitalized at Montefiore Medical Center

Body Mass Index (kg/m2) median [IQR]	28.1 [23.7-32.3]	26.7 [23.1- 31.4]	30.4 [25.7- 34.7]	0.057
History of smoking	28 (35%)	18 (35%)	10 (36%)	0.97
Influenza vaccination	59 (83%)	43 (88%)	16 (73%)	0.12
Comorbidities				
Hypertension	77 (98%)	49 (96%)	28 (100%)	0.54
Diabetes mellitus	53 (67%)	32 (63%)	21 (75%)	0.27
Heart disease	15 (19%)	10 (20%)	5 (18%)	0.85
Lung disease	9 (11%)	4 (8%)	5 (18%)	0.18
Cancer	9 (11%)	5 (10%)	4 (14%)	0.55
Blood type				
А	28 (36%)	15 (30%)	13 (46%)	
В	16 (21%)	13 (26%)	3 (11%)	0.19
AB	3 (4%)	1 (2%)	2 (7%)	
0	31 (40%)	21 (42%)	10 (36%)	
Angiotensin converting				
enzyme inhibitor or	10 (240/)	10 (0.40/)	7 (050()	0.00
angiotensin receptor	19 (24%)	12 (24%)	7 (25%)	0.88
blocker prior to admission				
Statin prior to admission	49 (62%)	31 (61%)	18 (64%)	0.76
Baseline creatinine, median [IQR]	1.4 [1.2-2.0]	1.4 [1.1-2.0]	1.5 [1.2-2.0]	0.86
Symptoms on admission				
Fever	46 (58%)	31 (61%)	15 (54%)	0.53
Dyspnea	46 (58%)	26 (51%)	20 (71%)	0.078
Myalgias	32 (41%)	17 (33%)	15 (54%)	0.08
Cough	46 (58%)	31 (61%)	15 (54%)	0.53
Diarrhea	28 (35%)	18 (35%)	10 (36%)	0.97
Pulse oximetry on admission (%)	95 [92-98]	96 [94-98]	93 [89-96.5]	0.037
Temperature on admission (F)	99 [98.3-100.5]	99 [98.4- 100.1]	99.1 [98.3- 100.9]	0.87