

Manuscript No.: microorganisms-889985

Regulation of Chitin Dependent Growth and Natural Competence in *Vibrio parahaemolyticus*

ANUSUYA DEBNATH^{1*}, TAMAKI MIZUNO¹ and SHIN-ICHI MIYOSHI¹

¹Okayama University, Okayama, Japan

***Corresponding author**

Anusuya Debnath

Graduate School of Medicine, Dentistry and Pharmaceutical Sciences

Okayama University

1-1-1, Tsushimanaka, Kita-ku

Okayama 700-8530

Phone: +81-86-251-7959

e-mail: anusuyadebnath@yahoo.co.in, pu0139jq@okayama-u.ac.jp

Supplementary Figures




Figure S1. The relative expression of competence genes *tfoX*, *pilA*, *comEA* and *qstR* at mRNA level was checked for WT, ΔchiS and ΔtfoS mutant in the presence of chitin. *pvsA* was used as an internal control. Each bar indicates the mean \pm SE of three independent experiments.




Figure S2. The relative mRNA expression of competence genes *tfoX*, *pilA*, *comEA* and *qstR* was compared in the presence of GlcNAc₂ and GlcNAc₆ in wild type strain. *pvsA* was used as an internal control. Each bar indicates the mean \pm SE of three independent experiments.