

Supplementary Material

Isolation and FTIR-ATR and ^1H NMR Characterization of Alginates from the Main Alginophyte Species of the Atlantic Coast of Morocco

Zahira Belattmania ¹, Soukaina Kaidi ¹, Samir El Atouani ¹, Chaimaa Katif ¹, Fouad Bentiss ^{2,3}, Charafeddine Jama ³, Abdeltif Reani ¹, Brahim Sabour ¹ and Vitor Vasconcelos ^{4,5,*}

¹ R.U. Phycology, Blue Biodiversity & Biotechnology – P3B, Laboratory of Plant Biotechnology, Ecology and Ecosystem Valorization, Faculty of Sciences, Chouaib Doukkali University, PO Box 20, M-24000 El Jadida; belattmania.z@ucdac.ma (Z.B.); souk.kaidi@gmail.com (S.K.); elatouanisamir@gmail.com (S.E.); chaimaakatif25@gmail.com (C.K.); abreani@yahoo.fr (A.R.); sabour.b@ucd.ac.ma (B.S.)

² Laboratory of Catalysis and Corrosion of Materials, Faculty of Sciences, Chouaib Doukkali University, PO Box 20, M-24000 El Jadida, Morocco; fbentiss@gmail.com (F.B.)

³ University of Lille, CNRS, INRAE, Centrale Lille, UMR 8207, UMET – Unité Matériaux et Transformations, F-59000 Lille, France; charafeddine.jama@ensc-lille.fr (C.J.)

⁴ CIIMAR, Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Terminal de Cruzeiros do Porto de Leixões, Av. General Norton de Matos, s/n, 4450-208 Matosinhos, Portugal

⁵ Department of Biology, Faculty of Sciences, University of Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal

* Correspondence: vmvascon@fc.up.pt; Tel.: 351-223401817 (V.V.)




Figure S1. The anomeric region in the 400 MHz - ^1H NMR spectrum of the Sigma-Aldrich sodium alginate using D_2O as solvent.



Figure 2S. Morphological traits of the studied macroalgae collected on the Northwestern Atlantic coast of Morocco.