

Supplementary Data

SUPPLEMENTARY TABLE S1. THYROID FINE NEEDLE ASPIRATE IMMUNOPHENOTYPING DATA COMPARING IMMUNE CHECKPOINT INHIBITOR THYROIDITIS SAMPLES COLLECTED WITHIN TWO WEEKS OF DIAGNOSIS AND HEALTHY THYROID SAMPLES

<i>Immune cell profile (% of WBCs)</i>	<i>ICI-induced thyroiditis at diagnosis (n=8)</i>	<i>Healthy volunteers (n=5)</i>	<i>Difference of means</i>	<i>p-value (t-test)</i>	<i>ICI-induced thyroiditis 2 months later (n=1)</i>
WBCs (% of all cells)	5.90	0.52	5.34	0.285	1.54
Granulocytes	6.22	15.73	-9.51	0.004	44.65
Mononuclear cells	93.84	84.32	9.518	0.004	15.55
Lymphocytes	81.23	74.50	6.729	0.192	14.05
CD3 ⁺ T lymphocytes	61.30	20.10	41.20	0.00006	4.27
CD4 ⁺ T lymphocytes	34.52	13.19	21.33	0.005	1.26
CD8 ⁺ T lymphocytes	23.21	5.07	18.14	0.00006	1.29
CD4 ⁻ CD8 ⁻ T lymphocytes	1.89	0.72	1.174	0.006	0.11
CD19 ⁺ B lymphocytes	9.59	26.72	-17.13	0.007	1.33
CD16 ⁺ CD56 ⁺ NK cells	0.48	0.31	0.17	0.760	0
CD16 ⁻ CD56 ⁺ NK cells	0.25	0.23	0.02	0.932	0.01
CD16 ⁺ CD56 ⁻ cells	0.31	0.43	-0.12	0.589	0.01
Monocytes	2.62	5.09	-2.47	0.097	0.81
Classical monocytes	1.63	2.89	-1.26	0.288	0.46
Intermediate monocytes	0.49	1.32	-0.83	0.068	0.23
Nonclassical monocytes	0.24	0.82	-0.58	0.060	0.05

ICI, immune checkpoint inhibitor; WBC, white blood cell.