

SUPPLEMENTARY MATERIAL

Nanofibrous asymmetric collagen/curcumin membrane containing aspirin-loaded PLGA nanoparticles for guided bone regeneration

**Mohammad Ali Ghavimi ^{a,#}, Amirhossein Bani Shahabadi ^{a,#}, Seyedhosein Jarolmasjed ^b,
Mohammad Yousef Memar ^c, Solmaz Maleki Dizaj ^{d,*}, Simin Sharifi ^{d,e,*}**

^a *Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran*

^b *Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran*

^c *Infectious and Tropical Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran*

^d *Dental and Periodontal Research Center, Tabriz University of Medical Sciences, Tabriz, Iran*

^e *Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran*

Co-First/Equal authorship

*maleki.s.89@gmail.com

P* sharifi.ghazi@gmail.com






Figure 1. Osteogenic differentiation of DPSCs cultured on PACNs and CCNFs membranes at 21 day in the osteogenic medium. The cell lysates were prepared and used for western blot with GAPDH, Runx-2 and OCN antibodies. The figure shows (a) GAPDH (37 kDa), (b) Runx-2 (55 kDa) (c) OCN (6 kDa). The gels were not cropped.