

Supporting Information

**Antibacterial Activity of Chromomycins from a Marine-Derived
*Streptomyces microflavus***

**Eunji Cho ¹, Oh-Seok Kwon ², Beomkoo Chung ¹, Jayho Lee ¹, Jeongyoon Sun ¹, Jongheon Shin ^{2,*}
and Ki-Bong Oh ^{1,*}**

¹ *Department of Agricultural Biotechnology, College of Agriculture and Life Sciences, Seoul National University, Seoul 08826, Korea*

² *Natural Products Research Institute, College of Pharmacy, Seoul National University, Seoul 08826, Korea*

Contents

Figure S1. The ^1H NMR (400 MHz, chloroform-*d*) spectrum of **1**

Figure S2. The ^{13}C NMR (100 MHz, chloroform-*d*) spectrum of **1**

Figure S3. The COSY (400 MHz, CDCl_3) spectrum of **1**

Figure S4. The TOCSY (400 MHz, CDCl_3) spectrum of **1**

Figure S5. The eHSQC (400 MHz, chloroform-*d*) spectrum of **1**

Figure S6. The HMBC (400 MHz, chloroform-*d*) spectrum of **1**




Figure S1. The ^1H NMR (400 MHz, chloroform-*d*) spectrum of **1**




Figure S2. The ^{13}C NMR (100 MHz, chloroform-*d*) spectrum of 1




Figure S3. The COSY (400 MHz, CDCl₃) spectrum of **1**




Figure S4. The TOCSY (400 MHz, CDCl_3) spectrum of **1**




Figure S5. The eHSQC (400 MHz, chloroform-*d*) spectrum of 1




Figure S6. The HMBC (400 MHz, chloroform-*d*) spectrum of 1