

**Supplementary Table S1. Relationship between nitrogen balance and outcomes according to initial diagnosis**

	<b>Positive nitrogen balance, n (%)</b>	<b>Negative nitrogen balance, n (%)</b>	<b>P-value</b>
<b>IS (n = 26, 14.9%)</b>	1 (3.8)	25 (96.2)	
<b>Good outcome at 3 months (mRS = 0-3), n (%)</b>	1 (100)	10 (40.0)	0.423
<b>Neurological worsening, n (%)</b>	0 (0.0)	9 (36.0)	1.00
<b>In hospital mortality, n (%)</b>	0 (0.0)	8 (32.0)	1.000
<b>SAH (n = 30, 17.1%)</b>	9 (30.0)	21 (70.0)	
<b>Good outcome at 3 months (mRS = 0-3), n (%)</b>	6 (66.7)	11 (52.4)	0.691
<b>Neurological worsening, n (%)</b>	1 (11.1)	3 (14.3)	1.000
<b>In hospital mortality, n (%)</b>	1 (11.1)	3 (14.3)	1.000
<b>ICH (n = 33, 18.9%)</b>	2 (8.3)	22 (91.7)	
<b>Good outcome at 3 months (mRS = 0-3), n (%)</b>	2 (100.0)	7 (31.8)	0.130
<b>Neurological worsening, n (%)</b>	0 (0.0)	3 (13.6)	1.000
<b>In hospital mortality, n (%)</b>	0 (0.0)	3 (13.6)	1.000

IS: ischemic stroke, mRS: modified Rankin Scale, SAH: subarachnoid hemorrhage, ICH: intracerebral hemorrhage