

Supplementary Table 1. Logistic regression for any chemotherapy use, all factors included

Variable	OR	95% CI low	95% CI high	p-value
Mean age (<50 ref.) (y)				
51–60	0.825	0.617	1.103	0.194
61–70	0.682	0.513	0.906	0.008
>70	0.283	0.211	0.378	<0.001
Male (sex)	1.087	0.972	1.217	0.145
Race (white ref.)				
Black	1.035	0.835	1.283	0.755
Other	0.976	0.727	1.308	0.869
CCI (0 ref.)				
1	0.852	0.758	0.958	0.008
2	0.644	0.526	0.787	<0.001
3+	0.592	0.414	0.845	0.004
cT Stage (cT2 ref.)				
3	1.440	1.231	1.686	<0.001
4	1.487	1.260	1.754	<0.001
Income status (<\$38,000 ref.)				
\$38,000–47,999	1.077	0.918	1.263	0.365
\$48,000–62,999	1.217	1.039	1.425	0.015
≥\$63,000	1.352	1.154	1.583	<0.001
Low volume facility	1.087	0.945	1.250	0.242
Geographic location (Eastern ref.)				
Central	0.903	0.810	1.006	0.065
Western	0.904	0.783	1.043	0.167
Insurance status (uninsured ref.)				
Private	1.566	1.105	2.220	0.012
Medicaid	1.328	0.889	1.984	0.166
Medicare	1.346	0.945	1.917	0.100
Other govt or unknown	2.380	1.486	3.813	<0.001

OR, odds ratio; CI, confidence interval; Ref., reference; CCI, Charlson Comorbidity Index; Govt, government.