

Supplemental Table 1.

Patient and Clinical Variables	Incontinent Cohorts (N = 159)	SUI Cohort (N = 50)	UII Cohort (N=109)	<i>p</i> -value
Age (years), mean (SD)	59 (13)	54 (14)	61 (13)	0.001^a
Race/Ethnicity				0.07 ^b
-White	112 (70%)	33 (66%)	79 (72%)	
-Hispanic	20 (13%)	8 (16%)	12 (11%)	
-Black	20 (13%)	3 (6%)	17 (16%)	
-Asian	1 (1%)	1 (2%)	0 (0%)	
-Other	6 (4%)	5 (10%)	1 (1%)	
BMI (kg/m ²), mean (SD)	31.2 (8.3)	30.2 (6.9)	31.7 (8.8)	0.16 ^a
Sexually Active	53 (53%) (N=100)	27 (54%) (N=50)	26 (52%) (N=50)	0.84
Current Anticholinergic Treatment	10 (6%)	2 (4%)	8 (7%)	0.42
Use of Hormone Replacement Therapy	19 (12%)	4 (8%)	15 (14%)	0.30
Diabetes	21 (13%)	9 (18%)	13 (12%)	0.30
Heart Disease	17 (11%)	3 (6%)	14 (13%)	0.19
Hypertension	66 (42%)	18 (36%)	48 (44%)	0.34
Urine Dipstick:				
- White Blood Cells (WBC)	2 (1%)	0 (0%)	2 (2%)	0.74
- Nitrites	0 (0%)	0 (0%)	0 (0%)	1
- Red Blood Cells (RBC)	23 (14%)	6 (12%)	17 (16%)	0.55
Urinary Distress Inventory (UDI) Score, mean (SD)	95.04 (51.03)	97.95 (55.36)	93.71 (49.12)	0.31 ^a
EQUC-Positive Specimen	131 (82%)	43 (86%)	88 (81%)	0.42

Chi-square test used unless otherwise indicated. SD = standard deviation.

a – two sample t test, b – Fisher’s exact test

Bold – significant p value