

1 **Supplementary References**

- 2 S1 Waldherr R, Rambausek M, Duncker WD, et al. Frequency of mesangial IgA deposits in a
3 non-selected autopsy series. *Nephrol Dial Transplant.* 1989;4:943–946.
- 4 S2 Suzuki H, Kiryluk K, Novak J, et al. The pathophysiology of IgA nephropathy. *J Am Soc
5 Nephrol.* 2011;22:1795–1803.
- 6 S3 Gharavi AG, Moldoveanu Z, Wyatt RJ, et al. Aberrant IgA1 glycosylation is inherited in
7 familial and sporadic IgA nephropathy. *J Am Soc Nephrol.* 2008;19:1008–1014.
- 8 S4 Suzuki H, Fan R, Zhang Z, et al. Aberrantly glycosylated IgA1 in IgA nephropathy patients
9 is recognized by IgG antibodies with restricted heterogeneity. *J Clin Invest.* 2009;119:1668–
10 1677.
- 11 S5 Oka K, Nishimura K, Kishikawa H, et al. IgA1 dominant subclass of latent IgA mesangial
12 deposition in donated kidney. *Int J Nephrol Renovasc Dis.* 2016;9:313–317.
- 13 S6 Rizk D V., Saha MK, Hall S, et al. Glomerular immunodeposits of patients with IgA
14 nephropathy are enriched for IgG autoantibodies specific for galactose-deficient IgA1. *J Am
15 Soc Nephrol.* 2019;30:2017–2026.
- 16 S7 Sofue T, Inui M, Hara T, et al. Association between post-transplantation immunoglobulin a
17 deposition and reduced allograft function. *Transplant Proc.* 2015;47:332–336.
- 18 S8 Abrera-Abeleda MA, Xu Y, Pickering MC, et al. Mesangial immune complex
19 glomerulonephritis due to complement factor D deficiency. *Kidney Int.* 2007;71:1142–1147.

1 S9 Stuht S, Gwinner W, Franz I, et al. Lymphatic neoangiogenesis in human renal allografts:
2 results from sequential protocol biopsies. *Am J Transplant.* 2007;7:377–384.