

Supplementary Table 1. Differences in HP CNV and rs2000999 genotype CROMIS versus ALSPAC, genotype frequency

| | Hp1-1 | Hp2-1 | Hp2-2 | Total | p-value overall |
|--------|---------------------|---------------------|---------------------|--------------|------------------------|
| CROMIS | 151 17.9% | 413 49% | 279 33.1% | 843 100% | 0.01 |
| ALSPAC | 137 14.8% | 418 45.1% | 372 40.1% | 927 100% | |
| | rs2000999 AA | rs2000999 AG | rs2000999 GG | | |
| CROMIS | 29 3.5% | 267 32.6% | 523 63.9% | 843 100% | 0.47 |
| ALSPAC | 34 4.6% | 229 30.6% | 485 64.8% | 748 100% | |

Supplementary Table 2. Univariable analysis for functional outcome and death

| Variable | Unfavourable outcome | | | Death | | |
|------------------------------|-----------------------------|---------------|------------------|--------------|---------------|------------------|
| | OR | 95% CI | p-value | OR | 95% CI | p-value |
| Age, years | 1.05 | 1.03-1.06 | <0.001 | 1.1 | 1.07-1.13 | <0.001 |
| Female Sex | 2.3 | 1.69-3.14 | <0.001 | 1.69 | 1.13-2.53 | 0.01 |
| Ethnicity | | | 0.71 | | | |
| - White | Reference | | | Reference | | 0.89 |
| - Asian | 1.3 | 0.66-2.58 | | 0.74 | 0.22-2.53 | |
| - Black | 1.16 | 0.5-2.66 | | 0.99 | 0.4-2.43 | |
| Smoker | 0.83 | 0.61-1.12 | 0.21 | 1.2 | 0.79-1.8 | 0.39 |
| Hypertension | 1.79 | 1.31-2.45 | <0.001 | 1.23 | 0.79-1.9 | 0.36 |
| SBP on admission, mmHg | 1 | 0.99-1.002 | 0.33 | 0.99 | 0.99-1 | 0.15 |
| DM | 1.48 | 0.98-2.26 | 0.06 | 1.5 | 0.91-2.48 | 0.11 |
| Anti-hypertensive medication | 1.45 | 1.07-1.97 | 0.02 | 1.76 | 1.17-2.65 | 0.01 |
| Anticoagulation | 1.41 | 1.03-1.94 | 0.03 | 1.78 | 1.18-2.68 | 0.006 |
| Antiplatelets | 1.33 | 0.93-1.89 | 0.11 | 1.04 | 0.65-1.66 | 0.86 |
| Hypercholesterolaemia | 1.57 | 1.15-2.14 | 0.004 | 1.09 | 0.72-1.66 | 0.67 |
| GCS admission | | | | | | |
| - GCS 3-8 | 4.41 | 1.52-12.78 | 0.006 | 5.51 | 2.6-11.64 | <0.001 |
| - GCS 9-15 | Reference | | | Reference | | |

| | | | | | | |
|-------------------|-----------|-----------|------------------|-----------|-----------|------------------|
| ICH location | | | 0.63 | | | |
| - Cerebellar | Reference | | | Reference | | 0.1 |
| - Brainstem | 1.35 | 0.4-4.55 | | 0.34 | 0.04-3.08 | |
| - Deep | 0.77 | 0.39-1.52 | | 0.82 | 0.33-2.07 | |
| - Lobar | 0.84 | 0.42-1.68 | | 1.36 | 0.54-3.39 | |
| crICH volume, mL | 1.61 | 1.35-1.93 | <0.001 | 1.88 | 1.52-2.34 | <0.001 |
| IV extension | 2.38 | 1.64-3.45 | <0.001 | 2.62 | 1.72-4 | <0.001 |
| crPHO volume, mL | 1.39 | 1.17-1.66 | <0.001 | 1.82 | 1.44-2.29 | <0.001 |
| OED, cm | 1.32 | 0.68-2.54 | 0.41 | 2.76 | 1.23-6.17 | 0.01 |
| Neurosurgery | 2.99 | 1-8.94 | 0.05 | 1.23 | 0.41-3.72 | 0.71 |
| HP CNV | | | 0.63 | | | 0.15 |
| - 1-1 | Reference | | | Reference | | |
| - 2-1 | 0.82 | 0.54-1.24 | | 0.65 | 0.37-1.12 | |
| - 2-2 | 0.84 | 0.54-1.3 | | 0.96 | 0.55-1.65 | |
| HP CNV 1 vs all 2 | 0.83 | 0.56-1.22 | 0.34 | 1.29 | 0.85-1.95 | 0.23 |
| HP CNV all 1 vs 2 | 0.93 | 0.71-1.33 | 0.86 | 0.77 | 0.47-1.27 | 0.31 |
| HP SNP | | | 0.95 | | | 0.49 |
| - A:A | Reference | | | Reference | | |
| - A:G | 0.88 | 0.38-2 | | 0.91 | 0.32-2.54 | |
| - G:G | 0.9 | 0.4-2.02 | | 0.71 | 0.26-1.96 | |

Cm = centimetre; CNV = copy number variation; crICH = cube root intracerebral haemorrhage; crPHO = cube root perihematomal oedema; DM = diabetes mellitus; GCS = Glasgow Coma Scale; HP = Haptoglobin; ICH = intracerebral haemorrhage; IV = intraventricular; ml = millilitre; mmHg = millimetre of mercury; OED = oedema extension distance; OR = Odds Ratio; SNP = single nucleotide polymorphism; y = year

Supplementary Table 3. Factors associated with 6 months mortality after ICH in an adjusted multivariable logistic regression model, HP CNV dichotomized

| | OR | 95% CI | P value |
|---------------------------|------|------------|---------|
| Age (years) | 1.11 | 1.07-1.14 | <0.001 |
| Female Sex | 1.15 | 0.69-1.94 | 0.6 |
| Hypertension | 0.98 | 0.56-1.72 | 0.95 |
| Diabetes mellitus | 1.33 | 0.65-2.69 | 0.43 |
| Oral anticoagulation | 1.27 | 0.75-2.13 | 0.37 |
| GCS on admission (binary) | 4.37 | 1.39-13.73 | 0.01 |
| - GCS 3-8 | | | |
| - GCS 9-15 (reference) | | | |
| ICH location | | | |

| | | | | |
|---|-----------------|-----------|--|--------|
| - Cerebellar (reference) | | | | 0.39 |
| - Brainstem | No observations | | | |
| - Deep | 0.94 | 0.32-2.8 | | |
| - Lobar | 0.62 | 0.2-1.93 | | |
| Cr ICH volume (mL) | 1.99 | 1.45-2.74 | | <0.001 |
| OED (cm) | 3 | 1.07-8.39 | | 0.04 |
| IV extension | 1.57 | 0.9-2.75 | | 0.11 |
| <i>HP CNV</i> | | | | |
| - HP1 allele (<i>HP1-1/2-1</i> vs <i>HP2-2</i>) | 0.55 | 0.31-0.95 | | 0.03 |
| Rs2000999 | | | | 0.77 |
| - A:A (reference) | | | | |
| - A:G | 0.61 | 0.16-2.4 | | |
| - G:G | 0.61 | 0.16-2.41 | | |

Cm = centimetre; CNV = copy number variation, Cr = cube root; CT = computed tomography; GCS = Glasgow Coma Scale; *HP* = Haptoglobin; ICH = intracerebral haemorrhage; IV = intraventricular; OAC = oral anticoagulation; SBP = systolic blood pressure

Supplementary Table 4. Univariable analysis for ICH and OED volume

| Variable | ICH volume | | | OED volume | | |
|---|-------------|---------------|------------------|-------------|----------------|------------------|
| | Coefficient | 95% CI | p-value | Coefficient | 95% CI | p-value |
| Time event to CT | | | 0.5 | | | 0.13 |
| - Day 1 | Reference | | | | | |
| - Day 2 | 0.18 | -0.12-0.48 | | 0.07 | 0.001-0.15 | |
| - Day 3 | -0.17 | -0.61-0.27 | | 0.03 | -0.09-0.15 | |
| Age, years | 0.004 | -0.002-0.01 | 0.2 | 0.001 | -0.001-0.002 | 0.25 |
| Female Sex | 0.05 | -0.1-0.2 | 0.52 | 0.02 | -0.01-0.06 | 0.19 |
| Smoker | -0.03 | -0.11-0.04 | 0.39 | -0.008 | -0.03-0.01 | 0.4 |
| Hypertension | 0.06 | -0.1-0.22 | 0.45 | -0.005 | -0.04-0.03 | 0.77 |
| SBP on admission, mmHg | -0.002 | -0.004-0.0005 | 0.12 | 0.0003 | -0.0005-0.0007 | 0.66 |
| Anti-hypertensive medication | 0.08 | -0.07-0.23 | 0.29 | -0.07 | -0.1-0.03 | <0.001 |
| Anticoagulation | 0.09 | -0.06-0.25 | 0.25 | -0.05 | -0.09-(-0.01) | 0.01 |
| Antiplatelets | 0.01 | -0.16-0.18 | 0.89 | -0.03 | -0.07-0.02 | 0.21 |
| High cholesterol | 0.16 | 0.01-0.32 | 0.03 | -0.006 | -0.04-0.03 | 0.74 |
| Platelet level, x10 ⁹ /liter | 0.001 | -0.0002-0.002 | 0.13 | 0.0003 | 0.0001-0.001 | 0.002 |
| ICH location | | | <0.001 | | | <0.001 |

| | | | | | | |
|----------------------------|-----------|---------------|-------------|--------|------------|-------------|
| - Cerebellar | Reference | | | | | |
| - Brainstem | -0.86 | -1.34-(-0.37) | | -0.09 | -0.21-0.04 | |
| - Deep | -0.19 | -0.49-0.11 | | 0.15 | 0.07-0.23 | |
| - Lobar | 0.67 | 0.37-0.97 | | 0.23 | 0.15-0.31 | |
| IV extension | 0.46 | 0.29-0.62 | <0.001 | -0.03 | -0.07-0.01 | 0.16 |
| Neurosurgery | 0.62 | 0.15-1.09 | 0.01 | -0.003 | -0.12-0.12 | 0.95 |
| <i>HP</i> CNV | | | 0.35 | | | 0.84 |
| - 1-1 | Reference | | | | | |
| - 2-1 | 0.11 | -0.09-0.34 | | 0.004 | -0.05-0.06 | |
| - 2-2 | 0.02 | -0.2-0.24 | | -0.01 | -0.06-0.05 | |
| <i>HP</i> CNV all 1 vs 2-2 | -0.07 | -0.22-0.09 | 0.39 | -0.01 | -0.05-0.03 | 0.57 |
| <i>HP</i> CNV all 2 vs 1-1 | 0.08 | -0.12-0.28 | 0.42 | -0.001 | -0.05-0.05 | 0.96 |
| <i>HP</i> SNP | | | 0.82 | | | 0.98 |
| - A:A | Reference | | | | | |
| - A:G | 0.03 | -0.39-0.45 | | -0.002 | -0.1-0.1 | |
| - G:G | 0.08 | -0.33-0.49 | | -0.005 | -0.11-0.1 | |

CNV = copy number variation; *HP* = Haptoglobin; ICH = intracerebral haemorrhage; IV = intraventricular; OED = oedema extension distance; SBP = systolic blood pressure; SNP = single nucleotide polymorphism