

Supplementary material 10. Baseline demographic and clinical characteristics for the qualitative participants.

	N=25
Age (years)	69±10
Gender n (%)	Female 13 (52%)
	Male 12 (48%)
Ethnicity n (%)	White background 13 (52%)
	Asian or Asian British 10 (40%)
	Caribbean 1 (4%)
	Not stated 1 (4%)
Diagnosis	Diabetic nephropathy 11 (44%)
	Aetiology uncertain 6 (24%)
	Chronic pyelonephritis 3 (12%)
	Atypical hemolytic uremic syndrome 1 (4%)
	FSGS 1 (4%)
	Henoch-Schönlein Purpura 1 (4%)
	Minimal change nephropathy 1 (4%)
	Polycystic kidney disease 1 (4%)
CCI	6±2
Time on HD (months)	43 (IQR 16-85)
CFS n (%)	Vulnerable 9 (36%)
	Mildly frail 5 (20%)
	Moderately frail 8 (32%)
	Severely frail 3 (12%)
Number of falls in the last six months	3 (IQR 2-4)
Previous transplant n (%)	No 21 (84%)
	Yes 4 (16%)
Active on transplant list n (%)	No 22 (88%)
	Yes 3 (12%)

Abbreviations: CCI, Charlson comorbidity index; CFS, clinical frailty scale; FSGS, Focal segmental glomerulosclerosis; HD, haemodialysis.