

Supplementary Table 6a. Complications: MIS cases.

	Pre-ERAS	ERAS (early)	ERAS (late)	<i>p</i>
n	623	701	697	
Grade				
No complication	562 (90.2)	622 (88.7)	630 (90.4)	
1	17 (2.7)	27 (3.9)	19 (2.7)	
2	28 (4.5)	27 (3.9)	24 (3.4)	
3	16 (2.6)	20 (2.9)	23 (3.3)	
4	0 (0)	4 (0.6)	1 (0.1)	
5	0 (0)	1 (0.1)	0 (0)	
Category				
Cardiovascular system	2 (0.3)	5 (0.7)	1 (0.1)	0.22
Endocrine system	0 (0)	1 (0.1)	0 (0)	
Gastrointestinal system	24 (3.9)	32 (4.6)	35 (5.0)	
General	2 (0.3)	3 (0.4)	5 (0.7)	
Genitourinary system	7 (1.1)	2 (0.3)	4 (0.6)	
Hematologic or vascular system	6 (1.0)	10 (1.4)	1 (0.1)	
Infection	10 (1.6)	17 (2.4)	10 (1.4)	
Metabolic	0 (0)	1 (0.1)	0 (0)	
Musculoskeletal system	0 (0)	2 (0.3)	2 (0.3)	
Nervous system	2 (0.3)	0 (0)	0 (0)	
Pulmonary system	1 (0.2)	1 (0.1)	1 (0.1)	
Wound or skin	7 (1.1)	5 (0.7)	8 (1.1)	

ERAS, Enhanced Recovery after surgery. *p* = chi-square test.

Supplementary Table 6b. Surgical site infections: MIS cases, from SSI surveillance cases reported to National Healthcare Safety Network.

	Pre-ERAS	ERAS (early)	ERAS (late)	<i>p</i>
n reported (%)	361 (57.9)	611 (87.2)	596 (85.5)	
SSI type				
SIP (%)	3 (0.8)	5 (0.8)	1 (0.2)	0.25
DIP (%)	1 (0.3)	0 (0)	0 (0)	0.19
IAB (%)	5 (1.4)	14 (2.3)	15 (2.5)	0.49

ERAS, Enhanced Recovery after surgery. SSI, surgical site infection. SIP, superficial incisional primary.

DIP, deep incisional primary. IAB, intraabdominal infection. *p* = chi-square test.