

## Supplementary table

### Factors that Influence the Reintegration to Normal Living for Older Adults 2 Years Post-Hip Fracture

Supplement 1: Characteristics of caregivers of older adults at 18 and 24 months post-discharge.

<b>Variables</b>	<b>n (%) / mean <math>\pm</math> SD (n=76)</b>
<b>Socio-demographic characteristics</b>	
Age [n(%)]	
64-74 years	55 (73)
75-84 years	15 (20)
85 years and older	5 (7)
Age (mean $\pm$ SD)	65.4 $\pm$ 13.1
Gender, female [n(%)]	57 (75)
Employment [n(%)]	
Full time	20 (26)
Part time	2 (3)
Unemployed	3 (4)
Retired	50 (66)
Disability	1 (1)
<b>Relationship characteristics</b>	
Caregiver relationship [n(%)]	

Spouse	26 (35)
Friend	5 (7)
Offspring	30 (39)
In-law	15 (20)
Frequency of caregiver contact [n(%)]	
Daily	48 (65)
Weekly to Daily	2 (3)
Weekly	9 (12)
Monthly	15 (20)
Caregiver's own health Good to Excellent (vs Fair or Poor) [n(%)]	49 (64)
Caregiver knows older adults's health Well to Very Well (vs Pretty well to Not at all) [n(%)]	64 (85)
Caregiver Burden (Zarit Burden Interview) (mean $\pm$ SD)	
At 18 months post-discharge (mean $\pm$ SD) (n=44)	13.8 $\pm$ 10.3
At 24 months post-discharge (mean $\pm$ SD) (n=38)	10.2 $\pm$ 9.9