

Supplementary Table 1. Bivariate comparison of demographic and health characteristics between respondents with no experience of migraine (non-migraine), patients diagnosed with migraine and matched non-migraine respondents.

		Non-migraine (N=25,209)		Matched Non-migraine (N=1,265)		Diagnosed migraine (N=1,265)		p-value	p-value
		Mean or %	[SD] or (n)	Mean or %	[SD] or (n)	Mean or %	[SD] or (n)	Non-migraine vs. Diagnosed	Matched vs. Diagnosed
Age, mean [SD]		52.80	[16.53]	43.45	[14.59]	43.79	[14.27]	< .001	.559
CCI, mean [SD]		0.17	[0.53]	0.18	[1.05]	0.24	[1.00]	< .001	.117
Gender, % (n)	Female	47.1%	(11,861)	67.4%	(852)	66.6%	(842)	< .001	.673
Employment status, % (n)	Currently employed	54.7%	(13,801)	60.1%	(760)	59.8%	(756)	< .001	.871
Household income, % (n)	<¥3,000,000	18.3%	(4623)	17.9%	(227)	18.7%	(237)	.683	.948
	¥3,000,000 to <¥5,000,000	24.4%	(6154)	24.5%	(310)	25.0%	(316)		
	¥5,000,000 to <¥8,000,000	23.9%	(6027)	25.9%	(328)	24.6%	(311)		
	¥8,000,000 or more	18.7%	(4704)	16.9%	(214)	17.0%	(215)		
	decline to answer	14.7%	(3701)	14.7%	(186)	14.7%	(186)		
Currently smoking, % (n)	Yes	41.2%	(10,385)	44.3%	(561)	44.0%	(557)	.046	.873
Currently use alcohol, % (n)	Yes	66.2%	(16,685)	63.0%	(797)	63.2%	(800)	.031	.902

Abbreviations: CCI = Charlson Comorbidity Index