

Supplementary Table 2. Unadjusted bivariate comparison of health outcomes between migraine patients currently taking Rx and migraine patients not currently taking Rx

Outcomes	Non-Rx (N=678)		Rx (N=587)		p-values
	Mean	SD	Mean	SD	
HRQoL					
SF-12v2: PCS	48.70	12.42	47.96	12.16	.290
SF-12v2: MCS	45.26	10.13	43.95	10.24	.022
SF-12v2: RCS	41.93	14.40	41.61	15.40	.703
EQ-5D index score	.77	.16	.76	.16	.255
EQ-5D VAS	65.33	23.11	63.35	23.95	.134
WPAI					
Absenteeism %*	7.10	17.71	6.76	16.98	.795
Presenteeism %**	31.01	26.60	34.75	27.52	.063
Work Productivity Loss %***	33.27	28.78	36.68	28.84	.119
Daily Activity Impairment %	34.07	27.52	37.58	28.28	.026
HCRU					
Number of physician visits in the past 6 months	7.52	10.05	9.37	16.16	.013
Number of ER visits in the past 6 months	.13	.62	.10	.71	.515
Number of times hospitalized in the past 6 months	.98	4.71	1.11	8.85	.738

Abbreviations: Rx = prescription medication, SD = standard deviation, HRQoL = Health-related quality of life, SF-12v2 = Short Form 12 health survey version 2, PCS = physical component summary score, MCS = mental component summary score, RCS = role/social component summary score, EQ-5D = EuroQol 5-Dimension, WPAI = Work Productivity and Activity Impairment, HCRU = healthcare resource utilization, ER = emergency room

*Sample size of Non-Rx: N=385, Rx: N=320. **Sample size of Non-Rx: N=395, Rx: N=337. ***Sample size of Non-Rx: N=380 Rx: N=317.