

Supplementary materials for

“Tumor-targeted delivery of siRNA to silence Sox2 gene expression enhances therapeutic response in hepatocellular carcinoma”

Yu Xia^{a,*}, Guoyi Tang^c, Yi Chen^b, Changbing Wang^b, Min Guo^b, Tiantian Xu^b, Mingqi Zhao^b, Yongjian Zhou^{a,*}

^a Department of Gastroenterology and Hepatology, Guangzhou Digestive Disease Center, Guangzhou First People's Hospital, School of Medicine, South China University of Technology, Guangzhou, 510180, China

^b Central Laboratory, Guangzhou Institute of Pediatrics, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, 510120, China

^c Department of Obstetrics Gynecology, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, 510120, China

* Corresponding authors. Department of Gastroenterology and Hepatology, Guangzhou Digestive Disease Center, Guangzhou First People's Hospital, School of Medicine, South China University of Technology, Guangzhou, 510180, China

E-mail addresses: eyyuxia@scut.edu.cn (Y. Xia), eyzhouyongjian@scut.edu.cn (Y. Zhou).




Fig. S1. Agarose gel electrophoretogram showing siSox2 protection by RGDfC-Se for 1 and 2 h incubation compared to siSox2 exposed to 50% serum. Naked siSox2 not exposed to serum was used as control.




Fig. S2. (A) The uptake of RGDfC-Se@siSox2 in HepG2 cells was observed under a fluorescence microscope. (B) The uptake of RGDfC-Se@siSox2 in HepG2 cells pretreated with RGDfC were observed under a fluorescence microscope. Scale bars = 20 μ m.