

**On-line Table 1: Subgroup comparison of functional dependence versus independence (mRS ≤2 at 3 months), unsuccessful versus successful recanalization, and symptomatic versus asymptomatic intracerebral hemorrhage**

	Functional Dependence	Functional Independence	Functional Dependence Value	Functional Independence Value	Unsuccessful Recanalization	Successful Recanalization	P Value	sICH	No sICH	P Value
<b>Patient characteristics</b>										
Age (mean) (yr)	65.9 ± 11.5	61.4 ± 14.4	.002	64 ± 11.6	64.9 ± 13.2	.64	65 ± 13.1	65.5 ± 12.3	.7	
Left-hemisphere stroke (No./No.) (%)	28/51 (54.9%)	23/33 (69.7%)	.17	16/37 (43.2%)	40/58 (69%)	.01	16/26 (61.5%)	39/68 (57.4%)	.7	
Baseline NIHSS (mean)	18 ± 4.7	16.1 ± 5.4	.9	17.8 ± 4.2	16 ± 6.1	.43	18.3 ± 4.7	16.2 ± 5.7	.09	
Baseline ASPECTS (median) (range)	7 ± 2	9 ± 1.6	.12	7 ± 2	8 ± 1.9	.02	6.5 ± 2.2	8 ± 1.8	.02	
Intravenous thrombolysis (No./No.) (%)	24/51 (47.1%)	17/33 (51.5%)	.69	18/37 (48.6%)	27/58 (46.6%)	.8	13/26 (50%)	31/68 (45.6%)	.13	
Prestroke antiplatelet therapy (No./No.) (%)	9/51 (17.6%)	8/33 (24.2%)	.46	9/37 (24.3%)	12/58 (20.7%)	.68	3/26 (11.5%)	17/68 (25%)	.15	
<b>Procedural characteristics</b>										
Onset to groin time (min) (median)	270 ± 137.3	233 ± 186.3	.78	275 ± 141.2	236 ± 166.5	.01	277 ± 180	245 ± 148	.45	
Onset to recanalization (min) (median)	360 ± 141.7	317.5 ± 210	.07	362.5 ± 152.8	330 ± 177.9	.63	353 ± 192.9	350 ± 158.7	.84	
Procedural time (median) (min)	77.5 ± 62.5	84 ± 54	.3	75 ± 71.9	81 ± 47.9	.68	100.5 ± 58.1	75 ± 58.4	.47	
No. of MT attempts (median)	3 ± 2	2 ± 1.2	.26	3 ± 2	2 ± 1.4	.12	3 ± 2.2	2 ± 1.5	.39	
Intraprocedural heparin (No./No.) (%)	27/51 (52.9%)	12/33 (36.4%)	.14	11/37 (29.7%)	29/58 (50%)	.052	14/26 (53.8%)	25/68 (36.8%)	.13	
Intraprocedural single-antiplatelet therapy (No./No.) (%)	22/43 (51.2%)	16/27 (59.3%)	.51	16/30 (53.3%)	30/49 (61.2%)	.49	11/22 (50%)	35/56 (62.5%)	.31	
Intraprocedural dual-antiplatelet therapy (No./No.) (%)	5/26 (19.2%)	4/15 (26.7%)	.59	4/18 (22.2%)	6/25 (24%)	.89	3/14 (21.4%)	7/28 (25%)	.8	
Successful recanalization mTICI ≥2b (No./No.) (%)	27/51 (52.9%)	25/33 (75.8%)	.036	/	/	/	17/26 (65.4%)	40/68 (58.8%)	.56	
Atheromatous lesion (No./No.) (%)	28/46 (60.9%)	23/33 (69.7%)	.4	25/34 (73.5%)	33/55 (60%)	.45	13/23 (56.5%)	45/65 (69.2%)	.027	
<b>Postprocedural characteristics</b>										
Postprocedural single-antiplatelet therapy (No./No.) (%)	17/36 (47.2%)	3/12 (25%)	.18	7/19 (36.8%)	18/35 (51.4%)	.31	8/19 (42.1%)	17/34 (50%)	.58	
Postprocedural dual-antiplatelet therapy (No./No.) (%)	7/27 (25.9%)	19/28 (67.9%)	.001	12/25 (48%)	18/35 (51.4%)	.79	4/16 (25%)	26/43 (60.5%)	.015	
sICH (No./No.) (%)	21/50 (42%)	3/33 (9.1%)	.001	9/37 (24.3%)	17/57 (29.8%)	.56	—	—	—	
Early stent patency (No./No.) (%)	25/31 (80.6%)	20/24 (83.3%)	.8	19/28 (67.9%)	31/36 (86.1%)	.08	17/18 (94.4%)	33/46 (71.7%)	.049	

**Note:**— indicates no need; mTICI, modified TICI.

**On-line Table 2: Univariate logistic regression analyses with ORs for functional independence, successful recanalization, stent patency, and symptomatic intracerebral hemorrhage**

	Functional Independence		Successful Recanalization		Stent Patency		sICH	
	OR (95% CI)	P Value	OR (95% CI)	P Value	OR (95% CI)	P Value	OR (95% CI)	P Value
Patient characteristics								
Age	0.97 (0.94-1)	.11	1 (0.97-1.03)	.73	0.96 (0.32-1)	.23	0.98 (0.95-1.02)	.5
Left-hemisphere stroke	1.89 (0.75-4.76)	.18	2.91 (1.23-6.86)	.01	2.12 (0.63-7.08)	.22	1.19 (0.4-3)	.8
Baseline NIHSS	0.92 (0.84-1)	.12	0.94 (0.86-1.02)	.15	0.97 (0.86-1.09)	.67	1.07 (0.97-1.18)	.12
Baseline ASPECTS	1.53 (1.16-2)	.002	1.13 (0.91-1.39)	.25	0.94 (0.7-1.26)	.68	0.72 (0.56-0.92)	.009
Intravenous thrombolysis	1.19 (0.49-2.87)	.69	0.91 (0.4-2.09)	.84	0.58 (0.17-1.95)	.38	1.19 (0.48-2.95)	.7
Prestroke antiplatelet therapy	1.49 (0.51-4.36)	.46	0.81 (0.3-2.17)	.67	0.69 (0.15-3.07)	.63	0.39 (0.1-1.46)	.15
Procedural characteristics								
Onset-to-groin time	1 (0.99-1)	.79	1 (0.99-1)	.75	1 (0.99-1)	.44	1 (0.99-1)	.63
Onset-to-recanalization	1 (0.99-1)	.81	1 (0.99-1)	.51	1 (0.99-1)	.84	1 (0.99-1)	.41
Procedural time	0.99 (0.99-1)	.66	0.99 (0.99-1)	.33	0.99 (0.99-1)	.13	1 (0.99-1)	.21
No. of MT attempts	0.71 (0.55-0.99)	.04	0.69 (0.51-0.93)	.01	0.71 (0.5-1)	.05	1.3 (0.98-1.75)	.06
Intraprocedural heparin	0.5 (0.2-1.24)	.18	2.3 (0.98-5.65)	.058	2.93 (0.81-10.61)	.13	2 (0.8-5)	.16
Intraprocedural single-antiplatelet therapy	1.38 (0.52-3.67)	.62	1.3 (0.55-3.46)	.64	0.51 (0.13-2)	.49	0.6 (0.22-1.62)	.32
Intraprocedural dual-antiplatelet therapy	1.52 (0.34-6.86)	.7	1.1 (0.26-4.67)	1	0.57 (0.08-3.9)	.61	0.81 (0.17-3.8)	1
Successful recanalization mTICI $\geq$ 2b	2.77 (1.05-7.31)	.04	/	/	2.93 (0.85-10.1)	.12	1.32 (0.51-3.38)	.64
Atheromatous lesion (OR) (95% CI)	1.47 (0.57-3.82)	.42	0.54 (0.21-1.37)	.19	0.75 (0.2-2.7)	.69	0.57 (0.21-1.5)	.27
Postprocedural characteristics								
sICH	0.13 (0.37-0.51)	.001	1.31 (0.51-3.38)	.64	6.69 (0.8-55.59)	.89	/	-
Postprocedural single-antiplatelet therapy	0.37 (0.08-1.6)	.31	1.81 (0.57-5.69)	.39	0.84 (0.16-4.43)	1	0.72 (0.23-2.25)	.77
Postprocedural dual-antiplatelet therapy	6.03 (1.87-19.44)	.003	1.14 (0.41-3.2)	1	0.73 (0.15-3.46)	.71	0.21 (0.06-0.78)	.02
Dual-layer stents	0.59 (0.1-3.25)	.69	2 (0.38-10.58)	.47	0.05 (0.005-0.5)	.007	-	.1

**Note:** --- indicates no need; OR, odds ratio; mTICI, modified TICI.