

Serum protein profiling reveals a landscape of inflammation and immune signaling in early-stage COVID-19 infection

Xin Hou^{1,3}, Xiaomei Zhang^{2,3}, Xian Wu^{1,3}, Minya Lu^{1,3}, Dan Wang², Meng Xu², Hongye Wang², Te Liang², Jiayu Dai², Hu Duan², Yingchun Xu^{1,4}, Xiaobo Yu^{2,4}, Yongzhe Li^{1,4}.

¹Department of Clinical Laboratory & Beijing Key Laboratory for Mechanisms Research and Precision Diagnosis of Invasive Fungal Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical Science & Peking Union Medical College, Beijing 100730, China.

²State Key Laboratory of Proteomics, Beijing Proteome Research Center, National Center for Protein Sciences-Beijing (PHOENIX Center), Beijing Institute of Lifeomics, Beijing, 102206, China.

³These authors contributed equally to this work.

⁴Correspondence to yongzhelipumch@126.com, yuxiaobo@mail.ncpsb.org and xycpumch@139.com.

Supplementary Tables

Supplementary Table 1	All COVID-19 and influenza patients enrolled in this study
Supplementary Table 2	List of proteins that are detected by antibody microarray
Supplementary	List of differentially-expressed serum proteins identified in

Table 3	COVID-19 patients compared to the influenza patients
Supplementary Table 4	Biological processes of differentially-expressed serum proteins identified in COVID-19 patients
Supplementary Table 5	Function enrichment analysis of differentially-expressed serum proteins identified in COVID-19 patients using the STRING database
Supplementary Table 6	Correlation analysis of differentially-expressed COVID-19 proteins and clinical indices.
Supplementary Table 7	Protein-protein interaction analysis of serum proteins correlated with the neutrophil and lymphocyte using STRING database.