



Comparison of dietary oils with different polyunsaturated fatty acid n-3 and n-6 content in the rat model of cutaneous wound healing

Tomas Komprda¹, Zbysek Sladek², Zuzana Sevcikova³, Veronika Svehlova¹, Jan Wijacki², Roman Guran^{4,5}, Tomas Do⁴, Zuzana Lackova^{4,5}, Hana Polanska⁶, Lucie Vrlikova³, Vendula Popelkova¹, Petr Michalek^{4,5}, Ondrej Zitka^{4,5}, Marcela Buchtova^{3*}

¹ Department of Food Technology, Mendel University in Brno, Zemedelska 1, 613 00 Brno, Czech Republic;

² Department of Animal Morphology, Physiology and Genetics, Mendel University in Brno, Zemedelska 1, 613 00 Brno, Czech Republic;

³ Laboratory of Molecular Morphogenesis, Institute of Animal Physiology and Genetics, v.v.i., Czech Academy of Sciences, Brno, Veвери 97, 602 00 Brno, Czech Republic

⁴ Department of Chemistry and Biochemistry, Mendel University in Brno, Zemedelska 1, 613 00 Brno, Czech Republic;

⁵ Central European Institute of Technology, Brno University of Technology, Purkynova 123, 612 00 Brno, Czech Republic;

⁶ Department of Pathological Physiology, Faculty of Medicine, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic

⁷ Section of Animal Physiology and Immunology, Department of Experimental Biology, Faculty of Science, Masaryk University, Brno 625 00, Czech Republic

* Correspondence: buchtova@iach.cz

Received: date; Accepted: date; Published: date

SUPPLEMENTARY FILES




Figure S1. Fatty acid content in individual diets

A, B: Total fatty acids content (EPA in red, DHA in orange) in used oils and individual diets: a standard diet alone (control, C), or a standard diet enriched with 8% of palm oil (P), safflower oil (S), fish oil (F) and *Schizochytrium* extract (Sch).




Figure S2: Weight characteristics in experimental animals

A: Final live weight of experimental animals. **B:** Daily weight gain of rats fed 64 days. SFA – saturated fatty acids, MUFA – monounsaturated fatty acids, PUFA – polyunsaturated fatty acids; stars labeled group of fatty acids differ significantly from the control group (* $p < 0.05$; ** $p < 0.01$, *** $p < 0.001$, unpaired t-test; $n = 10$).