## Transcriptome and genome sequencing elucidates the molecular basis for the high yield and good quality of the hybrid rice variety Chuanyou6203

Juansheng Ren<sup>1\*</sup>, Fan Zhang<sup>2\*</sup>, Fangyuan Gao<sup>1</sup>, Lihua Zeng<sup>3</sup>, Xianjun Lu<sup>1</sup>, Xiuqin Zhao<sup>2</sup>, Jianqun Lv<sup>1</sup>, Xiangwen Su<sup>1</sup>, Liping Liu<sup>1</sup>, Mingli Dai<sup>1</sup>, Jianlong Xu<sup>2</sup>, Guangjun Ren<sup>1\*\*</sup>

<sup>1</sup>Crop Research Institute, Sichuan Academy of Agricultural Sciences, Chengdu, 610066, P.R. China

<sup>2</sup>Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing, 100081, P.R. China

<sup>&</sup>lt;sup>3</sup>Sichuan Normal University, Chengdu, 610066, P.R. China

<sup>\*</sup>These authors contributed equally to the work

<sup>\*\*</sup>Corresponding author email: guangjun61@sina.com.




Figure S8. DEGs belonging to green-yellow modules enrich the pathway of carbon fixation in photosynthetic organisms by PATHVIEW Wed (https://pathview.uncc.edu/). 1 and 2 represent panicle samples 7 days before heading. 3 and 4 represent panicle samples 3 days after heading. 5 and 6 represent panicle samples 15 days after heading. 7 and 8 represent sword leaf samples 7 days before heading. 9 and 10 represent sword leaf samples 3 days after heading.