

Manuscript title:Population Structure of Nepali Spring Wheat (*Triticum aestivum* L.) Germplasm**Journal:**

BMC Plant Biology

Authors:*Kamal Khadka¹, Davoud Torkamaneh^{1,2,3}, Mina Kaviani¹, Francois Belzile^{2,3}, Manish N. Raizada¹ and Alireza Navabi¹**Affiliation:***¹ Department of Plant Agriculture, University of Guelph, Guelph, Ontario, Canada, N1G 2W1**Corresponding author email address:**kamal.khadka011@gmail.com

Additional file 4: Table S4. The frequency of genotypes in the Nepali Wheat Diversity Panel as differentiated into different subpopulations based on Q-matrix obtained from fastSTRUCTURE

	Subpopulation 1	Subpopulation 2	Subpopulation 3	Subpopulation 4	Total
Landraces	12 (0.07)	62 (0.37)	70 (0.42)	22 (0.13)	166 (1.00)
CIMMYT lines	8 (0.07)	69 (0.60)	25 (0.22)	13 (0.11)	115 (1.00)
Released varieties	2 (0.06)	22 (0.65)	4 (0.12)	6(0.18)	34 (1.00)
Canadian varieties	0	1 (0.33)	0	2 (0.67)	3 (1.00)

Note: values in parentheses indicate the frequency within each seed source category