



HHS Public Access

Author manuscript

Gastroenterology. Author manuscript; available in PMC 2021 March 01.

Published in final edited form as:

Gastroenterology. 2020 March ; 158(4): 1154. doi:10.1053/j.gastro.2020.02.014.

Spotlight: US Multi-Society Task Force on Colorectal Cancer Recommendations for Follow-up After Colonoscopy and Polypectomy

Samir Gupta, MD, MSCS¹, David Lieberman, MD², Joseph C. Anderson, MD^{3,4}, Carol A. Burke, MD⁵, Jason A. Dominitz, MD⁶, Tonya Kaltenbach, MD⁷, Douglas J. Robertson, MD³, Aasma Shaukat, MD⁸, Sapna Syngal, MD⁹, Douglas K. Rex, MD¹⁰


¹VA San Diego Healthcare System, San Diego, CA; UC San Diego Division of Gastroenterology and Moores Cancer Center, La Jolla, CA ²Division of Gastroenterology, Department of Medicine, Oregon Health and Science University, Portland, OR ³Veterans Affairs Medical Center, White River Junction, VT; The Geisel School of Medicine at Dartmouth, Hanover, NH ⁴University of Connecticut Health Center, Farmington, CT ⁵Department of Gastroenterology and Hepatology, Cleveland Clinic, Cleveland, OH ⁶VA Puget Sound Health Care System, University of Washington School of Medicine, Seattle, WA ⁷San Francisco Veterans Affairs Medical Center, San Francisco, CA; University of California San Francisco, San Francisco, CA ⁸Minneapolis VA Medical Center, Minneapolis, MN; University of Minnesota, Minneapolis, MN ⁹Division of Gastroenterology, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Division of Cancer Genetics and Prevention, Dana-Farber Cancer Institute ¹⁰Division of Gastroenterology and Hepatology, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



Recommendations for post-colonoscopy follow-up in average risk adults

Baseline finding	Recommended interval for first surveillance	Finding at first surveillance	Recommended interval for next surveillance
1–2 tubular adenomas < 10 mm	7–10 y	Normal colonoscopy*	10 y
		1–2 tubular adenomas < 10 mm	7–10 y
		3–4 tubular adenomas < 10 mm	3–5 y
		Adenoma ≥ 10 mm in size; or adenoma with tubulovillous/villous histology; or adenoma with high grade dysplasia; or 5–10 adenomas < 10 mm	3 y
3–4 tubular adenomas < 10 mm	3–5 y	Normal colonoscopy*	10 y
		1–2 tubular adenomas < 10 mm	7–10 y
		3–4 tubular adenomas < 10 mm	3–5 y
		Adenoma ≥ 10 mm in size; or adenoma with tubulovillous/villous histology; or adenoma with high grade dysplasia; or 5–10 adenomas < 10 mm	3 y
Adenoma ≥ 10 mm in size; or adenoma with tubulovillous/villous histology; or adenoma with high grade dysplasia; or 5–10 adenomas < 10 mm	3 y	Normal colonoscopy*	5 y
		1–2 tubular adenomas < 10 mm	5 y
		3–4 tubular adenomas < 10 mm	3–5 y
		Adenoma ≥ 10 mm in size; or adenoma with tubulovillous/villous histology; or adenoma with high grade dysplasia; or 5–10 adenomas < 10 mm	3 y

Recommendations for second surveillance stratified by adenoma findings at baseline and first surveillance

*Normal colonoscopy is defined as colonoscopy where no adenoma, SSR, or CRC is found.

Recommendations for post-colonoscopy follow-up in average risk adults with normal colonoscopy or adenomas ¹					Recommendations for post-colonoscopy follow-up in average risk adults with serrated polyps ¹					
Baseline finding	Risk compared with normal colonoscopy	Recommended interval for surveillance colonoscopy	Strength of recommendation	Quality of evidence	Baseline finding	Risk compared with normal colonoscopy	Recommended interval for surveillance colonoscopy	Strength of recommendation	Quality of evidence	
Normal	n/a	10 years ²	Strong	High	≤ 20 hyperplastic polyps in rectum or sigmoid colon < 10mm ⁷	↔	10 years ²	Strong	Moderate	
1 to 2 tubular adenomas < 10mm	↔	7 to 10 years ²	Strong	Moderate	≤ 20 hyperplastic polyps proximal to sigmoid colon < 10mm ⁷	↔	10 years	Weak	Very low	
3 to 4 tubular adenomas < 10mm	↑	3 to 5 years	Weak	Very low	1 to 2 SSPs < 10mm	↔ or ↑	5 to 10 years	Weak	Very low	
5 to 10 tubular adenomas < 10mm	↑↑	3 years	Strong	Moderate	3 to 4 SSPs < 10mm	↑	3 to 5 years	Weak	Very low	
> 10 adenomas on single exam ⁴	↑↑↑	1 year	Weak	Very low	5 to 10 SSPs < 10mm	↑↑	3 years	Weak	Very low	
Advanced adenoma	Any adenoma ≥ 10mm	↑↑↑↑	3 years	Strong	High	Hyperplastic polyp ≥ 10mm	↑↑	3 to 5 years ⁵	Weak	Very low
	Any adenoma with tubulovillous or villous histology	↑↑	3 years ⁵	Strong	Moderate	SSP ≥ 10mm	↑↑↑↑	3 years	Weak	Very low
	Any adenoma with high grade dysplasia	↑↑	3 years ⁵	Strong	Moderate	SSP with dysplasia	↑↑↑↑	3 years ⁵	Weak	Very low
	Piecemeal resection of adenoma ≥ 20mm	↑↑↑↑	6 months	Strong	Moderate ⁶	Traditional serrated adenoma	↑↑↑↑	3 years ⁵	Weak	Very low
						Piecemeal resection of SSP ≥ 20mm	↑↑↑↑	6 months	Strong	Moderate ⁶
					Large or advanced serrated polyp					

¹ All recommendations assume exam complete to cecum with bowel preparation adequate to detect lesions > 5mm in size; recommendations do not apply to individuals with a hereditary CRC syndrome, personal history of inflammatory bowel disease, personal history of hereditary cancer syndrome, serrated polyposis syndrome, malignant polyp, personal history of CRC, or family history of CRC, and must be judiciously applied to such individuals, favoring the shortest indicated interval based on either history or polyp findings.

² Follow up may be with colonoscopy or other screening modality for average risk individuals.

³ Patients with recommendations issued prior to 2020 for shorter than 7 to 10 year follow up after diagnosis of 1 to 2 tubular adenomas may follow original recommendations. If feasible, physicians may re-evaluate patients previously recommended an interval shorter than 10 years and reasonably choose to provide an updated recommendation for 7 to 10 year follow up, taking into account factors such as quality of baseline exam, prior polyp history, and patient preferences.

⁴ Patients with > 10 adenomas or lifetime > 10 cumulative adenomas may need to be considered for genetic testing based on absolute/cumulative adenoma number, patient age, and other factors such as family history of CRC (see text).

⁵ Assumes high confidence of complete resection.

⁶ See USMSTF recommendations for endoscopic removal of colorectal lesions (Kaltenbach TA et al., Endoscopic Removal of Colorectal Lesions—Recommendations by the US Multi-Society Task Force on Colorectal Cancer. At press 2020).

⁷ If cumulative more than 20 serrated polyps distributed throughout the colon, with at least 5 being proximal to the rectum, or 5 or more serrated polyps proximal to the rectum, all being 5mm or more in size, with 2 or more being 10mm or larger in size, consider a diagnosis of Serrated Polyposis Syndrome and alternate management (World Health Organisation. Classification of Tumours of the Digestive Tract. IARC Press, Lyon; 2019)

⁸ A 3-year follow up interval is favored if concern about consistency in distinction between SSP and HP locally, bowel preparation, or complete excision, whereas a 5-year interval is favored if low concerns for consistency in distinction between SSP and HP locally, adequate bowel preparation, and confident complete excision.

HP, hyperplastic polyp; SSP, sessile serrated adenoma/polyp/lesion