

Supplementary Data

SUPPLEMENTARY TABLE S1. DISTRIBUTION OF *SALMONELLA* ISOLATES ACCORDING TO SEROVAR AND REGION

<i>Province/region Serovar</i>	<i>Atlantic</i>	<i>British Columbia</i>	<i>Ontario</i>	<i>Prairies</i>	<i>Quebec</i>	<i>Total</i>
Kentucky	29	102	252	85	194	662
Heidelberg	57	17	113	14	116	317
Other ^a	42	26	62	42	41	213
Enteritidis	0	61	11	28	3	103
Infantis	2	4	6	29	4	45
Thompson	6	1	8	7	20	42
Typhimurium	1	0	4	24	12	41
Hadar	2	5	11	8	13	39
Schwarzengrund	0	1	4	25	3	33
Total	139	217	471	262	406	1495

^aThere were other serovars isolated, representing less than 2% of the total each.

SUPPLEMENTARY TABLE S2. DISTRIBUTION OF *SALMONELLA* ISOLATES ACCORDING TO SAMPLE TYPE AND STAGE IN THE PROCESSING CHAIN

	<i>Slaughter</i>	<i>Processing</i>	<i>Retail</i>	<i>Total</i>
Caeca	807	0	0	807
Parts	0	273	174	447
Whole carcasses	0	168	73	241
Total	807	441	247	1495