

SUPPLEMENTARY TABLES

Supplementary Table 1. Study design and animal usage.

Parts	Groups	BBB	Brain Ebema	WB/ RT-qPCR	IHC/IF	Death	Mortality (SAH)	Exclude (SAH)	Sum
Exp.1	sham	-	-	6	3	-		-	
	SAH3h,6h,12h,24h,48h,72h	-	-	36	3	7		3	
Exp.2	sham	6	6	6	3	-		-	
	SAH	6	6	6	3	2		-	
	SAH+AAV-NC	6	6	6	6	3		2	
Exp.3	SAH+AAV-Tim-3	6	6	6	6	9		1	
	sham	6	6	6	3	-	11.19%	-	
	SAH	6	6	6	3	-		-	365
	SAH+siRNA-scramble	6	6	6	6	2		1	
	SAH+Tim-3-siRNA	6	6	6	6	1		-	
Exp.4	sham	6	6	6	3	-		-	
	SAH	6	6	6	3	1		-	
	SAH+AAV-Tim-3+DMSO	6	6	6	6	11		2	
	SAH+AAV-Tim-3+NK-252	6	6	6	6	2		-	
	Subtotal	72	72	114	60	38		9	

Supplementary Table 2. Mortality of different groups.

Groups	Rats used	Mortality
sham	72	(0) 0
SAH	112	(10) 8.92%
SAH+AAV-NC	27	(3) 11.11%
SAH+AAV-Tim-3	33	(9) 27.27%
SAH+siRNA-scramble	26	(2) 7.69%
SAH+Tim-3-siRNA	25	(1) 4.00%
SAH+AAV-Tim-3+DMSO	35	(11) 16.12%
SAH+AAV-Tim-3+NK-252	26	(2) 7.69%