

Table S1. Scores on CERAD measures at the Time of Diagnosis of MCI, Probable Dementia and Either by Depressive Disorder Status Median (Min-Max)

CERAD							
Battery / Depressive Disorder	Possible Range	MCI (N=216)	P value	Probable Dementia (N=102)	P value	MCI/Probable dementia (N=285)	p-value¹
CERAD Word							
List-Delayed	0-10						
Yes		4 [0-8]	0.61	3 [0-7]	0.06	4 [0-8]	0.43
No		4 [0-10]		2 [0-7]		3 [0-10]	
CERAD							
Constructional Praxis-Recall	0-11						
Yes		9 [5-11]	0.04	8 [6-11]	0.33	9 [5-11]	0.09
No		9 [2-11]		9 [5-11]		9 [2-11]	
CERAD							

Category	unlimited						
Fluency							
Yes		11 [3-17]	0.02	9 [4-16]	0.86	11 [3-17]	0.05
No		12 [3-25]		10 [2-20]		12 [3-25]	
Boston							
Naming Test- 15 item							
Yes	0-15	13 [7-15]	0.88	12 [7-15]	0.09	13 [7-15]	0.68
No		13 [2-15]		13 [2-15]		13 [2-15]	
Trail Making							
Test-Part A							
Yes	0-300	67 [30-194]	0.08	77 [40-174]	0.85	70 [30-194]	0.20
No		56 [25-300]		78 [31-300]		60 [25-300]	
Trail Making							
Test-Part B							
Yes	0-300	151 [78-240]	0.26	174 [107-300]	0.41	158 [78-300]	0.34
No		152 [61-300]		218 [50-300]		179 [50-300]	

Notes: CERAD: The Consortium to Establish A Registry for Alzheimer's Disease; MCI: mild cognitive impairment; ¹ Wilcoxon test with significance level set to be 0.01 to control for multiple comparisons