

Supplementary

Table S1 Comparison of symptoms, physical activity, perceived health, and quality of life between the groups

Variable	Total (N=6,934)	Subjects without bronchiectasis (n=5,978)	Subjects with bronchiectasis (n=52)	P value
Symptoms				
Cough	5.0 (4.1–6.0)	4.9 (4.6–5.9)	21.0 (10.0–39.1)	<0.001
Sputum	9.4 (7.9–11.0)	9.3 (7.9–10.9)	24.2 (11.8–43.2)	0.008
Dyspnea	0.9 (0.6–13.7)	0.9 (0.7–1.3)	8.6 (2.0–30.5)	<0.001
Limitation of physical activity due to respiratory disease	5.2 (4.0–66.8)	4.8 (3.7–6.2)	34.8 (14.2–63.2)	<0.001
Perceived health				
Good/very good	41.4 (39.8–43.1)	41.5 (39.8–43.1)	28.3 (15.0–47.1)	0.061
Not good/not bad	39.4 (37.7–41.1)	39.4 (37.7–41.1)	37.3 (23.3–53.9)	
Bad/very bad	19.2 (18.1–20.3)	19.1 (17.9–20.2)	34.3 (21.3–50.2)	
EQ-5D component				
Mobility	12.9 (12.0–13.8)	12.8 (11.9–13.8)	22.7 (13.0–36.6)	0.040
Self-care	2.9 (2.5–3.3)	2.9 (2.5–3.3)	3.2 (0.6–15.7)	0.891
Usual activity	8.1 (7.4–8.8)	7.9 (7.3–8.7)	27.5 (15.8–43.4)	<0.001
Pain/discomfort	29.8 (28.3–31.4)	28.8 (27.2–30.5)	44.2 (28.1–61.6)	0.001
Anxiety/depression	12.6 (11.7–13.6)	12.5 (11.6–13.6)	28.2 (16.2–44.3)	0.004
EQ-5D index	0.92 (0.92–0.93)	0.93 (0.92–0.93)	0.87 (0.82–0.93)	<0.001

Data are presented as weighted mean (95% confidence interval) or weighted percentage (95% CI). EQ-5D, EuroQoL five dimension; CI, confidence interval.